(19) World Intellectual Property Organization International Bureau



- -

(43) International Publication Date 14 March 2002 (14.03.2002)

PCT

(10) International Publication Number WO 02/20569 A2

(51) International Patent Classification7:

C07K 14/00

- (21) International Application Number: PCT/US01/28013
- (22) International Filing Date:

7 September 2001 (07.09.2001)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/231,267 8 September 2000 (08.09.2000)

- (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).
- (72) Inventors: PARHAM, Christi, L.; 2385 30th Avenue, San Francisco, CA 94116 (US). GORMAN, Daniel, M.; 6371 Central Avenue, Newark, CA 94560 (US). KURATA, Hirokazu; 1091 Tanland Drive, #212, Palo Alto, CA 94303 (US). ARAI, Naoko; 648 Georgia Avenue, Palo Alto, CA 94306 (US). SANA, Theodore, R.; 1046 Pomeroy Avenue, Santa Clara, CA 95051 (US). MATTSON, Jeanine, D.; 559 Alvarado Street, San Francisco, CA 94114 (US). MURPHY, Erin, E.; 180 Emerson Street, Palo Alto, CA 94301 (US). SAVKOOR, Chetan; 4402 Silverberry Drive, San Jose, CA 95136-2415 (US). GREIN, Jeffery; 1083-A Alta Mira Drive, Santa Clara, CA 95051 (US). SMITH, Kathleen, M.; 275 Ventura #6, Palo Alto, CA 94304 (US). MCCLANAHAN, Terrill, K.; 1081 Westchester Drive, Sunnyvale, CA 94087 (US).

- (74) Agent: SCHRAM, David, B.; Schering Corporation, Patent Dept., K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN, YU, ZA.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

 as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Best Available Copy

(54) Title: MAMMALIAN GENES; RELATED REAGENTS AND METHODS

(57) Abstract: Nucleic acids encoding mammalian, e.g., primate or rodent, genes, purified proteins and fragments thereof. Anti-bodies, both polyclonal and monoclonal, are also provided. Methods of using the compositions for both diagnostic and therapeutic utilities are provided.

1

MAMMALIAN GENES; RELATED REAGENTS AND METHODS

FIELD OF THE INVENTION

The present invention relates to compositions and methods for affecting mammalian physiology, including morphogenesis or immune system function. In particular, it provides nucleic acids, proteins, and antibodies which regulate development and/or the immune system. Diagnostic and therapeutic uses of these materials are also disclosed.

5

10

15

20

25

30 /

BACKGROUND OF THE INVENTION

Recombinant DNA technology refers generally to techniques of integrating genetic information from a donor source into vectors for subsequent processing, such as through introduction into a host, whereby the transferred genetic information is copied and/or expressed in the new environment. Commonly, the genetic information exists in the form of complementary DNA (cDNA) derived from messenger RNA (mRNA) coding for a desired protein product. The carrier is frequently a plasmid having the capacity to incorporate cDNA for later replication in a host and, in some cases, actually to control expression of the cDNA and thereby direct synthesis of the encoded product in the host. See, e.g., Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols. 1-3, CSH Press, NY.

For some time, it has been known that the mammalian immune response is based on a series of complex cellular interactions, called the "immune network". Recent research has provided new insights into the inner workings of this network. While it remains clear that much of the immune response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines, play critical roles in controlling these cellular interactions. The interferons are generally considered to be members of the cytokine family. Thus, there is considerable interest in the isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which will lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system disorders.

Lymphokines apparently mediate cellular activities in a variety of ways. See, e.g., Paul (ed. 1998) <u>Fundamental Immunology</u> 4th ed., Lippincott; and Thomson (ed. 1998) <u>The</u>

2

Cytokine Handbook 3d ed., Academic Press, San Diego. They have been shown to support the proliferation, growth, and/or differentiation of pluripotential hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular lineages which make up a complex immune system. Proper and balanced interactions between the cellular components are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

5

10

15

20

25

30

Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These lymphocytes interact with many other cell types.

One means to modulate the effect of a cytokine upon binding to its receptor, and therefore potentially useful in treating inappropriate immune responses, e.g., autoimmune, inflammation, sepsis, and cancer situations, is to inhibit the receptor signal transduction. In order to characterize the structural properties of a cytokine receptor in greater detail and to understand the mechanism of action at the molecular level, purified receptor will be very useful. The receptors provided herein, by comparison to other receptors or by combining structural components, will provide further understanding of signal transduction induced by ligand binding.

An isolated receptor gene should provide means to generate an economical source of the receptor, allow expression of more receptors on a cell leading to increased assay sensitivity, promote characterization of various receptor subtypes and variants, and allow correlation of activity with receptor structures. Moreover, fragments of the receptor may be useful as agonists or antagonists of ligand binding. See, e.g., Harada, et al. (1992) J. Biol. Chem. 267:22752-22758. Often, there are at least two critical subunits in the functional receptor. See, e.g., Gonda and D'Andrea (1997) Blood 89:355-369; Presky, et al. (1996) Proc. Nat'l Acad. Sci. USA 93:14002-14007; Drachman and Kaushansky (1995) Curr. Opin. Hematol. 2:22-28; Theze (1994) Eur. Cytokine Netw. 5:353-368; and Lemmon and Schlessinger (1994) Trends Biochem. Sci. 19:459-463. Other receptor types, e.g., TLR-like, will similarly be useful.

3

Likewise, identification of novel ligands will be useful. Members of the tumor necrosis factor (TNF) family and transforming growth factor (TGF) family of ligands have identified physiological effects.

Finally, genes which exhibit disease associated expression patterns will be useful in diagnostic or other uses. The molecular diagnostic utility may be applied to identify patients who will be responsive to particular therapies, or to predict responsiveness to treatment.

5

10

15

20

25

30

From the foregoing, it is evident that the discovery and development of new soluble proteins and their receptors, including ones similar to lymphokines, should contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or indirectly involve development, differentiation, or function, e.g., of the immune system and/or hematopoietic cells. Moreover, novel markers will be useful in molecular diagnosis or therapeutic methods. In particular, the discovery and understanding of novel receptors or lymphokine-like molecules which enhance or potentiate the beneficial activities of other lymphokines would be highly advantageous. The present invention provides these and related compounds, and methods for their use.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1C show a sequence alignment of related IFN receptor family members. Tissue Factor is SEQ ID NO: 4; hIFNabR is SEQ ID NO: 5; CRF2-4 is SEQ ID NO: 6; cytor x is SEQ ID NO: 7; and cytor7 is SEQ ID NO: 8.

Figure 2 shows an alignment of TNF-x and TNF-y polypeptides (SEQ ID NO:9, 11, and 13); p is primate, r is rodent.

Figures 3A-3E show an alignment of primate and rodent TLR-like protein sequences. Figure 4 shows an Alignment of primate and rodent 5685C6 polypeptide sequences.

Figure 5 shows an alignment of Claudin homologs: D2 (SEQ ID NO:34); D8 (SEQ ID NO:37); D17 (SEQ ID NO:39); D7.2 (SEQ ID NO:41).

Figures 6A-6E show an alignment of Schlafen homologs: schlafen B (SEQ ID NO:43); schlafen C (SEQ ID NO:45); schlafen D (SEQ ID NO:47); schlafen E (SEQ ID NO:49); and schlafen F (SEQ ID NO:51).

4

SUMMARY OF THE INVENTION

The present invention is directed to novel genes, e.g., primate embodiments. These genes include receptors related to cytokine receptors, e.g., cytokine receptor like molecular structures, designated DNAX Interferon-like Receptor Subunit 4 (DIRS4); TNF related cytokines designated TNFx and TNFy; Toll-like receptor like molecules designated TLR-L1, TLR-L2, TLR-L3, TLR-L4, and TLR-L5; a TGF related molecule designated TGFx; a soluble Th2 cell produced entity designated 5685C6; a group of genes related to ones whose expression patterns correlate with medical conditions designated claudins, herein referred to as claudins D2, D8, D17, and D7.2; and a second group of genes related to ones whose expression patterns correlate with medical conditions designated schlafens, herein referred to as schlafens B, C, D, E, and F.

10

15

20

25

30

In particular, the present invention provides a composition of matter selected from: a substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of: SEQ ID NO: 2 (DIRS4); SEQ ID NO: 9, 11, 13, or 53 (TNFx or TNFy); SEQ ID NO: 15, 17, 19, 21, 23, 25, or 27 (TLR-L1 through TLR-L5); SEQ ID NO: 29 (TGFx): SEQ ID NO: 31 or 33 (5685C6); SEQ ID NO: 35, 37, 39, or 41 (claudins); SEQ ID NO: 43, 45, 47, 49, or 51 (schlafens). In preferred embodiments, the distinct nonoverlapping segments of identity: include one of at least eight amino acids; include one of at least four amino acids and a second of at least five amino acids; include at least three segments of at least four, five, and six amino acids; or include one of at least twelve amino acids. In certain embodiments, the polypeptide: is unglycosylated; is from a primate, such as a human; comprises at least contiguous seventeen amino acids of the SEO ID NO; exhibits at least four nonoverlapping segments of at least seven amino acids of the SEQ ID NO; has a length at least about 30 amino acids; has a molecular weight of at least 30 kD with natural glycosylation; is a synthetic polypeptide; is attached to a solid substrate; is conjugated to another chemical moiety; or comprises a detection or purification tag, including a FLAG, His6, or Ig sequence. In other embodiments, the composition comprises: a substantially pure polypeptide; a sterile polypeptide; or the polypeptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

5

Kit embodiments include those comprising such a polypeptide, and: a compartment comprising the polypeptide; or instructions for use or disposal of reagents in the kit.

Binding compound embodiments include those comprising an antigen binding site from an antibody, which specifically binds to a described polypeptide, wherein: the binding compound is in a container; the polypeptide is from a human; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to another chemical moiety; or the antibody: is raised to a recombinant polypeptide; is raised to a purified polypeptide; is immunoselected; is a polyclonal antibody; binds to a denatured antigen; exhibits a Kd to antigen of at least 30 _M; is attached to a solid substrate, including a bead or plastic membrane; is in a sterile composition; or is detectably labeled, including a radioactive or fluorescent label.

5

10

15

20

25

30

Kit embodiments include those comprising such a binding compound, and: a compartment comprising the binding compound; or instructions for use or disposal of reagents in the kit.

Methods are provided, e.g., for producing an antigen:antibody complex, comprising contacting under appropriate conditions a primate polypeptide with such a described antibody, thereby allowing the complex to form. Also provided are methods of producing an antigen:antibody complex, comprising contacting under appropriate conditions a polypeptide with an antibody which binds thereto, thereby allowing the complex to form. And methods are provided to produce a binding compound comprising: immunizing an immune system with a polypeptide described; introducing a nucleic acid encoding the described polypeptide to a cell under conditions leading to an immune response, thereby producing said binding compound; or selecting for a phage display library for those phage which bind to the desired polypeptide.

Further compositions are provided, e.g., comprising: a sterile binding compound, or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

Nucleic acid embodiments are provided, e.g., an isolated or recombinant nucleic acid encoding a polypeptide described, wherein the: polypeptide is from a primate; or the nucleic acid: encodes an antigenic polypeptide; encodes a plurality of antigenic polypeptide

6

sequences of SEQ ID NO:2, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, or 53; exhibits identity over at least thirteen nucleotides to a natural cDNA encoding the segment; is an expression vector; further comprises an origin of replication; is from a natural source; comprises a detectable label; comprises synthetic nucleotide sequence; is less than 6 kb, preferably less than 3 kb; is a hybridization probe for a gene encoding the polypeptide; or is a PCR primer, PCR product, or mutagenesis primer.

Various embodiments also include cells comprising the recombinant nucleic acids, particularly wherein the cell is: a prokaryotic cell; a eukaryotic cell; a bacterial cell; a yeast cell; an insect cell; a mammalian cell; a mouse cell; a primate cell; or a human cell.

5

10

15

20

25

30

Kit embodiments include those comprising a described nucleic acid, and: a compartment comprising the nucleic acid; a compartment further comprising a primate polypeptide; or instructions for use or disposal of reagents in the kit.

Other nucleic acids are provided which: hybridize under wash conditions of 30 minutes at 37° C and less than 2M salt to the coding portion of SEQ ID NO: 1, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50 or 52; or exhibit identity over a stretch of at least about 30 nucleotides to a SEQ ID NO: 1, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, or 52. Preferably, the wash conditions are at 45° C and/or 500 mM salt, or at 55° C and/or 150 mM salt; or the stretch is at least 55 or 75 nucleotides.

Methods are provided, e.g., for making: a duplex nucleic acid comprising contacting: a described nucleic acid with a complementary nucleic acid, under appropriate conditions, thereby resulting in hybridization to form the complex; or a nucleic acid complementary to a described nucleic acid with its complementary nucleic acid, under appropriate conditions, thereby resulting in hybridization to form the complex; or a polypeptide comprising culturing a cell comprising a described nucleic acid under conditions resulting in expression of the nucleic acid.

And methods are provided to: modulate physiology or development of a cell comprising contacting the cell with a polypeptide comprising SEQ ID NO: 9, 11, 13, 29, 31, or 33; modulate physiology or development of a cell comprising contacting the cell with a binding compound which binds to SEQ ID NO: 9, 11, 13, 29, 31, 33 or 53, thereby blocking signaling mediated by a protein comprising the SEQ ID NO; label a cell comprising contacting

7

the cell with a binding compound which binds to SEQ ID NO: 15, 17, 19, 21, 13, 15, or 37; or diagnose a medical condition comprising a step of evaluating expression of nucleic acid comprising SEQ ID NO: 34, 36, 38, 40, 42, 44, 46, 48, or 50.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. General

5

10

15

20

25

30

The present invention provides the amino acid sequences and nucleic acid sequences of mammalian, herein primate, genes. Among them is an interferon receptor-like subunit molecule, one designated DNAX Interferon Receptor family Subunit 4 (DIRS4), having particular defined properties, both structural and biological. Others include molecules designated TNFx and TNFy; Toll like receptor like molecules TLR-L1, TLR-L2, TLR-L3, TLR-L4, and TLR-L5; TGFx; 5685C6; claudins D2, D8, D17, and D7.2; and schlafens B, C, D, E, and F. Various cDNAs encoding these molecules were obtained from primate, e.g., human, cDNA sequence libraries. Other primate or other mammalian counterparts would also be desired. In certain cases, alternative splice variants should be available.

Some of the standard methods applicable are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols. 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York; each of which is incorporated herein by reference.

A nucleotide and corresponding amino acid sequence for a primate, e.g., human DIRS4 coding segment is shown in SEQ ID NO: 1 and 2, respectively. The new DIRS4 lacks a transmembrane segment, which suggests that the subunit acts as a soluble subunit, and would thus be an alpha receptor subunit. Alternatively, or in addition, a splice variant would exist which contains a transmembrane segment. This is consistent with the observation that two transcripts are found in many cell types. Interferon receptor like subunits may be receptors for the IL-10 family of ligands, e.g., IL-10, AK155, IL-19, IL-20/mda-7, AK155, IL-D110, IL-D210, etc. See, e.g., Derwent patent sequence database.

10

15

20

25

30

Also provided are nucleotide (SEQ ID NO: 8, 10, 12, and 52) and corresponding amino acid sequences (SEQ ID NO: 9, 11, 13, and 53) for primate and rodent forms of TNFx and primate and rodent forms of TNFy. Features for primate TNFx include: cAMP PKsites about 38, 74, 79, 205; Cas Phos sites about 41, 61; Cyt_c-Mesite about 43; Histone-Me site about 35; Myristoly sites about 5, 57, 220, 232 N-GLYCOSYL site about 229; PHOS2 sites about 38-41, 79-82, 134-136; PKC ph sites about 77, 142. Also segments 119-250, and 209-221 are notable. For rodent TNFx, features include: A predicted signal 1-19; mature would begin at about 20. Other features: cAMP PK sites at about 34, 93, 132, 229, 248, 263; Cas Phos sites about 119, 232, 251; Cyt_c-Me sites about 26, 90, 172; Histone-Me site about 82; Myristoly sites around 278, 290, 303; N-GLYCOSYL: 3 sites about 39, 287, 297; PHOS2 sites about 26-29, 34-37, 90-92, 93-96, 138-140, 192-194, 248-251; and PKC ph sites about 43, 51, 80, 81, 152; TyKinsite about 154. Signal cleavage site predicted between pos. 19 and 20: AGA-GA. Other significant segments include from about 74-132, 94-118, 168-308, and 193-201.

Nucleotide and corresponding amino acid sequences for TLR-L1 through TLR-L5 are provided in SEQ ID NO:14-27. The EST distribution for TLR1 suggests mRNA expression is restricted to brain tissue; chromosome Xq27.1-28 coding region is on a single exon. Features for primate TLR1 (SEQ ID NO:15) include: Tyr Kin site about 704 (KEGDPVAY); Tyr Kin sites about 713 (RNLQEFSY), 825(KPQSEPDY); N-GLYCOSYL sites about 84 (NYS), 219 (NCT), 294 (NPT), 366 (NIS), 421 (NLT), 583 (NLS); likely a Type Ia membrane protein; a possible uncleavable N-term signal sequence; and a transmembrane prediction of about 618-634 <612-646>. For rodent TLR-L1(SEQ ID NO:17), the features include: A predicted transmembrane segment from about residues 56-75; and predicted TyKin sites at about residues 136 and 145.

For primate TLR-L2 (SEQ ID NO:19) features include: N-glycosyl sites about 82 (NYT), 217 (NCS), 623 (NST), 674 (NQS); TyKin sites about 889 (RLREPVLY), 450 (RLSPELFY), 917 (KLNVEPDY); TyKin site about 889 (RLREPVLY), 917 (KLNVEPDY). Structurally this molecule has homology to type Ia membrane proteins.

Primate TLR-L3 (SEQ ID NO:23) has the following features: SIGNAL 1-26; TRANS 14-34; Pfam:LRRNT 43-73; Pfam:LRR 78-101; LRR_TYP 100-123; Pfam:LRR 102-125; LRR_TYP 124-147; Pfam:LRR 126-149; LRR_TYP 148-171; Pfam:LRR 150-173;

9

LRR_TYP 172-195; LRR_PS 172-194; Pfam:LRR 174-197; LRR_TYP 196-219; LRRCT 232-282; Pfam:LRRCT 232-282 with SEG 331-349 or SEG 365-379; Pfam:LRRNT 372-405; LRRNT 372-410; Pfam:LRR 409-432; LRR_TYP 431-454; Pfam:LRR 433-456; LRR_PS 455-477; LRR_TYP 455-478; Pfam:LRR 457-480; LRR_TYP 479-502; Pfam:LRR 481-504 with SEG 502-519; LRR_TYP 503-526; LRR_PS 503-525; Pfam:LRR 505-528; Pfam:LRRCT 562-612; LRRCT 562-612; TRANS 653-673; SEG 653-676; SEG 712-723; SEG 760-776; SEG 831-855. Structurally this molecule has homology to type Ia membrane proteins.

Primate TLR-L4 (SEQ ID NO:25) EST distributions suggest mRNA expression is restricted to brain tissue; human chromosome Xq26.3-28; predicted features at about, e.g., 10 SIGNAL 1-18; SEG 22-38; Pfam:LRR 60-83; LRR TYP 82-105; Pfam:LRR 84-107; LRR_PS 106-128; LRR TYP 106-129; Pfam:LRR 108-131; LRR TYP 130-153; Pfam:LRR 132-155; LRR SD22 154-174; LRR PS 154-176; LRR TYP 154-177; Pfam:LRR 156-178; LRR_SD22 177-198; LRR PS 177-198; LRR TYP 178-201; Pfam:LRR 179-200; Pfam:LRRCT 213-263; LRRCT 213-263; LRRNT 341-379; Pfam:LRRNT 341-374; Pfam:LRR 378-401; LRR_TYP 15 400-423; LRR SD22 400-421; Pfam:LRR 402-425; LRR_TYP 424-447; LRR_SD22 424-450; LRR_PS 424-447; Pfam:LRR 426-449; LRR_TYP 448-471; LRR_PS 448-470; Pfam:LRR 450-473; LRR TYP 472-495; LRR PS 472-494; Pfam:LRR 474-497; SEG 474-488; LRRCT 531-581; Pfam:LRRCT 531-581; SEG 617-643; TRANS 623-643; N-GLYCOSYL sites about 81 (NFS), 216 (NCS), 308 (NPS), 325 (NLS), 423 (NLT); 20 chromosome Xq26.3-28; coding region is on a single exon. Stucturally this molecule appears to be a Type Ia membrane protein.

For primate TLR-L5 (SEQ ID NO:27) the entire coding region lies on a single exon on human chromosome 13; predicted features at about, e.g., SIGNAL 1-20; Pfam:LRR 65-88; LRR_TYP 87-110; Pfam:LRR 89-112; LRR_TYP 111-134; Pfam:LRR 113-136; LRR_PS 135-157; LRR_SD22 135-156; LRR_TYP 135-158; Pfam:LRR 137-160; LRR_TYP 159-182; LRR_SD22 159-177; LRR_PS 159-181; Pfam:LRR 161-184; LRR_SD22 182-203; LRR_TYP 185-206; Pfam:LRR 185-205; LRRCT 218-268; Pfam:LRRCT 218-268; Hybrid:LRRNT 328-364; Pfam:LRRNT 328-360; LRR_SD22 386-407; Pfam:LRR 388-411; LRR_TYP 389-409; LRR_PS 410-432; LRR_TYP 410-433; LRR_SD22 410-428; Pfam:LRR 412-435; LRR_SD22 434-453; LRR_PS 434-457; LRR_TYP 434-457; Pfam:LRR 436-459; SEG 436-445; LRR_PS

25

30

10

458-480; LRR_SD22 458-484; LRR_TYP 458-481; SEG 459-476; Pfam:LRR 460-483; SEG 503-516; LRRCT 517-567; Pfam:LRRCT 517-567; SEG 585-596; TRANS 607-627; SEG 701-710; N-GLYCOSYL 3 sites about 292 (NDS), 409 (NLT), 572 (NPS); TyKin site about 798 (KLMETLMY).

5

10

15

20

25

30

Nucleotide and corresponding amino acid sequences for a primate, e.g., human, TGFx coding segment, are represented by SEQ ID NO:28 and 29, respectively. Human TGFx maps to chromosome 5 (clone CITB-H1_2319M24). Predicted features (SEQ ID NO: 29) include: TGFB domain 115-212; Pfam:TGF-beta 115-167; Pfam:TGF-beta 205-212; TGF-beta like conserved Cys residues at positions 115, 144, 148, 177, 209, 211.

Nucleotide and corresponding amino acid sequences for 5685C6 coding segments are presented in SEQ ID NO:30-33. The primate clone maps to chromosome 21q22.1. Features of primate 5685C6 (SEQ ID NO:31) include: N-GLYCOSYL sites about 10 (NST), 23 (NCS), 76 (NFT), 169 (NVT), 191 (NKS); most likely cleavage site predicted between pos. 19 and 20: VFA-LN. The secreted protein produced by Th2 cells. The corresponding rodent polypeptide (SEQ ID NO:33) has the following features Predicted features: N-GLYCOSYL sites about 6 (NNT), 19 (NCS), 159 (NRS); most likely cleavage site between pos. 26 and 27: TKA-QN. 5685C6 molecules appear to be soluble entities which are expressed in Th2 clones. The entities are useful markers of Th2 cells, and will be useful in characterizing such cell types.

Nucleotide and corresponding amino acid sequences for claudins D2, D8, D17, and D7.2 are SEQ ID NO:34-41 (See, e.g., Simon, et al. (1999) <u>Science</u> 285:103-106).

Nucleotide and corresponding amino acid sequences for schlafens B, C, D, E, and F (see, e.g., see Schwarz, et al. (1998) <u>Immunity</u> 9:657-668) are SEQ ID NO:42-51.

As used herein, the term DIRS4 shall be used to describe a protein comprising a protein or peptide segment having or sharing the amino acid sequence shown in the SEQ ID NOs noted above, or a substantial fragment thereof. The invention also includes a protein variation of the respective DIRS4 allele whose sequence is provided, e.g., a mutein or soluble extracellular construct. Typically, such agonists or antagonists will exhibit less than about 10% sequence differences, and thus will often have between 1- and 11-fold substitutions, e.g., 2-, 3-, 5-, 7-fold, and others. It also encompasses allelic and other variants, e.g., natural polymorphic, of the protein described. Typically, it will bind to its corresponding biological

11

ligand, perhaps in a dimerized state with a second receptor subunit, with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. The term shall also be used herein to refer to related naturally occurring forms, e.g., alleles, polymorphic variants, and metabolic variants of the mammalian protein.

5

10

15

20

25

30

Likewise, reference to the other genes described herein will be made. General descriptions directed to the methods of making or structural features will often be applicable to the other entities provided herein, e.g., the TNFx, TNFy, TLR-L1, TLR-L2, TLR-L3, TLR-L4, TLR-L5, TGFx, 5685C6, claudins D2, D8, D17, D7.2, and schlafens B, C, D, E, and F. Antibodies thereto, nucleic acids encoding them, etc., will be similarly applicable to the different entities.

This invention also encompasses proteins or peptides having substantial amino acid sequence identity with the amino acid sequences. It will include sequence variants with relatively few substitutions, e.g., preferably less than about 3-5.

A substantial polypeptide "fragment", or "segment", is a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids. Sequences of segments of different proteins can be compared to one another over appropriate length stretches.

Fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at residues 1, 2, 3, etc., and ending at, e.g., the carboxy-terminus (N), N-1, N-2, etc., in all practical combinations of different lengths. Particularly interesting polypeptides have one or both ends corresponding to structural domain or motif boundaries, as described, or of the designated lengths with one end adjacent one of the described boundaries. In nucleic acid embodiments, often segments which encode such polypeptides would be of particular interest.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches. In some comparisons, gaps may be introduces, as required. See, e.g.,

12

Needleham, et al. (1970) J. Mol. Biol. 48:443-453; Sankoff, et al. (1983) chapter one in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group (GCG), Madison, WI; each of which is incorporated herein by reference. This analysis is especially important when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are intended to include natural allelic and interspecies variations in the cytokine sequence. Typical homologous proteins or peptides will have from 50-100% homology (if gaps can be introduced), to 60-100% homology (if conservative substitutions are included) with an amino acid sequence segment of the appropriate SEQ ID NOs noted above. Homology measures will be at least about 70%, generally at least 76%, more generally at least 81%, often at least 85%, more often at least 88%, typically at least 90%, more typically at least 92%, usually at least 94%, more usually at least 95%, preferably at least 96%, and more preferably at least 97%, and in particularly preferred embodiments, at least 98% or more. The degree of homology will vary with the length of the compared segments. Homologous proteins or peptides, such as the allelic variants, will share most biological activities with the embodiments described individually, e.g., in the various tables.

10

15

20

25

30

As used herein, the term "biological activity" is used to describe, without limitation, effects on inflammatory responses, innate immunity, and/or morphogenic development by cytokine-like ligands. For example, the receptors typically should mediate phosphatase or phosphorylase activities, which activities are easily measured by standard procedures. See, e.g., Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature 363:736-738. The receptors, or portions thereof, may be useful as phosphate labeling enzymes to label general or specific substrates.

13

The terms ligand, agonist, antagonist, and analog of, e.g., a DIRS4_include molecules that modulate the characteristic cellular responses to cytokine ligand proteins, as well as molecules possessing the more standard structural binding competition features of ligand-receptor interactions, e.g., where the receptor is a natural receptor or an antibody. The cellular responses likely are typically mediated through receptor tyrosine kinase pathways.

Also, a ligand is a molecule which serves either as a natural ligand to which said receptor, or an analog thereof, binds, or a molecule which is a functional analog of the natural ligand. The functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists, see, e.g., Goodman, et al. (eds. 1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics, Pergamon Press, New York.

Rational drug design may also be based upon structural studies of the molecular shapes of a receptor or antibody and other effectors or ligands. See, e.g., Herz, et al. (1997) <u>J.</u>

Recept. Signal Transduct. Res. 17:671-776; and Chaiken, et al. (1996) <u>Trends Biotechnol.</u>

14:369-375. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) <u>Protein Crystallography</u>, Academic Press, New York, which is hereby incorporated herein by reference.

25 II. Activities

5

10

15

20

30

The cytokine receptor-like proteins will have a number of different biological activities, e.g., modulating cell proliferation, or in phosphate metabolism, being added to or removed from specific substrates, typically proteins. Such will generally result in modulation of an inflammatory function, other innate immunity response, or a morphological effect. The subunit will probably have a specific low affinity binding to the ligand.

Different receptors may mediate different signals. The TLR-L receptors may signal similar biology to the TLRs, which mediate fundamental innate immune or developmental responses. See, e.g., Aderem adn Ulevitch (2000) Nature 406:782-787. The TNFs and TGF are likely to signal as cytokines, as may the 5685C6, which seemingly is expressed by Th2 cells. The 5685C6 genes appear to be secreted proteins, which exhibit a cleavable signal sequence.

5

10

15

20

25

30

The claudins appear to be membrane proteins exhibiting 4 transmembrane segments, and seem to be associated with tight junctions and/or paracellular transport. They may also affect epithelial permeability or conductances, e.g., ion, across membranes. The claudin-D2 member of the claudin family is found to have regulated expression correlating with Crohn's disease. The other family members exhibit differential regulation in disease states, e.g., in Crohn's disease, ulcerative colitis, and various interstitial lung diseases. This is consistent with an important role in these disease processes. A functional role in the tight junctions/paracellular transport is consistent with problems in intestinal physiology.

Claudins define a structurally related multi-gene family of 4 TM proteins with distinct tissue distribution patterns. The claudins are major structural proteins of tight junctions (TJs) and can promote their formation. Their expression is necessary but not sufficient for tight junction formation. When expressed in fibroblasts, claudin-1 is capable of inducing a continuous association of adjacent cells, resulting in a cobblestone like pattern. However, this continuous barrier is not a tight junction. Claudins can be found outside of tight junction in certain cells. Claudin-3 and claudin-4 are receptors for Clostridium perfringens enterotoxin, a causative agent of fluid accumulation in the intestinal tract, causing diarrhea. Claudin-5 is deleted in Velo-cardio-facial syndrome (VCFS). Claudin-5 is only expressed in endothelial cells, and in some tissues it is even further restricted to arterials.

Mutations in Paracellin-1, claudin family member and a major renal tight junction protein, cause renal magnesium wasting with nephrocalcinosis. Thus, claudins may play important roles in selective paracellular conductance by determining the permeability of different epithelia.

The schlafens are members of a family of proteins of whose members are growth regulatory genes. See, e.g., Schwarz, et al. (1998) <u>Immunity</u> 9:657-668. These novel human sequences are related to the mouse Schlafen2 gene. It was observed to be differentially

15

regulated in mouse IBD: Rag Hh+ (IL-10 treated) colon expression was higher than Rag Hh+ alone and mimicked the expression of Rag Hh-.

The DIRS4 has the characteristic extracellular motifs of a receptor signaling through the JAK pathway. See, e.g., Ihle, et al. (1997) Stem Cells 15(suppl. 1):105-111; Silvennoinen, et al. (1997) APMIS 105:497-509; Levy (1997) Cytokine Growth Factor Review 8:81-90; Winston and Hunter (1996) Current Biol. 6:668-671; Barrett (1996) Baillieres Clin. Gastroenterol. 10:1-15; and Briscoe, et al. (1996) Philos. Trans. R. Soc. Lond. B. Biol. Sci. 351:167-171.

The biological activities of the cytokine or other receptor subunits will be related to addition or removal of phosphate moieties to substrates, typically in a specific manner, but occasionally in a non specific manner. Substrates may be identified, or conditions for enzymatic activity may be assayed by standard methods, e.g., as described in Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature 363:736-738.

III. Nucleic Acids

5

10

15

20

25

30

This invention contemplates use of isolated nucleic acid or fragments, e.g., which encode these or closely related proteins, or fragments thereof, e.g., to encode a corresponding polypeptide, preferably one which is biologically active. In addition, this invention covers isolated or recombinant DNAs which encode such proteins or polypeptides having characteristic sequences of the DIRS4 or the other genes. Typically, the nucleic acid is capable of hybridizing, under appropriate conditions, with a nucleic acid sequence segment shown in the appropriate SEQ ID NOs noted above, but preferably not with other genes. Said biologically active protein or polypeptide can be a full length protein, or fragment, and will typically have a segment of amino acid sequence highly homologous, e.g., exhibiting significant stretches of identity, to ones described. Further, this invention covers the use of isolated or recombinant nucleic acid, or fragments thereof, which encode proteins having fragments which are equivalent to the described proteins. The isolated nucleic acids can have

16

the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others from the natural gene.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially pure, e.g., separated from other components which naturally accompany a native sequence, such as ribosomes, polymerases, and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates, which are thereby distinguishable from naturally occurring compositions, and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule, either completely or substantially pure.

5

10

15

20

25

30

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain heterogeneity, preferably minor. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

A "recombinant" nucleic acid is typically defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence. Typically this intervention involves in vitro manipulation, although under certain circumstances it may involve more classical animal breeding techniques. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants as found in their natural state. Thus, for example, products made by transforming cells with an unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such a process is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a restriction enzyme sequence recognition site. Alternatively, the process is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms, e.g., encoding a fusion protein. Restriction enzyme recognition sites are often the target of such artificial

17

manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. This will include a dimeric repeat. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode equivalent polypeptides to fragments of the described sequences and fusions of sequences from various different related molecules, e.g., other cytokine receptor family members.

5

10

15

20

25

30

A "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least 21 nucleotides, more generally at least 25 nucleotides, ordinarily at least 30 nucleotides, more ordinarily at least 35 nucleotides, often at least 39 nucleotides, more often at least 45 nucleotides, typically at least 50 nucleotides, more typically at least 55 nucleotides, usually at least 60 nucleotides, more usually at least 66 nucleotides, preferably at least 72 nucleotides, more preferably at least 79 nucleotides, and in particularly preferred embodiments will be at least 85 or more nucleotides. Typically, fragments of different genetic sequences can be compared to one another over appropriate length stretches, particularly defined segments such as the domains described below.

A nucleic acid which codes for, e.g., a DIRS4, will be particularly useful to identify genes, mRNA, and cDNA species which code for itself or closely related proteins, as well as DNAs which code for polymorphic, allelic, or other genetic variants, e.g., from different individuals or related species. Other genes will be useful as markers for particular cell types, or diagnostic of various physiological conditions. Preferred probes for such screens may, in certain circumstances, be those regions of the gene which are conserved between different polymorphic variants or which contain nucleotides which lack specificity, and will preferably be full length or nearly so. In other situations, polymorphic variant specific sequences will be more useful.

This invention further covers recombinant nucleic acid molecules and fragments having a nucleic acid sequence identical to or highly homologous to the isolated DNA set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. Alternatively, recombinant clones derived from the genomic sequences, e.g., containing introns, will be useful for transgenic studies, including, e.g., transgenic cells and organisms, and for gene therapy. See, e.g., Goodnow

18

(1992) "Transgenic Animals" in Roitt (ed.) Encyclopedia of Immunology Academic Press, San Diego, pp. 1502-1504; Travis (1992) Science 256:1392-1394; Kuhn, et al. (1991) Science 254:707-710; Capecchi (1989) Science 244:1288; Robertson (1987)(ed.) Teratocarcinomas and Embryonic Stem Cells: A Practical Approach IRL Press, Oxford; and Rosenberg (1992) J. Clinical Oncology 10:180-199. Operable association of heterologous promoters with natural gene sequences is also provided, as are vectors encoding, e.g., the DIRS4 with a receptor partner. See, e.g., Treco, et al. WO96/29411 or USSN 08/406,030.

5

10

15

20

25

30

Homologous, or highly identical, nucleic acid sequences, when compared to one another, e.g., DIRS4 sequences, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. Comparative hybridization conditions are described in greater detail below.

Substantial identity in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 60% of the nucleotides, generally at least 66%, ordinarily at least 71%, often at least 76%, more often at least 80%, usually at least 84%, more usually at least 88%, typically at least 91%, more typically at least about 93%, preferably at least about 95%, more preferably at least about 96 to 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides, including, e.g., segments encoding structural domains such as the segments described below. Alternatively, substantial identity will exist when the segments will hybridize under selective hybridization conditions, to a strand or its complement, typically using a described sequence. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, more typically at least about 65%, preferably at least about 75%, and more preferably at least about 90%. See, Kanehisa (1984) Nucl. Acids Res. 12:203-213, which is incorporated herein by reference. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, generally at least about 20 nucleotides, ordinarily at least about 24 nucleotides, usually at least about 28 nucleotides, typically at least about 32 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably at least about 75 to 100 or more

19

nucleotides. This includes, e.g., 125, 150, 175, 200, 225, 250, 275, 300, 400, 500, 700, 900, and other lengths.

5

10

15

20

25

30

Stringent conditions, in referring to homology in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters typically controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30° C, more usually in excess of about 37° C, typically in excess of about 45° C, more typically in excess of about 55° C, preferably in excess of about 65° C, and more preferably in excess of about 70° C. Stringent salt conditions will ordinarily be less than about 500 mM, usually less than about 400 mM, more usually less than about 300 mM, typically less than about 200 mM, preferably less than about 100 mM, and more preferably less than about 80 mM, even down to less than about 20 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370, which is hereby incorporated herein by reference.

The isolated DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode this protein or its derivatives. These modified sequences can be used to produce mutant proteins (muteins) or to enhance the expression of variant species. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant derivatives include predetermined or site-specific mutations of the protein or its fragments, including silent mutations using genetic code degeneracy. "Mutant DIRS4" as used herein encompasses a polypeptide otherwise falling within the homology definition of the DIRS4 as set forth above, but having an amino acid sequence which differs from that of other cytokine receptor-like proteins as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant DIRS4" encompasses a protein having substantial sequence identity with a protein of SEQ ID NO:2, and typically shares most of the biological activities or effects of the forms disclosed herein.

Although site specific mutation sites are predetermined, mutants need not be site specific. Mammalian DIRS4 mutagenesis can be achieved by making amino acid insertions or deletions in the gene, coupled with expression. Substitutions, deletions, insertions, or many

combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy- terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mammalian DIRS4 mutants can then be screened for the desired activity, providing some aspect of a structure-activity relationship. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and periodic Supplements).

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The phosphoramidite method described by Beaucage and Carruthers (1981) <u>Tetra.</u>

<u>Letts.</u> 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polymerase chain reaction (PCR) techniques can often be applied in mutagenesis.

Alternatively, mutagenesis primers are commonly used methods for generating defined mutations at predetermined sites. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (1995; eds.) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY.

Antisense and other technologies for blocking expression of these genes are also available. See, e.g., Misquitta and Paterson (1999) <u>Proc. Nat'l Acad. Sci. USA</u> 96:1451-1456.

IV. Proteins, Peptides

5

10

15

20

25

30

As described above, the present invention encompasses primate DIRS4, e.g., whose sequences are disclosed in SEQ ID NO:2, and described above. Allelic and other variants are also contemplated, including, e.g., fusion proteins combining portions of such sequences with others, including epitope tags and functional domains. Analogous methods and applications exist directed to the other genes described herein.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins. A heterologous fusion protein is a fusion of

21

proteins or segments which are naturally not normally fused in the same manner. Thus, e.g., the fusion product of a DIRS4 with another cytokine receptor is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties, e.g., sequence or antigenicity, derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences.

5

10

15

20

25

30

In addition, new constructs may be made from combining similar functional or structural domains from other related proteins, e.g., cytokine receptors or Toll-like receptor like genes, including species variants. For example, ligand-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992, each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of receptor-binding specificities. For example, the ligand binding domains from other related receptor molecules may be added or substituted for other domains of this or related proteins. The resulting protein will often have hybrid function and properties. For example, a fusion protein may include a targeting domain which may serve to provide sequestering of the fusion protein to a particular subcellular organelle.

Candidate fusion partners and sequences can be selected from various sequence data bases, e.g., GenBank, c/o IntelliGenetics, Mountain View, CA; and BCG, University of Wisconsin Biotechnology Computing Group, Madison, WI, which are each incorporated herein by reference.

The present invention particularly provides muteins which bind cytokine-like ligands, and/or which are affected in signal transduction. Structural alignment of human DIRS4 with other members of the cytokine receptor family show conserved features/residues. Alignment of the human DIRS4 sequence with other members of the cytokine receptor family indicates various structural and functionally shared features. See also, Bazan, et al. (1996) Nature 379:591; Lodi, et al. (1994) Science 263:1762-1766; Sayle and Milner-White (1995) TIBS 20:374-376; and Gronenberg, et al. (1991) Protein Engineering 4:263-269. Similarly, the other genes have related family members.

Substitutions with either mouse sequences or human sequences are particularly preferred. Conversely, conservative substitutions away from the ligand binding interaction

regions will probably preserve most signaling activities; and conservative substitutions away from the intracellular domains will probably preserve most ligand binding properties.

5

10

15

20

25

30

"Derivatives" of the various proteins include amino acid sequence mutants, glycosylation variants, metabolic derivatives, and covalent or aggregative conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in amino acid side chains or at the N- or C- termini, e.g., by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties, including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the proteins or fragments thereof with other proteins of polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the proteins and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different proteins, resulting in, for instance, a hybrid protein exhibiting binding specificity for multiple different cytokine ligands, or a receptor which may have broadened or weakened specificity of substrate effect. Likewise, heterologous fusions may be constructed which would exhibit a

combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a receptor, e.g., a ligand-binding segment, so that the presence or location of a desired ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609, which is hereby incorporated herein by reference. Other gene fusion partners include glutathione-S-transferase (GST), bacterial β-galactosidase, trpE, Protein A, β-lactamase, alpha amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

5

10

15

20

25

30

The phosphoramidite method described by Beaucage and Carruthers (1981) <u>Tetra.</u>

<u>Letts.</u> 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory, and Ausubel, et al. (eds. 1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York, which are each incorporated herein by reference. Techniques for synthesis of polypeptides are described, for example, in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; each of which is incorporated herein by reference. See also Dawson, et al. (1994) Science 266:776-779 for methods to make larger polypeptides.

This invention also contemplates the use of derivatives of these proteins other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into three

classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of a receptor or other binding molecule, e.g., an antibody. For example, a cytokine ligand can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated Sepharose, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of an cytokine receptor, antibodies, or other similar molecules. The ligand can also be labeled with a detectable group, for example radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays.

A polypeptide of this invention can be used as an immunogen for the production of antisera or antibodies. These may be specific, e.g., capable of detecting or distinguishing between other related family members or various fragments thereof. The purified proteins can be used to screen monoclonal antibodies or antigen-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, the term "antibodies" also encompasses antigen binding fragments of natural antibodies, e.g., Fab, Fab2, Fv, etc. The purified proteins can also be used as a reagent to detect antibodies generated in response to the presence of elevated levels of expression, or immunological disorders which lead to antibody production to the endogenous receptor. Additionally, fragments may also serve as immunogens to produce the antibodies of the present invention. For example, this invention contemplates antibodies having binding affinity to or being raised against the amino acid sequences provided, fragments thereof, or various homologous peptides. In particular, this invention contemplates antibodies having binding affinity to, or having been raised against, specific fragments which are predicted to be, or actually are, exposed at the exterior protein surfaces.

The blocking of physiological response to the receptor ligands may result from the inhibition of binding of the ligand to the receptor, likely through competitive inhibition.

Antibodies to ligands may be antagonists. Thus, in vitro assays of the present invention will often use antibodies or antigen binding segments of these antibodies, or fragments attached to

25

solid phase substrates. Assays will also allow for the diagnostic determination of the effects of mutations and modifications, e.g., which affect signaling or enzymatic function.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to the receptor or fragments compete with a test compound for binding to a ligand or other antibody. In this manner, the neutralizing antibodies or fragments can be used to detect the presence of a polypeptide which shares one or more binding sites to a receptor and can also be used to occupy binding sites on a receptor that might otherwise bind a ligand.

10 V. Making Nucleic Acids and Protein

1.5

20

25

30

DNA which encodes the protein or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. Natural sequences can be isolated using standard methods and the sequences provided herein. Other species counterparts can be identified by hybridization techniques, or by various PCR techniques, or combined with or by searching in sequence databases, e.g., GenBank.

This DNA can be expressed in a wide variety of host cells which can, in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified constructs; and for structure/function studies. Variants or fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The protein, or portions thereof, may be expressed as fusions with other proteins.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired receptor gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression

control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

5

10

15

20

25

30

The vectors of this invention include those which contain DNA which encodes a protein, as described, or a fragment thereof encoding a biologically active equivalent polypeptide. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for such a protein in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the receptor is inserted into the vector such that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the protein or its fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of the protein encoding portion or its fragments into the host DNA by recombination.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but all other forms of vectors which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual, Elsevier, N.Y., and Rodriguez, et al. (eds. 1988) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Buttersworth, Boston, which are incorporated herein by reference.

Transformed cells are cells, preferably mammalian, that have been transformed or transfected with receptor vectors constructed using recombinant DNA techniques.

27

Transformed host cells usually express the desired protein or its fragments, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to express the subject protein. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the receptor to accumulate in the cell membrane. The protein can be recovered, either from the culture or, in certain instances, from the culture medium.

5

10

15

20

25

30

For purposes of this invention, nucleic sequences are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., <u>E. coli</u> and <u>B. subtilis</u>. Lower eukaryotes include yeasts, e.g., <u>S. cerevisiae</u> and <u>Pichia</u>, and species of the genus <u>Dictyostelium</u>. Higher eukaryotes include established tissue culture cell lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, <u>E. coli</u> and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the receptor or its fragments include, but are not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in <u>Vectors: A Survey of Molecular Cloning Vectors and Their Uses</u>, (eds. Rodriguez and Denhardt), Buttersworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.

. 1

28

Lower eukaryotes, e.g., yeasts and <u>Dictyostelium</u>, may be transformed with DIRS4 sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, <u>Saccharomyces cerevisiae</u>. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the receptor or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEp-series); integrating types (such as the YIp-series), or mini-chromosomes (such as the YCp-series).

5

10

15

20

25

30

Higher eukaryotic tissue culture cells are normally the preferred host cells for expression of the functionally active interleukin protein. In principle, many higher eukaryotic tissue culture cell lines are workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred. Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo PolyA, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

For secreted proteins, an open reading frame usually encodes a polypeptide that consists of a mature or secreted product covalently linked at its N-terminus to a signal

29

peptide. The signal peptide is cleaved prior to secretion of the mature, or active, polypeptide. The cleavage site can be predicted with a high degree of accuracy from empirical rules, e.g., von-Heijne (1986) Nucleic Acids Research 14:4683-4690 and Nielsen, et al. (1997) Protein Eng. 10:1-12, and the precise amino acid composition of the signal peptide often does not appear to be critical to its function, e.g., Randall, et al. (1989) Science 243:1156-1159; Kaiser et al. (1987) Science 235:312-317.

It will often be desired to express these polypeptides in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, the gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable in prokaryote or other cells.

10

15

20

25

30

The source of protein can be a eukaryotic or prokaryotic host expressing recombinant gene, such as is described above. The source can also be a cell line such as mouse Swiss 3T3 fibroblasts, but other mammalian cell lines are also contemplated by this invention, with the preferred cell line being from the human species.

Now that the sequences are known, the primate protein, fragments, or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York; and Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York; all of each which are incorporated herein by reference. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes. Similar techniques can be used with partial polypeptide sequences.

The various proteins, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either by a

30

so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the coupling reaction typically must be protected to prevent coupling at an incorrect location.

5

10

15

20

25

30

If a solid phase synthesis is adopted, the C-terminal amino acid is bound to an insoluble carrier or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonylhydrazidated resins, and the like.

An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in <u>J. Am. Chem. Soc.</u> 85:2149-2156, which is incorporated herein by reference.

The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation, electrophoresis, various forms of chromatography, and the like. The proteins of this invention can be obtained in varying degrees of purity depending upon desired uses. Purification can be accomplished by use of the protein purification techniques disclosed herein, see below, or by the use of the antibodies herein described in methods of immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate cells, lysates of other cells expressing the receptor, or lysates or supernatants of cells producing the protein as a result of DNA techniques, see below.

Generally, the purified protein will be at least about 40% pure, ordinarily at least about 50% pure, usually at least about 60% pure, typically at least about 70% pure, more typically at least about 80% pure, preferable at least about 90% pure and more preferably at least about 95% pure, and in particular embodiments, 97%-99% or more. Purity will usually

31

be on a weight basis, but can also be on a molar basis. Different assays will be applied as appropriate.

VI. Antibodies

5

10

15

20

25

30

Antibodies can be raised to the various mammalian, e.g., primate DIRS4, proteins and fragments thereof, both in naturally occurring native forms and in their recombinant forms, the difference being that antibodies to the active receptor are more likely to recognize epitopes which are only present in the native conformations. Denatured antigen detection can also be useful in, e.g., Western analysis. Anti-idiotypic antibodies are also contemplated, which would be useful as agonists or antagonists of a natural receptor or an antibody.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the protein can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity. These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M, preferably at least about 10 μ M, and more preferably at least about 3 μ M or better.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent agonists or antagonists, e.g., that bind to the receptor and inhibit or simulate binding to ligand, or inhibit the ability of the receptor to elicit a biological response, e.g., act on its substrate. They also can be useful as non-neutralizing antibodies or for use as markers for detection or diagnosis, and can be coupled to toxins or radionuclides to bind producing cells. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker.

The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they might bind to the antigen without inhibiting, e.g., ligand or substrate binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying antigen. They may be

32

used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the respective protein.

Protein fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Mammalian cytokine receptors, cytokines, enzymes, marker proteins, and fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York; each of which are incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

5

10

15

20

25

30

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256: 495-497, which discusses one method of generating monoclonal antibodies. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve <u>in vitro</u> exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin

33

Repertoire in Phage Lambda," <u>Science</u> 246:1275-1281; and Ward, et al. (1989) <u>Nature</u> 341:544-546. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant or chimeric immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) <u>Nature Genetics</u> 15:146-156.

10

15

20

25

30

The antibodies of this invention can also be used for affinity chromatography in isolating the proteins or peptides. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified protein will be released. Conversely, the protein may be used to purify antibody by immunoselection.

The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

Antibodies raised against a protein will also be used to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the protein or cells which express the protein. They also will be useful as agonists or antagonists of a ligand, which may be competitive inhibitors or substitutes for naturally occurring ligands.

A target protein that specifically binds to or that is specifically immunoreactive with an antibody generated against it, such as an immunogen consisting of a described amino acid sequence, is typically determined in an immunoassay. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a protein of SEQ ID NO: 2. This antiserum is selected to have low crossreactivity against other cytokine receptor family members, e.g., IFN

1 34

receptor subunits, preferably from the same species, and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

5

10

15

20

25

30

In order to produce antisera for use in an immunoassay, the protein, e.g., of SEQ ID NO: 2, is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred strain of mice such as Balb/c, is immunized with the selected protein, typically using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, supra). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used an immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10⁴ or greater are selected and tested for their cross reactivity against other cytokine receptor family members, e.g., receptors aligned in Figure 1, using a competitive binding immunoassay such as the one described in Harlow and Lane, supra, at pages 570-573. Preferably at least two cytokine receptor family members are used in this determination. These cytokine receptor family members can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein.

Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, the protein of SEQ ID NO: 2 can be immobilized to a solid support. Proteins added to the assay compete with the binding of the antisera to the immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to selected other receptor subunits. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein. In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein

required is less than twice the amount of the protein of the selected protein or proteins that is required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

It is understood that these proteins are members of families of homologous proteins. For a particular gene product, such as the DIRS4, the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are allelic, non-allelic, or species variants. It is also understood that the terms include nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding the respective proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations typically will substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring DIRS4 protein. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring the appropriate effect, e.g., upon transfected lymphocytes. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for the cytokine receptor family as a whole. By aligning a protein optimally with the protein of the cytokine receptors and by using the conventional immunoassays described herein to determine immunoidentity, one can determine the protein compositions of the invention.

VII. Kits and quantitation

5

10

15

20

25

30

Both naturally occurring and recombinant forms of the molecules of this invention are particularly useful in kits and assay methods. For example, these methods would also be applied to screening for binding activity, e.g., ligands or receptors for these proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK automated workstation, Beckman Instruments, Palo Alto, California, and Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes means for testing binding by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays to screen for a ligand or agonist/antagonist homologous proteins can be greatly

36

facilitated by the availability of large amounts of purified, soluble cytokine receptors in an active state such as is provided by this invention. Alternatively, production of large amounts of ligand will be useful in screening for receptor. Markers will also be available in large amounts to generate specific reagents.

Purified protein, e.g., DIRS4, can be coated directly onto plates or otherwise presented for use in the ligand or antibody screening techniques. However, non-neutralizing antibodies to these proteins can be used as capture antibodies to immobilize the respective receptor on the solid phase, useful, e.g., in diagnostic uses.

5

10

15

20

25

30

This invention also contemplates use of, e.g., DIRS4, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of the protein or its ligand. Alternatively, or additionally, antibodies against the molecules may be incorporated into the kits and methods. Typically the kit will have a compartment containing either a peptide or gene segment or a reagent which recognizes one or the other. Typically, recognition reagents, in the case of peptide, would be a receptor or antibody, or in the case of a gene segment, would usually be a hybridization probe. Diagnostic applications will be useful for the markers, as described.

A preferred kit for determining the concentration of, e.g., DIRS4, in a sample would typically comprise a labeled compound, e.g., ligand or antibody, having known binding affinity for DIRS4, a source of DIRS4 (naturally occurring or recombinant) as a positive control, and a means for separating the bound from free labeled compound, for example a solid phase for immobilizing the DIRS4 in the test sample. Compartments containing reagents, and instructions, will normally be provided.

Antibodies, including antigen binding fragments, specific for mammalian claudins or schlafens or a peptide fragment, or receptor fragments are useful in diagnostic applications to detect the presence of elevated levels of protein and/or its fragments. Diagnostic assays may be homogeneous (without a separation step between free reagent and antibody-antigen complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA) and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to a

37

cytokine receptor or to a particular fragment thereof. These assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) <u>Antibodies: A Laboratory Manual</u>, CSH., and Coligan (ed. 1991 and periodic supplements) <u>Current Protocols In Immunology</u> Greene/Wiley, New York.

Anti-idiotypic antibodies may have similar use to serve as agonists or antagonists of cytokine receptors or ligands. These should be useful as therapeutic reagents under appropriate circumstances.

5

10

15

20

25

30

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled ligand is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent, and will contain instructions for proper use and disposal of reagents. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium having appropriate concentrations for performing the assay.

The aforementioned constituents of the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In many of these assays, a test compound, cytokine receptor, ligand, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ¹²⁵I, enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Both of the patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The cytokine receptor can be immobilized on various matrixes followed by washing. Suitable matrices include plastic such

38

as an ELISA plate, filters, and beads. Methods of immobilizing the receptor to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of antibody/antigen complex by any of several methods including those utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) <u>Clin. Chem.</u> 30(9):1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678, each of which is incorporated herein by reference.

5

10

15

20

25

.30

Methods for linking protein or fragments to various labels are well reported in the literature. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequences provided. These sequences can be used as probes for detecting levels of the respective genes or transcripts in patients suspected of having an immunological or other medical disorder. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly ³²P. However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex

4 39

can be detected. The use of probes to the novel anti-sense RNA may be carried out in conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) <u>Progress in Growth Factor Res.</u> 1:89-97.

10

15

20

25

30

5

VIII. Therapeutic Utility

This invention provides reagents with significant therapeutic value. See, e.g., Levitzki (1996) Curr. Opin. Cell Biol. 8:239-244. The cytokine receptors (naturally occurring or recombinant), fragments thereof, mutein receptors, and antibodies, along with compounds identified as having binding affinity to the receptors or antibodies, should be useful in the treatment of conditions exhibiting abnormal expression of the receptors of their ligands. Such abnormality will typically be manifested by immunological or other disorders. Additionally, this invention should provide therapeutic value in various diseases or disorders associated with abnormal expression or abnormal triggering of response to the ligand. The biology of interferons, IL-10, TNFs, and TGFs are well described. Conversely, the TLRs have also been the subject of much interest, and the described homologs described herein will also be of similar interest. Associations with significant medical conditions for the claudins and schlafens is described below.

Recombinant proteins, muteins, agonist or antagonist antibodies thereto, or antibodies can be purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, along with physiologically innocuous stabilizers and excipients. These combinations can be sterile, e.g., filtered, and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof which are not complement binding.

40

Ligand screening using receptor or fragments thereof can be performed to identify molecules having binding affinity to the receptors. Subsequent biological assays can then be utilized to determine if a putative ligand can provide competitive binding, which can block intrinsic stimulating activity. Receptor fragments can be used as a blocker or antagonist in that it blocks the activity of ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of ligand, e.g., inducing signaling. This invention further contemplates the therapeutic use of antibodies to cytokine receptors as antagonists.

5

10

15

20

25

30

Conversely, receptor screening for receptors for ligands can be performed. However, ligands can also be screened for function using biological assays, which are typically simple due to the soluble nature of the ligands.

The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, reagent physiological life, pharmacological life, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; each of which is hereby incorporated herein by reference. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 µM concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

41

Cytokines, receptors, fragments thereof, and antibodies or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to be 5 administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or 10 parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing 15 Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents, e.g., agonists or antagonists of other cytokine 20 receptor family members.

IX. Screening

25

30

Drug screening using DIRS4, TLR-L receptors, or fragments thereof can be performed to identify compounds having binding affinity to the receptor subunits, including isolation of associated components. See, e.g., Emory and Schlegel (1996) Cost-Effective Strategies for Automated and Accelerated High-Throughput Screening IBC, Inc., Southborough, MA. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is therefore a blocker or antagonist in that it blocks the activity of the ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of a cytokine ligand. This invention

42

further contemplates the therapeutic use of antibodies to the receptor as cytokine agonists or antagonists.

Conversely, for ligands, receptors may be screened. Orphan receptor subunits, or testing of known receptor subunits in known or novel pairings may be performed.

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing the DIRS4 or TLR-L receptors. Cells may be isolated which express a receptor in isolation from other functional receptors, or in combination with other specific subunits. Such cells, either in viable or fixed form, can be used for standard ligand/receptor binding assays. See also, Parce, et al. (1989) Science 246:243-247; and Owicki, et al. (1990) Proc. Nat'l Acad. Sci. USA 87:4007-4011, which describe sensitive methods to detect cellular responses. Competitive assays are particularly useful, where the cells (source of putative ligand) are contacted and incubated with a labeled receptor or antibody having known binding affinity to the ligand, such as 125_Iantibody, and a test sample whose binding affinity to the binding composition is being measured. The bound and free labeled binding compositions are then separated to assess the degree of ligand binding. The amount of test compound bound is inversely proportional to the amount of labeled receptor binding to the known source. Any one of numerous techniques can be used to separate bound from free ligand to assess the degree of ligand binding. This separation step could typically involve a procedure such as adhesion to filters followed by washing, adhesion to plastic followed by washing, or centrifugation of the cell membranes. Viable cells could also be used to screen for the effects of drugs on cytokine mediated functions, e.g., second messenger levels, i.e., Ca⁺⁺; cell proliferation; inositol phosphate pool changes; and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system. Calcium sensitive dyes will be useful for detecting Ca⁺⁺ levels, with a fluorimeter or a fluorescence cell sorting apparatus.

X. Ligands

5

10

15

20

25

30

The descriptions of the DIRS4 and TLR-L receptors herein provide means to identify ligands, as described above. Such ligand should bind specifically to the respective receptor with reasonably high affinity. Various constructs are made available which allow either labeling of the receptor to detect its ligand. For example, directly labeling cytokine receptor,

43

fusing onto it markers for secondary labeling, e.g., FLAG or other epitope tags, etc., will allow detection of receptor. This can be histological, as an affinity method for biochemical purification, or labeling or selection in an expression cloning approach. A two-hybrid selection system may also be applied making appropriate constructs with the available cytokine receptor sequences. See, e.g., Fields and Song (1989) Nature 340:245-246.

Generally, descriptions of cytokine receptors will be analogously applicable to individual specific embodiments directed to DIRS4 or TLR-L reagents and compositions. Conversely, soluble ligands, e.g., TNFs and TGFs, will be characterized for biological activity.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the inventions to the specific embodiments.

EXAMPLES

I. General Methods

5

10

15

20

25

30

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols. 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Coligan, et al. (ed. 1996) and periodic supplements, Current Protocols In Protein Science Greene/Wiley, New York; Deutscher (1990) "Guide to Protein Purification" in Methods in Enzymology, vol. 182, and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1989) Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering,

44

<u>Principle and Methods</u> 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) <u>QIAexpress:</u> The High Level Expression & Protein Purification System QUIAGEN, Inc., Chatsworth, CA.

Computer sequence analysis is performed, e.g., using available software programs, including those from the GCG (U. Wisconsin) and GenBank sources. Public sequence databases were also used, e.g., from GenBank and others.

Many techniques applicable to IL-10 or IL-12 receptors may be applied to the DIRS4 or other receptor subunits, as described, e.g., in USSN 08/110,683 (IL-10 receptor), which is incorporated herein by reference.

10 II. Computational Analysis

5

15

20

25

30

Human sequences were identified from genomic sequence database using, e.g., the BLAST server (Altschul, et al. (1994) Nature Genet. 6:119-129). Standard analysis programs may be used to evaluate structure, e.g., PHD (Rost and Sander (1994) Proteins 19:55-72) and DSC (King and Sternberg (1996) Protein Sci. 5:2298-2310). Standard comparison software includes, e.g., Altschul, et al. (1990) J. Mol. Biol. 215:403-10; Waterman (1995) Introduction to Computational Biology: Maps, Sequences, and Genomes Chapman & Hall; Lander and Waterman (eds. 1995) Calculating the Secrets of Life: Applications of the Mathematical Sciences in Molecular Biology National Academy Press; and Speed and Waterman (eds. 1996) Genetic Mapping and DNA Sequencing (Ima Volumes in Mathematics and Its Applications, Vol 81) Springer Verlag.

III. Cloning of full-length cDNAs; Chromosomal localization

PCR primers derived from the sequences are used to probe a human cDNA library. Full length cDNAs for primate, rodent, or other species DIRS4 are cloned, e.g., by DNA hybridization screening of _gt10 phage. PCR reactions are conducted using T. aquaticus Taqplus DNA polymerase (Stratagene) under appropriate conditions.

Chromosome spreads are prepared. In situ hybridization is performed on chromosome preparations obtained from phytohemagglutinin-stimulated human lymphocytes cultured for 72 h. 5-bromodeoxyuridine was added for the final seven hours of culture (60 g/ml of medium), to ensure a posthybridization chromosomal banding of good quality.

45

A PCR fragment, amplified with the help of primers, is cloned into an appropriate vector. The vector is labeled by nick-translation with ³H. The radiolabeled probe is hybridized to metaphase spreads at final concentration of 200 ng/ml of hybridization solution as described in Mattei, et al. (1985) <u>Hum. Genet.</u> 69:327-331.

After coating with nuclear track emulsion (KODAK NTB₂), slides are exposed. To avoid any slipping of silver grains during the banding procedure, chromosome spreads are first stained with buffered Giemsa solution and metaphase photographed. R-banding is then performed by the fluorochrome-photolysis-Giemsa (FPG) method and metaphases rephotographed before analysis. Alternatively, mapped sequence tags may be searched in a database.

Similar appropriate methods are used for other species.

IV. Localization of mRNA

5

10

15

20

25

30

Human multiple tissue (Cat # 1, 2) and cancer cell line blots (Cat # 7757-1), containing approximately 2 μg of poly(A)⁺ RNA per lane, are purchased from Clontech (Palo Alto, CA). Probes are radiolabeled with[α-32P] dATP, e.g., using the Amersham Rediprime random primer labeling kit (RPN1633). Prehybridization and hybridizations are performed at 65° C in 0.5 M Na₂HPO₄, 7% SDS, 0.5 M EDTA (pH 8.0). High stringency washes are conducted, e.g., at 65° C with two initial washes in 2 x SSC, 0.1% SDS for 40 min followed by a subsequent wash in 0.1 x SSC, 0.1% SDS for 20 min. Membranes are then exposed at -70° C to X-Ray film (Kodak) in the presence of intensifying screens. More detailed studies by cDNA library Southerns are performed with selected human DIRS4 clones to examine their expression in hemopoietic or other cell subsets.

Alternatively, two appropriate primers are selected, e.g., from the tables. RT-PCR is used on an appropriate mRNA sample selected for the presence of message to produce a cDNA, e.g., a sample which expresses the gene.

Full length clones may be isolated by hybridization of cDNA libraries from appropriate tissues pre-selected by PCR signal. Northern blots can be performed.

Message for genes encoding each gene will be assayed by appropriate technology, e.g., PCR, immunoassay, hybridization, or otherwise. Tissue and organ cDNA preparations are

46

available, e.g., from Clontech, Mountain View, CA. Identification of sources of natural expression are useful, as described. And the identification of functional receptor subunit pairings will allow for prediction of what cells express the combination of receptor subunits which will result in a physiological responsiveness to each of the cytokine ligands.

5

10

15

20

25

30

For mouse distribution, e.g., Southern Analysis can be performed: DNA (5 µg) from a primary amplified cDNA library was digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and transferred to a nylon membrane (Schleicher and Schuell, Keene, NH).

Samples for mouse mRNA isolation may include: resting mouse fibroblastic L cell line (C200); Braf:ER (Braf fusion to estrogen receptor) transfected cells, control (C201); T cells, TH1 polarized (Mel14 bright, CD4+ cells from spleen, polarized for 7 days with IFN-γ and anti IL-4; T200); T cells, TH2 polarized (Mel14 bright, CD4+ cells from spleen, polarized for 7 days with IL-4 and anti-IFN-γ; T201); T cells, highly TH1 polarized (see Openshaw, et al. (1995) J. Exp. Med. 182;1357-1367; activated with anti-CD3 for 2, 6, 16 h pooled; T202); T cells, highly TH2 polarized (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367; activated with anti-CD3 for 2, 6, 16 h pooled; T203); CD44- CD25+ pre T cells, sorted from thymus (T204); TH1 T cell clone D1.1, resting for 3 weeks after last stimulation with antigen (T205); TH1 T cell clone D1.1, 10 μg/ml ConA stimulated 15 h (T206); TH2 T cell clone CDC35, resting for 3 weeks after last stimulation with antigen (T207); TH2 T cell clone CDC35, 10 µg/ml ConA stimulated 15 h (T208); Mel14+ naive T cells from spleen, resting (T209); Mel14+ T cells, polarized to Th1 with IFN-y/IL-12/anti-IL-4 for 6, 12, 24 h pooled (T210); Mel14+ T cells, polarized to Th2 with IL-4/anti-IFN-γ for 6, 13, 24 h pooled (T211); unstimulated mature B cell leukemia cell line A20 (B200); unstimulated B cell line CH12 (B201); unstimulated large B cells from spleen (B202); B cells from total spleen, LPS activated (B203); metrizamide enriched dendritic cells from spleen, resting (D200); dendritic cells from bone marrow, resting (D201); monocyte cell line RAW 264.7 activated with LPS 4 h (M200); bone-marrow macrophages derived with GM and M-CSF (M201); macrophage cell line J774, resting (M202); macrophage cell line J774 + LPS + anti-IL-10 at 0.5, 1, 3, 6, 12 h pooled (M203); macrophage cell line J774 + LPS + IL-10 at 0.5, 1, 3, 5, 12 h pooled(M204); aerosol challenged mouse lung tissue, Th2 primers, aerosol OVA challenge 7, 14, 23 h pooled (see Garlisi, et al. (1995) Clinical Immunology and Immunopathology 75:75-83; X206);

5

10

15

20

25

30

Nippostrongulus-infected lung tissue (see Coffman, et al. (1989) Science 245:308-310; X200); total adult lung, normal (O200); total lung, rag-1 (see Schwarz, et al. (1993) Immunodeficiency 4:249-252; O205); IL-10 K.O. spleen (see Kuhn, et al. (1991) Cell 75:263-274; X201); total adult spleen, normal (O201); total spleen, rag-1 (O207); IL-10 K.O. Peyer's patches (O202); total Peyer's patches, normal (O210); IL-10 K.O. mesenteric lymph nodes (X203); total mesenteric lymph nodes, normal (O211); IL-10 K.O. colon (X203); total colon, normal (O212); NOD mouse pancreas (see Makino, et al. (1980) Jikken Dobutsu 29:1-13; X205); total thymus, rag-1 (O208); total kidney, rag-1 (O209); total heart, rag-1 (O202); total brain, rag-1 (O203); total testes, rag-1 (O204); total liver, rag-1 (O206); rat normal joint tissue (O300); and rat arthritic joint tissue (X300).

Samples for human mRNA isolation may include: peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); T cell, TH1 clone HY06, resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti IFN-γ, TH2 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13, Tc783.58, Tc782.69, resting (T118); T cell random γδ T cell clones, resting (T119); Splenocytes, resting (B100); Splenocytes, activated with anti-CD40 and IL-4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); NK 20 clones pooled, resting (K100); NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101); NKL clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); NK cytotoxic clone 640-A30-1, resting (K107); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101);

48

elutriated monocytes, activated with LPS, IFNy, anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFNy, IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFNy, anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFNy, IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated monocytes, activated LPS for 6 h (M109); DC 70% CD1a+, from CD34+ GM-CSF, TNFα 12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNFα 12 days, activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNFα 12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D104); DC 95% CD14+, ex CD34+ GM-CSF, TNFα 12 days FACS sorted, activated with PMA and ionomycin 1, 6 hr pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNFα, monocyte supe for 4, 16 h pooled (D110); leiomyoma L11 benign tumor (X101); normal myometrium M5 (O115); malignant leiomyosarcoma GS1 (X103); lung fibroblast sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102); kidney fetal 28 wk male (O100); lung fetal 28 wk male (O101); liver fetal 28 wk male (O102); heart fetal 28 wk male (O103); brain fetal 28 wk male (O104); gallbladder fetal 28 wk male (O106); small intestine fetal 28 wk male (O107); adipose tissue fetal 28 wk male (O108); ovary fetal 25 wk female (O109); uterus fetal 25 wk female (O110); testes fetal 28 wk male (O111); spleen fetal 28 wk male (O112); adult placenta 28 wk (O113); and tonsil inflamed, from 12 year old (X100).

10

15

20

25

30

For the DIRS4, southern blot analysis revealed expression in several cDNA libraries, including resting MOT72 (Th0 clone); resting, activated, and anti-peptide HY06 (Th1 clone); activated T cells CD4+, Th2 polarized; resting pooled T cell clones; resting and activated splenocytes; resting EBV B cells; activated JY (B cell line); cytotoxic NK cells; TF1 cells; resting and activated U937 cells; monocytes treated with anti-IL-10; monocytes (anti-IL-10 and IL-10 stimulated); activated monocytes; dendritic cells (activated and resting); MRC5

49

(lung fibroblast sarcoma line); CHA (kidney epithelial carcinoma line); normal and asthmatic monkey lung; normal and smoker lung; normal colon; fetal lung; liver; gall bladder; and small intestine. There were two transcript sizes, about 500 bp and about 1.8 kb bands, suggesting two different transcripts, possibly soluble and membrane spanning forms.

5

10

The primate, e.g., human, TNFx expression, by PCR, is high in allergic lung and normal lung; much lower in adult placenta, fetal spleen, and normal skin. Essentially no expression in gut samples and fetal organs. In cells, high expression was detected in resting HY06 cells and TF-1; lower in activated HY06 cell and JY cells, and no significant expression in the other human samples tested, e.g., most in the list above. Table 1 shows additional TaqMan expression data for human TNFx.

Table 1:

LIBRARY	Ct_gene	LIBRARY	Ct_gene
PBMC resting	44.64 mono + anti-IL-10		22.47
PBMC activated	40.48	mono + IL-10	21.04
Mot 72 resting	26.29	M1	40.52
Mot 72 activated	24.51	M6	21.75
Mot 72 anti-peptide	20.72	70% DC resting	26.27
HY06 resting	15.86	D1	37.94
HY06 activated	18.3	D6	25.05
HY06 anti-peptide	24.27	CD1a+ 95%	26.87
HY935 resting	25.97	CD14+ 95%	35.17
HY935 activated	25.03	CD1a+ CD86+	27.48
B21 resting	26.3	DC/GM/IL-4	32.33
B21 activated	24.53	DC LPS	27.81
Te gamma delta	45	DC mix	27.32
Jurkat resting pSPORT	45	fetal kidney	26.41
Jurkat activated pSPORT	28.09	fetal lung	31.16
Splenocytes resting	23.51	fetal liver	26.28
Splenocytes activated	26.19	fetal heart	34.28
Вс	23.88	fetal brain	25.02
JY	19.29	fetal small intestine	37.89
NK pool	38.21	fetal adipose tissue	26.41
NK pool activated	37.54	fetal ovary	37.49
NKA6 pSPORT	34.39	fetal uterus	26.03
NKL/IL-2	25.71	fetal testes	36.65
NK cytotox.	23.28	fetal spleen	23.2
NK non cytotox.	26.35	adult placenta	24.06
U937/CD004 resting	28.18	inflammed tonsil	26.21
U937 activated	26.21	TF1	23.48
C-	27	MRC5	33.99

LIBRARY	Ct_gene	LIBRARY	Ct_gene
C+	23.13	СНА	28.27
mast cell pME	28.65	Taq_control_genomic_2	50
TC1080 CD28- pMET7	38.1	Crohns colon 403242A	28.32
RV-C30 TR1 pMET7	24.97	lung 080698-2	27.42
DC resting mono-derived	28.12	18 hr. Ascaris lung	28.06
DC CD40L activ. mono-deriv.	27.07	hi dose IL-4 lung	34.01
DC resting CD34-derived	28.9	normal colon #22	44.6
DC TNF/TGFb act CD34-der.	36.74	ulcerative colitis colon #26	38.12
allergic lung #19	20.21	normal thyroid	28.14
Pneumocystis carnii lung #20	36.33	Hashimotos thyroiditis	36.88
RA synovium pool	28	normal skin	24.12
Psoriasis skin	32.37	Crohns colon 4003197A	30.31
normal lung	35.68	lung 121897-1	36.25
4 hr. Ascaris lung	31.45	Crohns colon 9609C144	27.49
24 hr. Ascaris lung	26.34	A549 unstim.	28.03
normal lung pool	22.21	A549 activated	24.1
Taq_control_genomic_1	50	Taq_control_water	50

The rodent, e.g., mouse, TNFx is highly expressed in 5 month ApoE KO mouse aorta; C57B6 3 wk polarized Th1 cells; and C57B6 3 wk polarized Th2 cells. It is less highly expressed in Balb/c 3 wk polarized Th2 cells, LPS treated spleen, and various other Th2 polarized populations. In tissues, by PCR, it is expressed highly in TNK KO spleen, NZB/W spleen, NZB/W spleen, GF ears/skin; rag-1 testis, w.t. C57B6 spleen, w.t. C57B6 pancreas, and 2 mo. lung. It is expressed at lower levels in influenza lung, rag-1 lung, rag-1 spleen, spinal cord samples, lung samples, stomach, and lymph nodes. Table 2 shows additional TaqMan expression data for mouse TNFx.

Table 2:

LIBRARY	Ct_gene	LIBRARY	Ct_gene
L cell	26	rag-1 brain	24.47
TH1 7 day	26.63	rag-1 testes	38.4
TH2 7 day	24.56	rag-1 lung	22.81
TH1 3 week Balb/C	39.09	rag-1 liver	36.69
TH2 3 week Balb/C	24.48	rag-1 spleen	24.23
preT	36.92	rag-1 thymus	23.91
D1.1 resting	32.74	rag-1 kidney	22.32
D1.1 con A stim.	37.76	w.t. Peyers patches	25.48
CDC35 resting	30.8	w.t. mesenteric lymph nodes	25.59
CDC35 con A stim.	41.92	w.t. colon	28.7
Mel 14+ naive T	28.16	Braf:ER (-) oligo dT	38.53
Mel14+ TH1	29.2	TH1 3 week C57 Bl/6	23.12
Mel 14+ TH2	25.02	TH2 3 week C57 Bl/6	22.54
A20	37.61	TH1 3 week Balb/C fresh	28.02
CH12	25.29	TH2 3 week Balb/C fresh	37.73
lg. B cell	30.34	b.m. DC (YJL) resting	27.99
LPS spleen	24.04	b.m. DC (YJL) aCD40 stim.	40.47
macrophage	28.6	b.m. $mf + LPS + aIL-10R$	29.74
J774 resting	39.73	b.m. mf + LPS + IL-10	27.67
J774 +LPS + anti-IL-10	36.51	peritoneal mf	37.02
J774 +LPS + IL-10	40.53	MC-9/MCP-12 pMET7	39.68
Nippo-infected lung	25.87	EC	40.13
IL-10 K.O. spleen	24.18	EC + TNFa	40.54
IL-10 K.O. colon	36.97	bEnd3 + TNFa	41.26
asthmatic lung	26.61	bEnd3 + TNFa + IL-10	38.35
w.t. lung	24.06	ApoE aorta 5 month	21.03
w.t. spleen	28.87	ApoE aorta 12 month	34.28
rag-1 heart	26.48	NZ B/W kidney	21.02

LIBRARY	Ct_gene	LIBRARY	Ct_gene
Nippo IL-4 K.O. lung	28.59	NZ B/W spleen	21.2
Nippo anti IL-5 lung	25.73	tolerized & challenged lung	27.17
Influenza lung	23.93	Aspergillus lung	23.32
b common lung 2 month	24.53	Taq_control_water	50
IL-10 K.O. stomach	29.87	Taq_control_genomic_1	50
IL-10 K.O. MLN aIL-12	26.58	Taq_control_genomic_2	50
IL-10 K.O. MLN +IL-10	25.89	w.t. d17 spinal cord EAE model	22.87
Rag-2 Hh- colon	29.2	TNF K.O. d17 spinal cord EAE	22.84
		model	
Rag-2 Hh+ colon	27.1	TNF K.O. spinal cord	23.27
IL-7 K.O./Rag-2 Hh- colon	40	TNF K.O. spleen	20.78
IL-7 K.O./Rag-2 Hh+ colon	40	G.F. ears (skin)	20.7
transfer model IBD	28.1	w.t. spinal cord	22.74
w.t. C57 Bl/6 aorta	39.38	8 w.t. C57 Bl/6 spleen	22.15
w.t. thymus	27.05	w.t. C57 Bl/6 pancreas	24.75
w.t. stomach	26.49	MM2/MM3 activated. pME	37.67
MM2/MM3 resting pME	37.62	2	

The primate, e.g., human, TNFy is expressed in fetal adipose tissue and fetal ovary. It is expressed at a lower level in fetal brain, Hashimoto's thyroiditis, RA synovium pool, adult placenta, and fetal uterus. It is expressed at lower levels in fetal kidney, normal thyroid, and detectable in Crohn's colon, psoriasis skin, and fetal lung. It is essentially undetectable in other organs evaluated, including various Ascaris challenged lung samples. In cell libraries, it is expressed in TF-1 cells, and much lower in CHA cells, and was not significantly expressed in other cell lines tested. Table 3 provides additional TaqMan expression data for human TNFy.

Table 3:

LIBRARY	Ct_gene	LIBRARY	Ct_gene
PBMC resting	45	mono + IL-10	42.96
PBMC activated	44.16	5M1	41.25
Mot 72 resting	42.47	⁷ M6	45
Mot 72 activated	28.59	70% DC resting	40.37
Mot 72 anti-peptide	42.47	7D1	28.94
HY06 resting	43.19	9D6	28.38
HY06 activated	41.48	3 CD1a+ 95%	25.63
HY06 anti-peptide	43.28	3 CD14+ 95%	28.36
HY935 resting	45	5 CD1a+ CD86+	28.67
HY935 activated	43.62	DC/GM/IL-4	45
B21 resting	41.73	BDC LPS	38.8
B21 activated	44.35	5 DC mix	26.53
Tc gamma delta	43.2	l fetal kidney	27.98
Jurkat resting pSPORT	23.44	fetal lung	30.57
Jurkat activated pSPORT	25.19	efetal liver	43.92
Splenocytes resting	38.72	2 fetal heart	40.84
Splenocytes activated	44.09	9 fetal brain	26.02
Bc	44.83	3 fetal small intestine	40.05
JY	43.0	5 fetal adipose tissue	23.63
NK pool	39.09	9 fetal ovary	25.85
NK pool activated	44.32	2 fetal uterus	27.57
NKA6 pSPORT	42.5	8 fetal testes	45
NKL/IL-2	4:	5 fetal spleen	39.08
NK cytotox.	44.79	9 adult placenta	28.05
NK non cytotox.	4:	5 inflammed tonsil	45
U937/CD004 resting	24.1	7 TF 1	22.09
U937 activated	24.4	1 MRC5	26.18
C-	40.3	8 CHA	19.22
C+	41.1	7 mast cell pME	43.93

. 55

LIBRARY	Ct_gene	LIBRARY	Ct_gene
mono + anti-IL-10	45	TC1080 CD28- pMET7	41.62
DC resting mono-derived	45	RV-C30 TR1 pMET7	42.76
DC CD40L activ. mono-deriv.	45	4 hr. Ascaris lung	45
DC resting CD34-derived	45	24 hr. Ascaris lung	45
DC TNF/TGFb act CD34-der.	39.71	normal lung pool	45
allergic lung #19	43.22	normal skin	42.69
Pneumocystis carnii lung #20	43.81	Crohns colon 4003197A	29.82
normal colon #22	43.66	lung 121897-1	45
ulcerative colitis colon #26	45	Crohns colon 9609C144	41.86
normal thyroid	27.71	A549 unstim.	27.09
Hashimotos thyroiditis	27.4	A549 activated	29.01
RA synovium pool	28	Taq_control_water	50
Psoriasis skin	31.49	Taq_control_genomic_1	50
normal lung	45	Taq_control_genomic_2	50
Crohns colon 403242A	33.18	18 hr. Ascaris lung	44.16
lung 080698-2	30.01	hi dose IL-4 lung	43.59

Table 4 provides TaqMan expression data for rodent, e.g., moust TNFy.

LIBRARY	Ct_gene	LIBRARY	Ct_gene
L cell	40 rag-1 lung		40
TH1 7 day	40 rag-1 liver		40
TH2 7 day	27.11	rag-1 spleen	23.97
TH1 3 week Balb/C	40	rag-1 thymus	26.29
TH2 3 week Balb/C	26.95	rag-1 kidney	40
preT	40	w.t. Peyers patches	27.04
D1.1 resting	40	w.t. mesenteric lymph nodes	40
D1.1 con A stim.	40	w.t. colon	26.63
CDC35 resting	40	Braf:ER (-) oligo dT	40
CDC35 con A stim.	39.83	TH1 3 week C57 Bl/6	26.78
Mel 14+ naive T	40	TH2 3 week C57 Bl/6	40
Mel14+ TH1	40	TH1 3 week Balb/C fresh	40
Mel 14+ TH2	31.22	TH2 3 week Balb/C fresh	40
A20	27.39	b.m. DC (YJL) resting	40
CH12	28.18	b.m. DC (YJL) aCD40 stim.	40
lg. B cell	26.35	b.m. mf + LPS + aIL-10R	40
LPS spleen	21.58	b.m. mf + LPS + IL-10	40
macrophage	40	peritoneal mf	40
J774 resting	24.99	MC-9/MCP-12 pMET7	40
J774 +LPS + anti-IL-10	28.41	EC	40
J774 +LPS + IL-10	27.57	EC + TNFa	40
Nippo-infected lung	26.98	bEnd3 + TNFa	40
IL-10 K.O. spleen	25.43	bEnd3 + TNFa + IL-10	40
IL-10 K.O. colon	23.68	ApoE aorta 5 month	35.16
asthmatic lung	37.45	ApoE aorta 12 month	35.47
w.t. lung	40	NZ B/W kidney	37.17
w.t. spleen	39.95	SNZ B/W spleen	25.25
rag-1 heart	40) tolerized & challenged lung	40
rag-1 brain	40	Aspergillus lung	39.26

57

LIBRARY	Ct_gene	LIBRARY	Ct_gene
rag-1 testes	40	Nippo IL-4 K.O. lung	26.13
Influenza lung	37.13	Nippo anti IL-5 lung	34.73
b common lung 2 month	39.33	w.t. thymus	40
IL-10 K.O. stomach	27.3	w.t. stomach	30.14
IL-10 K.O. MLN aIL-12	40	MM2/MM3 resting pME	40
IL-10 K.O. MLN +IL-10	37.97	MM2/MM3 activated. pME	40
Rag-2 Hh- colon	26.95	Taq_control_water	50
Rag-2 Hh+ colon	22.94	Taq_control_genomic_1	50
IL-7 K.O./Rag-2 Hh- colon	26.77	Taq_control_genomic_2	50
IL-7 K.O./Rag-2 Hh+ colon	24.24	w.t. d17 spinal cord EAE	40
•		model	
transfer model IBD	23.01	TNF K.O. d17 spinal cord	40
		EAE model	
w.t. C57 Bl/6 aorta	40	TNF K.O. spinal cord	27.99
w.t. spinal cord	38.8	TNF K.O. spleen	24.93
w.t. C57 Bl/6 spleen	26.38	G.F. ears (skin)	40
w.t. C57 Bl/6 pancreas	40		

The primate, e.g., human, TLR-L1 is expressed in TF-1 cells, D6 cells, and barely detectable in resting U937 cells, resting Jurkat cells, and pooled NK cells. In tissues, it is found in fetal uterus, fetal ovary, allergic lung, and fetal testis. Lower levels are found in fetal kidney, fetal small intestine, fetal brain, fetal adipose tissue, normal lung pool, and fetal lung.

5

The primate, e.g., human, TLR-L2, TLR-L3, and TLR-L4 seem to be expressed in brain tissue.

The primate, e.g., human, TLR-L5 seems to be expressed in unstimulated A549,

activated A549, MRC5, and Bc cell lines. Among tissues, it is most highly expressed in fetal uterus, fetal small intestine, and lesser in fetal lung, fetal kidney, fetal liver, and fetal ovary. It is just detectable in fetal brain, fetal adipose, fetal testes, psoriasis skin, and various intestinal samples.

58

The 5685C6 probes show positive hybridization to subtraction libraries of Th2 minus Th1 polarized cells, and absence of hybridization to libraries of Th1 minus Th2 polarized cells. This suggests that the probe is present selectively in Th2 polarized cells, and can serve as a marker for such cell type. PCR techniques should confirm the expression profile.

Structurally, this protein exhibits similarities to other proteins possessing a thioredoxin fold, including a peroxidase protein, e.g., glutathione peroxidase. See Choi, et al. (1998) Nature Structural Biol. 5:400-406. Thioredoxin has been reported to exhibit certain chemoattractant activities. See Bertini, et al. (1999) J. Expt'l Med. 189:1783-1789.

5

10

15

20

25

30

TaqMan primers were designed for all four novel claudin transcripts. These primer sets were used to screen a panel of human libraries representing different cell types, tissues, and disease states, and two extended cDNA panels. The cDNA panels were composed of samples derived from either normal or diseased human lung or intestine. The claudin genes are some of the most highly regulated genes detected. Moreover, claudin D8 shows the greatest reciprocal regulation between Crohn's and Ulcerative colitis samples, making it a good candidate in future diagnostic panels for these diseases.

claudin-D2: In library southerns, expression is highest in one Crohn's colon, the fetal intestine, and two epithelial cell lines, lower level expression in fetal lung, kidney, ovary and testes. In human cDNA panels, this is highly up-regulated in 8/9 Crohn's disease, both with and without steroid treatment (mean induction = 53x, n=9). In addition, claudin-D2 is also induced in 9/12 ulcerative colitis samples (mean induction = 8.2x), but this induction is significantly less than that observed in the Crohn's disease samples. Also up-regulated (mean induction=29 x) in 12/13 interstitial lung disease samples (idiopathic pulmonary fibrosis, hypersensitive pneumonitis, and eosinophilic granuloma).

claudin-D8: In library southerns, expression is highest in fetal kidney and normal colon. Also, expressed in ulcerative colitis colon, thyroid, and fetal lung. No expression is observed in the cells on the panel. In human cDNA panels, high level expression in the gut. Little to no expression in all Crohn's disease samples mean reduction 130×10^{-2} . Some ulcerative colitis samples also have reduced claudin-D8 expression, but the pattern is heterogeneous. In contrast, claudin-D8 is up-regulated in several interstitial lung disease samples (12/15, mean induction = 9x), but the level of expression in these samples is on the

order of ten fold lower than in normal colon. It is also induced in primary human bronchial epithelial cells by I-309.

5

10

15

20

25

30

claudin-D17: In library southerns, overall the expression level measured is low relative to the other claudins described here, on the order of 100 fold lower. It is unclear whether the expression level is actually lower or whether the primers for this gene are insensitive (non-optimal). Expression is highest in one of the asthma lungs and in psoriatic skin. No expression is observed in the cell lines on the panel. In human cDNA panels, the expression is increased in 8/11 ulcerative colitis samples (mean induction = 13x), while the expression is unchanged in Crohn's disease samples. Expressed at low level in primary bronchial epithelial cell lines, induced by I-309. Otherwise, level is too low to detect except in sporadic samples.

claudin-D7.2: In library southerns, expressed at highest level in human fetal and adult lung, monkey lungs, and in one Crohn's colon sample. Lower level expression in the two epithelial (A549 and CHA) and one fibroblast (MRC5) cell lines on the panel. In human cDNA panels, expressed at a high level in the gut and an even higher level in the lung. Upregulated in Crohn's disease samples from patients which have not been treated with steroids (mean induction = 3.7x, n=4). No consistent modulation of this gene in any of the lung diseases examined on this panel.

Claudin family structure: If the genomic structural organization of Claudin family members is based upon that of Paracellin-1, then the proteins would all be encoded by 5 exons. The putative splice sites and exon numbers are predictable, corresponding to the residues of D2 about: 2 codons upstream from M1; A43, A75, G129, and C182; and transmembrane segments corresponding to about G17-V36, M83-C104, V117-H141, and L164-Q188. Paracellin has an extra 60 amino acids at its N-terminus, which is located on the cytoplasmic side of the membrane.

Disease Associations: Claudin-D2 is up-regulated in 8/9 Crohn's disease relative to the control samples, while claudin-D8 is down-regulated. All claudins, described in this invention disclosure, show disease association as described above.

The claudins may form part of a diagnostic panel of genes that could distinguish Crohn's disease from ulcerative colitis, or assist in the determination of disease severity in either or both diseases. For example, claudin-D2 is expressed at higher levels in Crohn's disease than in ulcerative colitis. In contrast, the claudin-D8, cluster 1645577, is expressed at

60

very low levels in Crohn's disease samples, and is less dramatically reduced in most ulcerative colitis samples. See, e.g., Simon, et al. (1999) Science 285:103-106; Hirano, et al. (19xx)

Genome Research 10:659-663; Morita, et al. (1999) Proc. Nat'l Acad. Sci. USA 96:511-516;

Anderson and Van Itallie (1999) Current Biology 9:R922-R924; and Furuse, et al. (1999) J.

Cell Biol. 147:891-903.

Introduction of an adenovirus or another expression vector expressing the claudin-D8 ortholog into the intestines of patients with inflammatory bowel disease may improve intestinal barrier function and ameliorate disease.

5

10

15

20

25

30

In contrast, antibodies to one of the claudins described here may be able to: induce an intracellular signal that could promote tight junction formation and lead to improved intestinal barrier function; block entry of pathogenic agents, which may play a causative role in initiation or maintenance of either Crohn's disease or ulcerative colitis; promote migration of myeloid cells across tight junctions and allow clearance of pathogenic agents prior to infection of the epithelium.

Expression of schlafen family members in fibroblasts/ thymoma cells retards or arrests cell growth. They guide cell growth and T-cell development, and are an integral component of the machinery that maintains T-cell quiescence. They may have important roles in the development or maintenance of autoimmune disorders. The mouse schlafens participate in the regulation of the cell cycle. This family is characterized by two splice variants: a short and a long form.

Schlafen B: 748 aa; ORF. Quantitative PCR analysis reveals in T cells, resting DC, M1 macrophage cell panel. Induced in Hashimoto's thyroiditis, fetal kidney, fetal uterus, and fetal spleen. Slightly induced in Crohn's colon.

Schlafen C: 891 aa, full ORF. Quantitative PCR data revealed this to be significantly up-regulated in all Crohn's samples, asthmatic lung, Ascaris lung, Hashimoto's thyroiditis, and fetal tissues compared to control.

Schlafen D: 578 aa, full ORF. The quantitative PCR data for human schlafen D revealed that it is significantly differentially regulated in Crohn's disease and Ulcerative Colitis compared to normal colon. Also it appears to be highly expressed in many developing tissues (fetal) and disease states (allergic, Ascaris and pneumocystis carnii lungs, Crohn's colon, ulcerative colitis, and Psoriasis skin) compared to cell lines.

61

Schlafen E: 897 aa, full ORF. Quantitative PCR analysis reveals expression in the colon, fetal liver, fetal lung, fetal ovary, and fetal uterus, and significantly upregulated in one Crohn's sample and highly induced in Hashimoto's thyroiditis.

Schlafen F: 358 aa; full ORF. Distribution analysis is not complete. Similar samples may isolated in other species for evaluation.

V. Cloning of species counterparts

5

10

20

25

30

Various strategies are used to obtain species counterparts of, e.g., the DIRS4, preferably from other primates or rodents. One method is by cross hybridization using closely related species DNA probes. It may be useful to go into evolutionarily similar species as intermediate steps. Another method is by using specific PCR primers based on the identification of blocks of similarity or difference between genes, e.g., areas of highly conserved or nonconserved polypeptide or nucleotide sequence.

15 VI. Production of mammalian protein

An appropriate, e.g., GST, fusion construct is engineered for expression, e.g., in E. coli. For example, a mouse IGIF pGex plasmid is constructed and transformed into E. coli. Freshly transformed cells are grown, e.g., in LB medium containing 50 _g/ml ampicillin and induced with IPTG (Sigma, St. Louis, MO). After overnight induction, the bacteria are harvested and the pellets containing, e.g., the DIRS4 protein, are isolated. The pellets are homogenized, e.g., in TE buffer (50 mM Tris-base pH 8.0, 10 mM EDTA and 2 mM pefabloc) in 2 liters. This material is passed through a microfluidizer (Microfluidics, Newton, MA) three times. The fluidized supernatant is spun down on a Sorvall GS-3 rotor for 1 h at 13,000 rpm. The resulting supernatant containing the cytokine receptor protein is filtered and passed over a glutathione-SEPHAROSE column equilibrated in 50 mM Tris-base pH 8.0. The fractions containing the DIRS4-GST fusion protein are pooled and cleaved, e.g., with thrombin (Enzyme Research Laboratories, Inc., South Bend, IN). The cleaved pool is then passed over a Q-SEPHAROSE column equilibrated in 50 mM Tris-base. Fractions containing DIRS4 are pooled and diluted in cold distilled H2O, to lower the conductivity, and passed back over a fresh Q-Sepharose column, alone or in succession with an immunoaffinity

antibody column. Fractions containing the DIRS4 protein are pooled, aliquoted, and stored in the -70° C freezer.

Comparison of the CD spectrum with cytokine receptor protein may suggest that the protein is correctly folded. See Hazuda, et al. (1969) <u>J. Biol. Chem.</u> 264:1689-1693.

For other genes, e.g., membrane proteins, the protein may be best expressed on cell surfaces. Those may be in prokaryote expression systems, or eukaryotes. Surface expressed forms will most likely have conformations consistent with the natural interaction with lipid.

VII. Determining physiological forms of receptors

5

10

15

20

25

30

The cellular forms of receptors for ligands can be tested with the various ligands and receptor subunits provided, e.g., IL-10 related sequences. In particular, multiple cytokine receptor like ligands have been identified, see, e.g., USSN 60/027,368, 08/934,959, and 08/842,659, which are incorporated herein by reference.

Cotransformation of the DIRS4 with putative other receptor subunits may be performed. Such cells may be used to screen putative cytokine ligands, such as the AK155, for signaling. A cell proliferation assay may be used.

In addition, it has been known that many cytokine receptors function as heterodimers, e.g., a soluble alpha subunit, and transmembrane beta subunit. Subunit combinations can be tested now with the provided reagents. In particular, appropriate constructs can be made for transformation or transfection of subunits into cells. Combinatorial transfections of transformations can make cells expressing defined subunits, which can be tested for response to the predicted ligands. Appropriate cell types can be used, e.g., 293 T cells, with, e.g., an NF b reporter construct.

Biological assays for receptors will generally be directed to the ligand binding feature of the protein or to the kinase/phosphatase activity of the receptor. The activity will typically be reversible, as are many other enzyme reactions, and may mediate phosphatase or phosphorylase activities, which activities are easily measured by standard procedures. See, e.g., Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature 363:736-738.

The family of cytokines contains molecules which are important mediators of hematopoiesis or inflammatory disease. See, e.g., Nelson and Martin (eds. 2000) Cytokines in Pulmonary Disease Dekker, NY; Ganser and Hoelzer (eds. 1999) Cytokines in the Treatment of Hematopoietic Failure Dekker, NY: Remick and Friedland (eds. 1997) Cytokines in Health and Disease Dekker, NY; Dinarello (1996) Blood 87:2095-2147; and Thomson (ed. 1994) The Cytokine Handbook Academic Press, San Diego. Ligand and receptors are very important in the signaling process.

VIII. Antibodies specific for proteins

5

10

15

20

25

30

Inbred Balb/c mice are immunized intraperitoneally with recombinant forms of the protein, e.g., purified DIRS4 or stable transfected NIH-3T3 cells. Animals are boosted at appropriate time points with protein, with or without additional adjuvant, to further stimulate antibody production. Serum is collected, or hybridomas produced with harvested spleens.

Alternatively, Balb/c mice are immunized with cells transformed with the gene or fragments thereof, either endogenous or exogenous cells, or with isolated membranes enriched for expression of the antigen. Serum is collected at the appropriate time, typically after numerous further administrations. Various gene therapy techniques may be useful, e.g., in producing protein in situ, for generating an immune response. Serum may be immunoselected to prepare substantially purified antibodies of defined specificity and high affinity.

Monoclonal antibodies may be made. For example, splenocytes are fused with an appropriate fusion partner and hybridomas are selected in growth medium by standard procedures. Hybridoma supernatants are screened for the presence of antibodies which bind to the DIRS4, e.g., by ELISA or other assay. Antibodies which specifically recognize specific DIRS4 embodiments may also be selected or prepared.

In another method, synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (ed. 1991) <u>Current Protocols in Immunology</u> Wiley/Greene; and Harlow and Lane (1989) <u>Antibodies: A Laboratory Manual Cold Spring Harbor Press.</u> In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods. Nucleic acids may also be introduced into cells in an animal to produce the antigen, which serves to elicit an immune response. See, e.g., Wang, et al. (1993)

64

<u>Proc. Nat'l. Acad. Sci.</u> 90:4156-4160; Barry, et al. (1994) <u>BioTechniques</u> 16:616-619; and Xiang, et al. (1995) <u>Immunity</u> 2: 129-135.

Moreover, antibodies which may be useful to determine the combination of the DIRS4 with a functional alpha subunit may be generated. Thus, e.g., epitopes characteristic of a particular functional alpha/beta combination may be identified with appropriate antibodies.

IX. Production of fusion proteins

Various fusion constructs are made, e.g., with DIRS4. A portion of the appropriate gene is fused to an epitope tag, e.g., a FLAG tag, or to a two hybrid system construct. See, e.g., Fields and Song (1989) <u>Nature</u> 340:245-246.

The epitope tag may be used in an expression cloning procedure with detection with anti-FLAG antibodies to detect a binding partner, e.g., ligand for the respective cytokine receptor. The two hybrid system may also be used to isolate proteins which specifically bind to DIRS4.

15

20

25

30

10

5

X. Structure activity relationship

Information on the criticality of particular residues is determined using standard procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the positions identified above, and evaluating biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to focus on specific positions to determine the residues which can be substituted to either retain, block, or modulate biological activity.

Alternatively, analysis of natural variants can indicate what positions tolerate natural mutations. This may result from populational analysis of variation among individuals, or across strains or species. Samples from selected individuals are analyzed, e.g., by PCR analysis and sequencing. This allows evaluation of population polymorphisms.

XI. Isolation of a ligand for receptor

A cytokine receptor can be used as a specific binding reagent to identify its binding partner, by taking advantage of its specificity of binding, much like an antibody would be used. Typically, the binding receptor is a heterodimer of receptor subunits. A binding reagent

65

is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods.

5

10

15

20

25

30

The binding composition is used to screen an expression library made from a cell line which expresses a binding partner, i.e., ligand, preferably membrane associated. Standard staining techniques are used to detect or sort surface expressed ligand, or surface expressing transformed cells are screened by panning. Screening of intracellular expression is performed by various staining or immunofluorescence procedures. See also McMahan, et al. (1991) EMBO J. 10:2821-2832.

For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in PBS, for 30 min at room temperature. Rinse once with PBS. Then plate COS cells at $2-3 \times 10^5$ cells per chamber in 1.5 ml of growth media. Incubate overnight at 37° C.

On day 1 for each sample, prepare 0.5 ml of a solution of 66 µg/ml DEAE-dextran, 66 _M chloroquine, and 4 µg DNA in serum free DME. For each set, a positive control is prepared, e.g., of DIRS4-FLAG cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free DME. Add the DNA solution and incubate 5 hr at 37° C. Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may be stored at -80° C after all liquid is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin (0.1%) with 32 _l/ml of 1 M NaN 3 for 20 min. Cells are then washed with HBSS/saponin 1X. Add appropriate DIRS4 or DIRS4/antibody complex to cells and incubate for 30 min. Wash cells twice with HBSS/saponin. If appropriate, add first antibody for 30 min. Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of

66

buffer plus 4 drops DAB plus 2 drops of H_2O_2 per 5 ml of glass distilled water. Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90° C.

Evaluate positive staining of pools and progressively subclone to isolation of single genes responsible for the binding.

5

10

15

20

25

Alternatively, receptor reagents are used to affinity purify or sort out cells expressing a putative ligand. See, e.g., Sambrook, et al. or Ausubel, et al.

Another strategy is to screen for a membrane bound receptor by panning. The receptor cDNA is constructed as described above. The ligand can be immobilized and used to immobilize expressing cells. Immobilization may be achieved by use of appropriate antibodies which recognize, e.g., a FLAG sequence of a DIRS4 fusion construct, or by use of antibodies raised against the first antibodies. Recursive cycles of selection and amplification lead to enrichment of appropriate clones and eventual isolation of receptor expressing clones.

Phage expression libraries can be screened by mammalian DIRS4. Appropriate label techniques, e.g., anti-FLAG antibodies, will allow specific labeling of appropriate clones.

All citations herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the specific embodiments that have been presented herein by way of example.

WHAT IS CLAIMED IS:

- 1. A substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of SEQ ID NO: 2 (DIRS4); SEQ ID NO: 9, 11, 13, or 53 (TNFx or TNFy); SEQ ID NO: 15, 17, 19, 21, 23, 25, or 27 (TLR-L1 through TLR-L5); SEQ ID NO: 29 (TGFx); SEQ ID NO: 31 or 33 (5685C6); SEQ ID NO: 35, 37, 39, or 41 (claudins); or SEQ ID NO: 43, 45, 47, 49, or 51 (schlafens).
- 10 2. The substantially pure or isolated antigenic polypeptide of Claim 1, wherein said distinct nonoverlapping segments of identity:
 - a) include one of at least eight amine acids;
 - b) include one of at least four amino acids and a second of at least five amino acids;
 - c) include at least three segments of at least four, five, and six amino acids; or
 - d) include one of at least twelve amino acids.
 - 3. The composition of matter of Claim 1, wherein said polypeptide:
 - a) is unglycosylated;

15

- b) is from a primate, such as a human;
- c) comprises at least contiguous seventeen amino acids of said SEQ ID NO;
 - d) exhibits at least four nonoverlapping segments of at least seven amino acids of said SEQ ID NO;
 - e) has a length at least about 30 amino acids;
 - f) has a molecular weight of at least 30 kD with natural glycosylation;
- 25 g) is a synthetic polypeptide;
 - h) is attached to a solid substrate;
 - i) is conjugated to another chemical moiety; or
 - j) comprises a detection or purification tag, including a FLAG, His6, or Ig sequence.
- 30 4. A composition comprising:
 - a) a substantially pure polypeptide of Claim 1;

68

- b) a sterile polypeptide of Claim 1; or
- c) said polypeptide of Claim 1 and a carrier, wherein said carrier is:
 - i) an aqueous compound, including water, saline, and/or buffer; and/or
 - ii) formulated for oral, rectal, nasal, topical, or parenteral administration.

5

- 5. A kit comprising a polypeptide of Claim 1, and:
 - a) a compartment comprising said polypeptide; or
 - b) instructions for use or disposal of reagents in said kit.
- 10 6. A binding compound comprising an antigen binding site from an antibody, which specifically binds to a polypeptide of Claim 1, wherein:
 - a) said binding compound is in a container;
 - b) said polypeptide is from a human;
 - c) said binding compound is an Fv, Fab, or Fab2 fragment;
 - d) said binding compound is conjugated to another chemical moiety; or
 - e) said antibody:
 - i) is raised to a recombinant polypeptide of Claim 1;
 - ii) is raised to a purified polypeptide of Claim 1;
 - iii) is immunoselected;
- 20

- iv) is a polyclonal antibody;
- v) binds to a denatured antigen;
- vi) exhibits a Kd to antigen of at least 30 μM;
- vii) is attached to a solid substrate, including a bead or plastic membrane;
- viii) is in a sterile composition; or
- ix) is detectably labeled, including a radioactive or fluorescent label.
- 25
- 7. A kit comprising said binding compound of Claim 6, and:
 - a) a compartment comprising said binding compound; or
 - b) instructions for use or disposal of reagents in said kit.

10

15

20

- 8. A method of producing an antigen:antibody complex, comprising contacting under appropriate conditions a primate polypeptide with an antibody of Claim 7, thereby allowing said complex to form.
- 5 9. A method of producing an antigen:antibody complex, comprising contacting under appropriate conditions a polypeptide of Claim 1 with an antibody which binds thereto, thereby allowing said complex to form.
 - 10. A method of producing a binding compound comprising:
 - a) immunizing an immune system with a polypeptide of Claim 1; or
 - b) introducing a nucleic acid encoding said polypeptide of Claim 1 to a cell under conditions leading to an immune response, thereby producing said binding compound; or
 - c) selecting for a phage display library for those phage which bind to said polypeptide of Claim 1.
 - 11. A composition comprising:
 - a) a sterile binding compound of Claim 7, or
 - b) said binding compound of Claim 7 and a carrier, wherein said carrier is:
 - i) an aqueous compound, including water, saline, and/or buffer, and/or
 - ii) formulated for oral, rectal, nasal, topical, or parenteral administration.
 - 12. An isolated or recombinant nucleic acid encoding said polypeptide of Claim 1, wherein said:
- a) polypeptide is from a primate; or
 - b) said nucleic acid:
 - i) encodes an antigenic polypeptide;
 - ii) encodes a plurality of antigenic polypeptide sequences of SEQ ID NO:2, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53;

WO 02/20569

30 16.

A nucleic acid which:

		iii) exhibits identity over at least thirteen nucleotides to a natural cDNA
		encoding said segment;
		iv) is an expression vector;
		v) further comprises an origin of replication;
5		vi) is from a natural source;
		vii) comprises a detectable label;
		viii) comprises synthetic nucleotide sequence;
		ix) is less than 6 kb, preferably less than 3 kb;
		x) is a hybridization probe for a gene encoding said polypeptide; or
10		xi) is a PCR primer, PCR product, or mutagenesis primer.
	13.	A cell comprising said recombinant nucleic acid of Claim 12.
	14.	The cell of Claim 13, wherein said cell is:
15		a) a prokaryotic cell;
		b) a eukaryotic cell;
		c) a bacterial cell;
		d) a yeast cell;
		e) an insect cell;
20		f) a mammalian cell;
		g) a mouse cell;
		h) a primate cell; or
		i) a human cell.
25	15.	A kit comprising said nucleic acid of Claim 12, and:
		a) a compartment comprising said nucleic acid;
		b) a compartment further comprising a primate polypeptide; or
		c) instructions for use or disposal of reagents in said kit.

- a) hybridizes under wash conditions of 30 minutes at 37° C and less than 2M salt to the coding portion of SEQ ID NO: 1, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, or 52; or
- b) exhibits identity over a stretch of at least about 30 nucleotides to a SEQ ID NO: 1, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, or 52.
- 17. The nucleic acid of Claim 16, wherein:
 - a) said wash conditions are at 45° C and/or 500 mM salt; or
- b) said stretch is at least 55 nucleotides.
 - 18. The nucleic acid of Claim 16, wherein:
 - a) said wash conditions are at 55° C and/or 150 mM salt; or
 - b) said stretch is at least 75 nucleotides.

15

20

19.

5

A method of making:

- a) a duplex nucleic acid comprising contacting:
 - i) a nucleic acid of Claim 12 with a complementary nucleic acid, under appropriate conditions, thereby resulting in hybridization to form said complex; or
 - ii) a nucleic acid complementary to said nucleic acid of Claim 12 with its complementary nucleic acid, under appropriate conditions, thereby resulting in hybridization to form said complex; or
- b) a polypeptide comprising culturing a cell comprising said nucleic acid of Claim 12 under conditions resulting in expression of said nucleic acid.

25

30

20. A method of:

- a) modulating physiology or development of a cell comprising contacting said cell with a polypeptide comprising SEQ ID NO: 9, 11, 13, 29, 31, 33, or 53;
- b) modulating physiology or development of a cell comprising contacting said cell with a binding compound of Claim 6 which binds to SEQ ID NO: 9, 11, 13, 29,

72

- 31, or 33, thereby blocking signaling mediated by a protein comprising said SEQ ID NO;
- c) labeling a cell comprising contacting said cell with a binding compound which binds to SEQ ID NO: 2, 15, 17, 19, 21, 23, 25, or 27; or
- d) diagnosing a medical condition comprising a step of evaluating expression of nucleic acid comprising SEQ ID NO: 34, 36, 38, 40, 42, 44, 46, 48, or 50.

WO 02/20569

73

PCT/US01/28013

SEQUENCE IDENTIFICATION NUMBERS

SEQ ID NO: 1 is primate DIRS4 nucleotide sequence.

SEQ ID NO: 2 is primate DIRS4 polypeptide sequence.

5 SEQ ID NO: 3 is tissue factor polypeptide sequence.

SEQ ID NO: 4 is primate IFN $\alpha\beta$ R polypeptide sequence.

SEQ ID NO: 5 is CRF1-4 polypeptide sequence.

SEQ ID NO: 6 is cytor x polypeptide sequence.

SEQ ID NO: 7 is cytor7 polypeptide sequence.

10 SEQ ID NO: 8 is primate TNFx nucleic acid sequence.

SEQ ID NO: 9 is primate TNFx polypeptide sequence.

SEQ ID NO: 10 is rodent TNFx nucleic acid sequence.

SEQ ID NO: 11 is rodent TNFx polypeptide sequence.

SEQ ID NO: 12 is primate TNFy nucleic acid sequence.

15 SEQ ID NO: 13 is primate TNFy polypeptide sequence.

SEQ ID NO: 14 is primate TLR-L1 nucleic acid sequence.

SEQ ID NO: 15 is primate TLR-L1 polypeptide sequence.

SEQ ID NO: 16 is rodent TLR-L1 nucleic acid sequence.

SEQ ID NO: 17 is rodent TLR-L1 polypeptide sequence.

20 SEQ ID NO: 18 is primate TLR-L2 nucleic acid sequence.

SEQ ID NO: 19 is primate TLR-L2 polypeptide sequence.

SEQ ID NO: 20 is rodent TLR-L2 nucleic acid sequence.

SEQ ID NO: 21 is rodent TLR-L2 polypeptide sequence.

SEQ ID NO: 22 is primate TLR-L3 nucleic acid sequence.

25 SEQ ID NO: 23 is primate TLR-L3 polypeptide sequence.

SEQ ID NO: 24 is primate TLR-L4 nucleic acid sequence.

SEQ ID NO: 25 is primate TLR-L4 polypeptide sequence.

SEQ ID NO: 26 is primate TLR-L5 nucleic acid sequence.

SEQ ID NO: 27 is primate TLR-L5 polypeptide sequence.

30 SEQ ID NO: 28 is primate TGFx nucleic acid sequence.

SEQ ID NO: 29 is primate TGFx polypeptide sequence.

74

SEQ ID NO: 30 is primate 5685C6 nucleic acid sequence.

SEQ ID NO: 31 is primate 5685C6 polypeptide sequence.

SEQ ID NO: 32 is rodent 5685C6 nucleic acid sequence.

SEQ ID NO: 33 is rodent 5685C6 polypeptide sequence.

SEQ ID NO: 34 is primate claudin-D2 nucleic acid sequence.

SEQ ID NO: 35 is primate claudin-D2 polypeptide sequence.

SEQ ID NO: 36 is primate claudin-D8 nucleic acid sequence.

SEQ ID NO: 37 is primate claudin-D8 polypeptide sequence.

SEQ ID NO: 38 is primate claudin-D17 nucleic acid sequence.

SEQ ID NO: 39 is primate claudin-D17 polypeptide sequence.

SEQ ID NO: 40 is primate claudin-D7.2 nucleic acid sequence.

SEQ ID NO: 41 is primate claudin-D7.2 polypeptide sequence.

SEQ ID NO: 42 is primate schlafen B nucleic acid sequence.

SEQ ID NO: 43 is primate schlafen B polypeptide sequence.

15 SEQ ID NO: 44 is primate schlafen C nucleic acid sequence.

SEQ ID NO: 45 is primate schlafen C polypeptide sequence.

SEQ ID NO: 46 is primate schlafen D nucleic acid sequence.

SEQ ID NO: 47 is primate schlafen D polypeptide sequence.

SEQ ID NO: 48 is primate schlafen E nucleic acid sequence.

20 SEQ ID NO: 49 is primate schlafen E polypeptide sequence.

SEQ ID NO: 50 is primate schlafen F nucleic acid sequence.

SEQ ID NO: 51 is primate schlafen F polypeptide sequence.

SEQ ID NO: 52 is rodent TNFy nucleic acid sequence.

SEQ ID NO: 53 is rodent TNFy polypeptide sequence.

TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	-METPAWPRVPRPETAVARTLLLGWVFAQVAGASGTTN-TMAGPERWGPLLLCLLQAAPGRPR-L MLLSQNAFIFRSLNLVLMVYISLVFGISYDSPDYTMAWSLGSWLGGCLLVSALGMVMMPKHCFLGFLISFFLTGVAGTQSTHESMRAPGRPALRPLPLPPLLLLLLAAPWGRAVPCVSGGL
TissueFactor 1274993aaR hIFNabR CRF2-4 cytor x cytor7	VAAYNLTWKSTNFKTILEWEPKPVN-QVYTVQISTKS APPQNVTLLSQNFSVYLTWLPGLGNPQD-VTYFVAYQSSP DESCTFKISLRNFRSILSWE-LKNHSIVPTHYTLLYTIMS PPPENVRMNSVNFKNILQWESPAFAKGN-LTFTAQYLSY- LKPQRVQFQSRNFHNILQWQPGRALTGNSSVYFVQYKIYG PKPANITFLSINMKNVLQWTPPEGLQGVKVTYTVQYFIYG
TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	GDWKSKCFYTTDTECDLTDEIVKDVKQTYLARVFSY TRRRWREVEECAGTKELLCSMMCLKKQDLYNKFKGRVRTV KPEDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGRIFQDKCMNTTLTECDFSSLS-KYGDHTLRVRAE -QRQWKNKEDCWGTQELSCDLTSET-SDIQEPYYGRVRAA -QKKWLNKSECRNINRTYCDLSAET-SDYEHQYYAKVKAI
TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	PAGNVESTGSAGEPLYENSPEFTPYLETNLGQPTIQSFEQ SPSSKSPWVESEYLDYLFEVEPAPP-VLVLTQ FSGNTTLFSCSHNFWLAIDMSFEPP-EFEIVG FADEHSDWVNIT-FCPVDDTIIGPP-GMQVEV SAGSYSEWSMTPRFTPWWETKIDPP-VMNITQ WGTKCSKWAESGRFYPFLETQIGPP-EVALTT
TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	VGTKVNVTVEDERTLVR-RNNTFLSLRDVFGKDLIYTLYY T-EEILSANATYQLPPCMPPLDLKYEVAF FTNHINVVVKFPSIVEEELQFDLSLVIE-EQSEGIVK LADSLHMRFLAPKIENEYETWTMKNVYN-SWTYNVQY VNGSLLVILHAPNLPYRYQKEKNVSIEDYYELLYRVFI DEKSISVVLTAPEKWKRNPEDLPVSMQQIYS-NLKYNVSV

FIG.1A

WO 02/20569 PCT/US01/28013 2/19

TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7 TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	WKSSSG-KKTAKTNTNEFLIDVDKGENYCFSVQAVIP WKEGAGNKVGSSFPAPRLGPLLHPFLLRFFSP KHKPEIKGNMSGNFTYIIDK-LIPNTNYCVSVYLEHS WKNGTDEKFQITPQYDFEVLRNLEPWTTYCVQVRGFLP INNSLEKEQKVYEGAHRAVEIEA-LTPHSSYCVVAEIYQP LNTKSNR-TWSQCVTNHTLVLTW-LEPNTLYCVHVESFVP SRTVNRKSTDS-PVECMGQEKGEFREIFYIISQPAPAPLLQEVFPVHSFREIFYII DEQAVIKS-PLKCTLLPPGQESESAESAKIGGIITVF DR-NKAGEWS-EPVCEQTTHDETVPSWMVAVIL ML-DRRSQRS-EERCVEIP
TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	GAVAFVVIILVIILAISLHKCRKAG LIALVLTSTIVTLKWIGYICLRNSLPKVLNFHNFLAW MASVFMVCLALLGCFSLLWCVYKKTKY
TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	PFPNLPPLEAMDMVEVIYINRKKKVWDYNYDDES-DSDTE AFS FFVPAEKIVINFITLNISDDSKISHQDMSLLGKSSDVSSL
TissueFactor 1274993R hIFNabR CRF2-4 cytor x cytor7	EN AAPRTSGGGYTMHGLTVRPLGQASATSTESQLIDPESEEEPRNSLPQHLKEFLGHPHHNTLLFFSFPLSDEN NDPQPSGNLRPPQEEEEVKHLGYASHLMEIFCDSEENTEG

TissueFactor 1274993R	SP
hIFNabR CRF2-4	PEEDYSSTEGSGGRITFNVDLNSVFLRVLDDEDSDDLEAP
cytor x	
cytor7	SLQEEVSTQGTLLESQAALAVLGPQTLQYSYTPQLQDLDP
TissueFactor	
1274993R	
hIFNabR CRF2-4	PDLPEVDVELPTMPKDSP-QQLELLSGPCERRKSPLQDPF
cytor x	
cytor7	TSLTQQESLSRTIPPDKTVIEYEYDVRTTDICAGPEEQEL
TissueFactor	LNVS
1274993R	
hIFNabR	LMLSSHLEEMVDPEDPDNVQSNHLLASGEGTQ
CRF2-4	LSVIAEDSESG-KQNPDS
cytor x	
cytor7	LAQEHTDSEEGPEEEPSTTLVDWDPQTGRLCIPSLSSFDQ
TissueFactor	
1274993R	
hIFNabR	PTFPSPSSEGLWSEDAPSDQSDTSES
CRF2-4	CSLGTPPGQGPQS
cytor x	
cytor7	DSEGCEPSEGDGLGEEGLLSRLXEEPAPDRPPGENETYLM
TissueFactor	
1274993R	
hIFNabR	DVDLGDGYIMR
CRF2-4 aa	
cytor x	
cytor7	OFMEEWGLYVOMEN

7 50 0	51 100 0	92 150 0	142 200 25	192 250 75	242 300 124	
AGREGEE- MWAWGWAAAALLWLQTAGARQELKKSRQLFARVDSPNITTSNREGFPG	PSQASGPEFSDAHMTWLNFVRRPDDGALRKRCGSRDKKPRDLFG SVKPPEASGPELSDAHMTWLNFVRRPDDGSSRKRCRGRDKKSRGLSGLPG	PPGPPGAEVTAETLLHEFQELLKEATERRFSGLLDPLLPQG PPGPPGPPGPPGSPGVGVTPEALLQEFQEILKEATELRFSGLPDTLLPQE	RGLRLVGEAFHCRLQGPRRVDKRTLVELHGFQAPAAQGAFLRGSGLSLAS PSQRLVVEAFYCRLKGPVLVDKKTLVELQGFQAPTTQGAFLRGSGLSLSL HELGVYYLPDAEGAFRRGPGLNLTS	GRFTAPVSGIFQFSASLHVDHSELQGKARLRARDVVCVLICIESLCQRHT GRFTAPVSAIFQFSASLHVDHSELQGRGRLRTRDMVRVLICIESLCHRHT GQYRAPVAGFYALAATLHVALGEPPRRGPPRPRDHLRLLICIQSRCQRNT	CLEAVSGLESNSRVFTLQVQGLLQLQAGQYASVFVDNGSGAVLTIQAGSS SLEAVSGLESNSRVFTVQVQGLLHLQSGQYVSVFVDNSSGAVLTIQNTSS SLEAIMGLESSSELFTISVNGVLYLQMGQWTSWACERPP-QALPLRGKWS	FIG. 2
TMTOT	FSDAH LSDAH	GSPGV	CRLQG	OFSAS OFSAS ALAAT	SRVFT SRVFT SELFT	250 308 135
MWAWGWAAAAL	PSQASGPE SVKPPEASGPE	PPGPPGP	RGLRLVGEAFH PSQRLVVEAFY	GRFTAPVSGIF GRFTAPVSAIF GQYRAPVAGFY	CLEAVSGLESN SLEAVSGLESN SLEAIMGLESS	FSGLLLGT FSGMLLGT TDLDNVWTVSE
	51	52 101 1	. 93 151 1	143 201 26	193 251 76	243 301 125
pTNF-x rTNF-x pTNF-y	ptne-x rtne-x ptne-y	ptne-x rtne-x ptne-y	pTNF-x rTNF-x pTNF-y	ptne-x rtne-x ptne-y	pTNF-x rTNF-x pTNF-y	pTNF-x rTNF-x pTNF-y

TLRL1 HU TLRL2 HU TLRL4 HU TLRL3 HU TLRL5 HU	MLSGVWFLSVLTVAGILQTESRKTAKDICKIRCLCEEKENVLNINMLQTLAFAVTSLVLSCAETIDYYGEICDNACPCEEKDGILTVSMELWLFLILSALISSTNADSDISVEICN-VCSCVSVENVLYVN MKPSIAEMLHRGRMLWIILLSTIALGWTTPIPLIEDSEEIDEPCFDPCYCEVKESLFHIHMKLWIHLFYSSLLACISLHSQTPVLSSRGSCDSLCNCEEKDGTMLIN * * * :::
TIRL1 HU TIRL2 HU TIRL4 HU TIRL3 HU TIRL5 HU	CENKGFTTVSLLQPPQYRIYQLFLNGNLLTRLYPNEFVNYSNAVTLHLGNNGLQEIRTGA CENRGIISLSEISPPRFPIYHLLLSGNLLNRLYPNEFVNYTGASILHLGSNVIQDIETGA CEKVSVYRPNQLKPPWSNFYHLNFQNNFLNILYPNTFLNFSHAVSLHLGNNKLQNIEGGA CDSKGFTNISQITEFWSRPFKLYLQRNSMRKLYTNSFLHLNNAVSINLGNNALQDIQTGA CEAKGIKMVSEISVPPSRPFQLSLLNNGLTMLHTNDFSGLTNAISIHLGFNNIADIEIGA *: * * * * * * * * * * * * * * * * * *
TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU	FSGLKTLKRLHLNNNKLEILREDTFLGLESLEYLQADYNYISAIEAGAFSKLNKLKVLIL FHGLRGLRRLHLNNNKLELLRDDTFLGLENLEYLQVDYNYISVIEPNAFGKLHLLQVLIL FLGLSALKQLHLNNNELKILRADTFLGIENLEYLQADYNLIKYIERGAFNKLHKLKVLIL FNGLKILKRLYLHENKLDVFRNDTFLGLESLEYLQADYNVIKRIESGAFRNLSKLRVLIL FNGLKILKQLHINHNSLEILKEDTFHGLENLEFLQADNNFITVIEPSAFSKLNRLKVLIL

FIG. 3A

TIRL1_HU TIRL2_HU TIRL4_HU TIRL3_HU TIRL3_HU	NDNLLLSLPSNVFRFVLLTHLDLRGNRLKVMPFAGVLEHIGG-IMEIQLEENPWNCTCDL NDNLLSSLPNNLFRFVPLTHLDLRGNRLKLLPYVGLLQHMDK-VVELQLEENPWNCSCEL NDNLISFLPDNIFRFASLTHLDIRGNRIQKLPYIGVLEHIGR-VVELQLEDNPWNCSCDL NDNLIPMLPTNLFKAVSLTHLDLRGNRLKVLFYRGMLDHIGRSLMELQLEENPWNCTCEI NDNLIPMLPTNLFRFVPLTHLDLRGNQLQTLPYVGFLEHIGR-ILDLQLEDNKWACNCDL *** : ***:** *************************
TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU TLRL5_HU	U ISLKAWLDTITVEVGEIVCETPERLHGKDVTQLTRQDLCPRKSASDSSQRGSHADTHV U ISLKDWLDSISYSALVGDVVCETPFRLHGRDLDEVSKQELCPRRLISDYEMRPQTPLSTT U LPLKAWLENMPYNIYIGEAICETPSDLYGRLLKETNKQELCPMGTGSDFDVR-ILPPSQL VQLKSWLERIPYTALVGDITCETPFHFHGKDLREIRKTELCPLLSDSEVEASLGIPHSSS U LQLKTWLENMPPQSIIGDVVCNSPPFFKGSILSRLKKESICPTPPVYEEHEDPSGS : ** **: : : * : : : : : : : : : : : :
TLRL1 HU TLRL2 HU TLRL4 HU TLRL3 HU TLRL5 HU	QRLSPTMNPALNPTRAPKASRPP-KMRNRPTPR-VTVSKDRQSF GYLHTTPASVNSVATSSSAVYKPPLKPPKGTRQPNKPRVRPTSRQPSKDLGYSNY ENGYTTPNGHTTQTSLHRLVTKPPKTTNPSKISGIVAGKALSNRNL SKENAWPTKPSSMLSSVHFTASSVEYKSSNKQPKPTKQPRTPRPPSTSQALYPGPNQ LHLAATSSINDSRMSTKTTSILKLPTKAPGL

FIG. 3B

SS-GLLIHCQERNIESLSDLRPPPQNPRKLILAG **::*: AVIQEGAFTNLTSLRRLYLNGNYLEVLYPSMFD ISMIQDRAFGDLTNLRRLYLNGNRIERLSPELFY ITVIKGDVFHNLTNLRRLYLNGNQIERLYPEIFS ISYVQDGAFINLPNLKSLFLNGNDIEKLTPGMFR IEVLEEGSFMNLTRLQKLYLNGNHLTKLSKGMFL * :: *:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*	<pre>IPYITKPSTQLPGPYCPIPCNCKVLSPS-GLLIHCQERNIESLSDLRPPPQNPRKLILAG</pre>	TLRL5_HU TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU TLRL3_HU
NGLNVNCQERKFTNISDLQPRPTSPKKLILG LGLNVNCQERKIESIAELQPKPYNPKKMYLTE LGLSVNCQEKNIQSMSELIPKPLNAKKLHVNG LGLTVNCKERGFNNISELLPRPLNAKKLYLSS -GLLIHCQERNIESLSDLRPPPQNPRKLILAG	OSC OIC HPG LSI	TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU TLRL5_HU

GLQSLQYLFLQYNLIREIQSGTFDPVPNLQLLFLNNNLLQAMPSGVFSGLTLLRLNLRSN GLHNLQYLYLEYNLIKEISAGTFDSMPNLQLLYLNNNLLKSLPVYIFSGAPLARLNLRNN GLQSLHYLYFEFNVIREIQPAAFSLMPNLKLLFLNNNLLRTLPTDAFAGTSLARLNLRKN GLHNLEYLYLEYNAIKEILPGTFNPMPKLKVLYLNNNLLQVLPPHIFSGVPLTKVNLKTN GLQSLQYLYLEYNVIKEIKPLTFDALINLQLLFLNNNLLRSLPDNIFGGTALTRLNLRNN **** * ****** * * ***

> TLRL2_HU TLRL4_HU TLRL3_HU TLRL5_HU

FIG. 3C

	•	
HFSHLPVKGVLDQLPAFIQIDLQENPWDCTCDIMGLKDWTEHANSPVIINEVTCESPAKH HFTSLPVSGVLDQLKSLIQIDLHDNPWDCTCDIVGMKLWVEQLKVGVLVDEVICKAPKKF KFMYLPVSGVLDQLQSLTQIDLEGNPWDCTCDLVALKLWVEKLSDGIVVKELKCETPVQF YFLYLPVAGVLEHINAIVQIDLNENPWDCTCDLVPFKQWIETISSVSVVGDVLCRSPENL QFTHLPVSNILDDLDLLTQIDLEDNPWDCSCDLVGLQQWIQKLSKNTVTDDILCTSPGHL * *** : * : * * * * * * * * * * * * * *	AGEILKFLGREAICPDSPNLSDGTVLSMNHNTDTPRSLSVSPSSYPELH J AETDMRSIKSELLCPDYSDVVVSTPTPSSIQVPARTSAVTPAVRLNSTGAPASLGAGGA J ANIELKSLKNEILCPKLLNKPSAPFTSPAPAITFTTPLGPIRSPPGG J THRDVRTIELEVLCPEMLHVAPAGESPAQPGDSHLIGAPTSASPYEFSPPG J DKKELKALNSEILCPGLVNNPSMPTQTSYLMVTTPATTTNTADTILRSLT	TEVPLSVLILGLLVVFILSVCFGAGLFVFVLKRR-KGVPSVPRNTNNLDVSSFQLQYGSY SSVPLSVLILSLLLVFIMSVFVAAGLFVLVMKRR-KKNQSDHTSTNNSDVSSFNMQYSVY -PVPLSILILSILVVLILTVFVAFCLLVFVLRRN-KKPTVKHEGLGNPDCGSMQLQLRKH GPVPLSVLILSLLVLFFSAVFVAAGLFAYVLRRRKKLPFRSKRQEGVDLTGIQMQCHRL DAVPLSVLILGLIMFITIVFCAAGIVVLVLHRR-RRYKKKQVDEQMRDNSPVHLQYSMY
TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU TLRL3_HU	TLRL1 HU TLRL2 HU TLRL4 HU TLRL3 HU TLRL5 HU	TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU

NTETHDKTDGHVYNYIPPPVGQMCQNPIYMQKEGDPVAYYR GGGGGTGGHPHAHVHHRGPALPKVKTPAGHVYEYIPHPLGHMCKNPIYRSREGNSVEDYK DHKTNKKDGLSTEAFIPQTIEQMSKSHTCGLKESETGFMFS FEDGGGGGGGGGGRPTLSSPEKAPPVGHVYEYIPHPVTQMCNNPIYKPREEEEVAVSS GHKTTHHTTERPSASLYEQHMVSPMVHVYRSPSFGPKHLEEEEERN	NLQELEEKKEEPDLATYSSNHHLQQQQQPPPPPQQPQQQ-DLHE	QATPREPELLYQNIAPPPQLQLQPGEEERRESHHLRSPAYSVSTIEPREDLLSPVQDADRFYRGIL
TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU TLRL5_HU	TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU	TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU

FIG. 3E

ERVKELPSAGLVHYNFCTLPKRQFAPSYESRRQNQDRINKTVLYGT EPDKHCSTTPAGNSLPEYPKFPCSPAAYTFSPNYDLRRPHQYLHPGAGDSRLREPVLYSP ESKKEYNSKSKKSLIGGN RPQPAPCTVGFVDCLYGTVPKLKELHVHPPGMQYPDLQQDARLKETLLFSA EKERELQQLGITEYLRKNIAQLQPDMEAHYPGAHEELKIMETLMYSR	PRKCFVGOS-KPNHPLLQAKPQSEPDYLEVLEKQTAISQL PSAVFVEPN-RNEYLELKAKLNVEPDYLEVLEKQTTFSQF HSKIVVEQR-KSEYFELKAKLQSSPDYLQVLEEQTALNKI EKGFTDHQTQKSDYLELRAKLQTKPDYLEVLEKTTYRF
TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU TLRL3_HU	TLRL1_HU TLRL2_HU TLRL4_HU TLRL3_HU

FIG 3F

LKSYSIDNISSLASDFPDFSYFKTSPMPSNRSYVVTVIY	r5685C6
LKSYSIENVTSIANNFPDFSYFRTFPMPSNKSYVVTFIY	p5685C6
GLQRLRIHTKARHPSRGQSLLIHSRREGSSLYKGWQTCMFISFLDVALFNGDSS GLKRLRINMEAKHPFPEQSLLIHSGGDSDSREKPMWLHKGWQPCMYISFLDMALFNRDSA **:****: **:** ***********************	r5685C6 p5685C6
CSCKSILPSAMEQTSYHGHLTIWFTDISTLGHVLKFTLVQDLKLSLCGSSTFPTKYLAIC	r5685C6
CNCKTVLPLAVERTSYNGHLTIWFTDTSALGHLLNFTLVQDLKLSLCSTNTLPTEYLAIC	p5685C6
MTSPSSFCLLLLQALGIVALGHFTKAQNN-TLIFTKGNTIKNCSCFVDIKDCDYSLANLL MAPPSRHCLLLISTLGVFALNCFTKGQKNSTLIFTRENTIRNCSCSADIRDCDYSLANLM *; ** .****; **************************	r5685C6 p5685C6

FIG. 4

117		D7.2
150	101 GMKQVQCTGSNERAKAYLLGTSGVLFILTGIFVLIPVSWTANIIRDFYN	D17
150	101 GMKCTRCTGDNEKVKAHILLTAGINLIITGMVGANPVNLVSNAIIRDFFT	D8
149	101 GMRCTVFCQES-RAKDRVAVAGGVFFILGGLLGFIPVAWNLHGILRDFYS 149	D2
	* * * * * *	
86		D7.2
100	51 WMNCIRQARVRLQCKFYSSLLALPPALETARALMCVAVALSLIALLIGIC	D17
100	51 WMNCVRQANIRMQCKIYDSLLALSPDLQAARGLMCAASVMSFLAFMMAIL 100	D8
100	51 WMECATHSTGITQCDIYSTLLGLPADIQGAQAMMVTSSAISSLACIISVV 100	D2
	**	
49	H	D7.2
50	1 MAFYPLQIAGLVLGFLGMVGTLATTLLPQWRVSAFVGSNIIVFERLWEGL	D17
20	1 MATHALEIAGLFLGGVGMVGTVAVTVMPQWRVSAFIENNIVVFENFWEGL	D8
20	1 MASLGLQLVGYILGLLGLLGTLVAMLLPSWKTSSYVGASIVTAVGFSKGL 50	D2

FIG. 5A

199 200 200 130					
150 PLVPDSMKFEIGEALYLGIISSLFSLIAGIILCFSCSSQRNRSNYYDAYQ 199 151 PIVNVAQKRELGEALYLGWTTALVLIVGGALFCCVFCCNEKSSSYRYSIP 200 151 PAIHIGQKRELGAALFLGWASAAVLFIGGGLLCGFCCCNRKKQGYRYPVP 200 118 SHLHQGGIGG	*	200 AQPLATRSSPRAGQPPKVKSEFNSYSLTGYV 230	SHRTTQKSYHTGKKSPSVYSRSQYV 225	GYRVPHTDKRRNTTMLSKTSTSYV 224	130
150 151 151 118		200	201	201	131
D2 D8 D17 D7.2		D2	D8	D17	D7.2

FIG. 5B

•	-1	1 MESLATUTEMFIFEVIVUGEVIFGEENARMINSCLARSENSALIKA 40	_
<i>r</i> >	Н	MEANHCSLGVYPSYPDLVIDVGEVTLGEENRKKLQKTQRDQ-ERARVIRA 49	_
_	Н	MNISVDLETNYAELVLDVGRVTLGENSRKKMKDCKLRKKQNERVSRA 47	_
r-7	Н	MSLRIDVDTNFPECVVDAGKVTLGTQQRQEMDPRLREK-QNEIILRA 46	
r	Н	2KIQRDQ-EKERVMRA 4	0
	•	**	
~	49	49 ICALLNSGGGVIKAEIDDKTYSYQCHGLGQDLETSFQKLLPS-GSQKYLD 97	_
<i>r</i> \	50	ACALLNSGGGVIQMEMANRDERPTEMGLDLEESLRKLIQYPYLQAFFE 9'	~
_	48	MCALLNSGGGVIKAEIENEDYSYTKDGIGLDLENSFSNILLF-VP-EYLD 95	. ~
r-3	47	VCALLNSGGGIIKAEIENKGYNYERHGVGLDVPPIFRSHLD 8.	_
r	50	ACALLNSGGGVIRMAKKVEHPVEMGLDLEQSLRELIQSSDLQAFFE 95	
		· * * * * * * * * * * * * * * * * * * *	
m	86	YMQQGHNLLIFVKSWSPDVFSLPLRICSLRSNLYRRDVTSAINLSA 143	~
<i>r</i> >	86	IKQHGRCFYIFVKSWSGDPFLKDGSFNSRICSLSSSLYCRSGTSVLHMNS 14'	_
\sim	96	FMQNGNYFLIFVKSWSLNTSGLRITTLSSNLYKRDITSAKVMNA 139	~
F-7	88	KMOKENHFLI FVKSWNTEAGVPLATLCSNLYHRERTSTDVMDS	$\overline{}$
۲• ·	96	TKQQGRCFYLFVKSWSSGPFPEDRSVKPRLCSLSSSLYRRSETSVRSMDS 145	

FIG. 6A

140 TAALEFLKDMKKTRGRLYLRPELLAKRPCVDIQEENNMKALA 181 131 QEALAFLKCRTQTPTNINVSNSLGPQAAQGSVQYEGNINVSA 172 146 REAFCFLKTKRKPKILEEG-PFHKIHKGVYQELPNSDPADPNSDPA 190 * * * 188 SEFFKKDKLMYKEKLNFTESTHVEFKRFTTKKVIPRIKEMLPHYVSAFAN 237 190 YEVFQTDTIEYGEILSFPESPSIEFKQFSTKHIQQYVENIIPEYISAFAN 239 182 GVFFDRTELDRKEKLTFTESTHVEIKNFSTEKLLQRIKEILPQYVSAFAN 231 173 AALFDRKRLQYLEKLNLPESTHVEFVMFST-DVSHCVKDRLPKCVSAFAN 221 191 DLIFQKDYLEYGEILPFPESQLVEFKQFSTKHFQEYVKRTIPEYVPAFAN 240
148 RQAFDFLKTKER-QŠKYNLINEGSPPSKIMKAVYQNISESNPA 189 140 TAALEFLKDMKKTRGRLYLRPELLAKRPCVDIQEENNMKALA 181 131 QEALAFLKCRTQTPTNINVSNSLGPQAAQGSVQYEGNINVSA 172 146 REAFCFLKTKRKPKILEEG-PFHKIHKGVYQELPNSDPADPNSDPA 190 * * * 188 SEFFKKDKLMYKEKLNFTESTHVEFKRFTTKKVIPRIKEMLPHYVSAFAN 237 190 YEVFQTDTIEYGEILSFPESPSIEFKQFSTKHIQQYVENIIPEYISAFAN 233 173 AALFDRKELQYLEKLNLPESTHVEFKNFST-DVSHCVKDRLPKCVSAFAN 221 173 AALFDRKRLQYLEKLNLPESTHVEFKQFSTKHFQEYVKRTIPEYVPAFAN 240

289 280 271 290

TEGGYLFIGVDDKSRKVLGCAKEQVDPDSLKNVIARAISKLPIVHFCSSK TDGGYLFIGLNED-KEIIGFKAEMSDLDDLEREIEKSIRKMPVHHFCMEK

240 232 222

日日日日日

TQGGYVLIGVDDKSKEVVGCKWEKVNPDLLKKEIENCIEKLPTFHFCCEK

TGGGYLFXGVDDKSREVLGCAKENXDPDSLRXKIEXAIYKLPCXHFCQPQ

TEGGYVFFGVHDETCQVIGCEKEKIDLTSLRASIDGCIKKLPVHHFCTQR

'n	Zaa	Z88 FRVNFTTENVIQEDGIVCVIQVEFFCCVVFAEAFDSWIMEDNSVT 337	33.1
F)	290	290 PRVEYSTKIVEVFCGKELYGYLCVIKVKAFCCVVFSEAPKSWMVREKYIR 339	339
_	281	KKINYSCKFLGVYDKGSLCGYVCALRVERFCCAVFAKEPDSWHVKDNRVM	330
רביז	272	PEIKYVINFLEVHDKGALRGYVCAIKVEKFCCAVFAKVPSSWQVKDNRVR	321
r.,	291	RPITFTLKIVDVLKRGELYGYACMIRVNPFCCAVFSEAPNSWIVEDKYVC	340
		· · · · · · · · · · · · · · · · · · ·	
~	338	338 RLTAEQWVVMMLDTQ	352
<i>r</i> \	340	PLTTEEWVEKMMDADPEFPPDFAEAFESQLSLSDSPSLCRPVYSKKGLEH	389
$\overline{}$	331	QLTRKEWIQFMVEAEPKFS-SSYEEVISQINTSLPAPHSWPLLEW	374
(-1	322	QLPTREWTAWMMEADPDLSRCPEMVLQLSLSSATPRSKPVCIHKNSEC 369	369
F+ -	341	341 SLITEKWVGMMIDIDPDLL-QLSEDFECQLSLSSGPPLSRPVYSKKGLEH 389	389
		* * * *	
m	353	SGKGK	357
r)	390	390 KADLQQHLFPVPPGHLECTPESLWKELSLQHEGLKELIHKQMRPFSQGIV	439
\sim	375	375 QRQRHHCPGLSGRITYTPENLCRKLFLQHEGLKQLICEEMDSVRKGSL	422
r-1	370	LKEQQKRYFPVFSDRVVYTPESLYKELFSQHKGLRDLINTEMRPFSQGIL 419	419
Γ÷.	390	390 KKELQQLLFSVPPGYLRYTPESLWRDLISEHRGLEELINKQMQPFFRGIV 439	439

FIG. 6C

m U D	358 440 423	358 440 ILSRSWAVDLNLQEKPGVICDALLIAQNSTPILYTILREQDAEGQDYCTR 423 IFSRSWSVDLGLQENHKVLCDALLISQDSPPVLYTFHMVQDEEFKGYSTQ	357 489 472
H	420		469
Ŀ	440	440 ILSRSWAVDLNLQEKPGVICDALLIAQNSTPILYTILREQDAEGQDYCTR	489
Д	358		357
ບ	490	490 TAFTLKQKLVNMGGYTGKVCVRAKVLCLSPESSAEALEAAVSPMDYPASY 539	539
Д	473	TALTLKOKLAKIGGYTKKVCVMTKIFYLSPEG	504
ഥ	470	470 VAYSLKQKLVNKGGYTGRLCITPLVCVLNSDRKAQSVYSSY-LQIYPESY	518
ᄪ	490	490 TAFTLKQKLVNMGGYTGKVCVRAKVLCLSPESSAEALEAAVSPMDYPASY	539
В	358		357
ပ	540	SLAGTQHMEALLQSLVIVLLGFRSLLSDQLGCEVLNLLTAQQYEIFSRSL	589
Д	505	MISCOYDLRSQVI	517
闰	519	519 NEMTPQHMEALLQSLVIVLLGFKSFLSEELGSEVLNLLTNKQYELLSKNL	568
۲ı	540	540 SLAGTOHMEALLOSLVIVLLGFRSLLSDQLGCEVLNLLTAQQYEIFSRSL 589	589

FIG. 6D

358 358 590 590 590 640 640 640 640 640 640 640 640 640 64
358 590 RKNRELFVHGLPGSGKTIMAMKIMEKIRNVFHCEAHRILYVCENQPLRNF 518 YPESYYFTRRKYLLKALFKALKRLKSLRDQFSFAENLYQIIG 569 RKTRELFVHGLPGSGKTILALRIMEKIRNVFHCEPANILYICENQPLKKL 590 RKNRELFVHGLPGSGKTIMAMKIMEKIRNVFHCEAHRILYVCENQPLRNF 640 ISDRNICRAETRETFLREKFEHIQHIVIDEAQNFRTEDGDWYRKAKTI 560IDCFQKNDKKMFKSCRRL 640 ISDRNICRAETRKTFLRENFEHIQHIVIDEAQNFRTEDGDWYGKAKFI 640 ISDRNICRAETRKTFLRENFEHIQHIVIDEAQNFRTEDGDWYGKAKFI 640 ISDRNICRAETRKTFLRENFEHIQHIVIDEAQNFRTEDGDWYGKAKFI 640 ISDRNICRAETRKTFLRENFEHIGHSGLPPLSAQYPREELTRVVRNAGPI 578 T 669 TRQQRDGPGVLWIFLDYFQTYHLSCSGLPPPSDQYPREEINRVVRNAGPI

FIG. 6E

357 787 578 768 748	357 833 578 818 748	357 877 578 868	
35 AEYIQQEMQLIIENPPINIPHGYLAILSEAKWVPGVPGNTKIIKNFTLEQ 78 ANYLQQVMQEARQNPPPNLPPGSLVMLYEPKWAQGVPGNLEIIEDLNLEE 76 AKYLQKENASN	ŎST	35 MVVQLSDACDMLGVHIVLDSVRRFSGLERSIVFGIHPRTADPAI 87 57 LHEESDLLLQIGDASDVLTDHIVLDSVCRFSGLERNIVFGINPGVAPPAG 86	FIG. 6F
SYLAILS	AVLVST AVLFTK	VLDSVR.	357 897 578 3V 891 748
•		• • •	LPNILICLASRAKQHLYIFL AYNLLLCLASRAKRHLYILKASV
358 738 579 719	358 788 579 769 749	358 834 579 819 749	358 878 579 869 749
日の口は下	шпопы	- 百CDロ 〒	поопы

1

SEQUENCE LISTING

<110>	Schering Corporation	
<120>	MAMMALIAN GENES; RELATED REAGENTS AND METHODS	
<130>	DX01169K	
<150>	60/231,267	
<151>	2000-09-08	
<160>	53	
<170>	PatentIn version 3.1	
<210>	1	
<211>	704	
<212>	DNA	
<213>	Homo sapiens	
<400> atggcg	1 gggc ccgagegctg gggccccctg ctcctgtgcc tgctgcaggc cgctccaggg	60
aggccc	egtc tggcccctcc ccagaatgtg acgctgctct cccagaactt cagcgtgtac	120
ctgaca	tggc tcccagggct tggcaacccc caggatgtga cctattttgt ggcctatcag	180
agctct	ccca cccgtagacg gtggcgcgaa gtggaagagt gtgcgggaac caaggagctg	240
ctatgt	tota tgatgtgcct gaagaaacag gacctgtaca acaagttcaa gggacgcgtg	300
cggacg	gttt ctcccagctc caagtccccc tgggtggagt ccgaatacct ggattacctt	360
tttgaa	gtgg agccggcccc acctgtcctg gtgctcaccc agacggagga gatcctgagt	420
gccaat	gcca cgtaccaget geceeetge atgeceecae tggatetgaa gtatgaggtg	480
gcattc	tgga aggaggggc cggaaacaag gtgggaagct cctttcctgc ccccaggcta	540
ggcccg	getee tecacecett ettactcagg ttetteteac ceteccagee tgeteetgea	600

2

cccctcctcc aggaagtctt ccctgtacac tcctgaactt ctggcagtca gccctaataa 660 aatctgatca aagtaaaaaa aaaaaaaag ggcggccgcc gact 704

<210> 2

<211> 211

<212> PRT

<213> Homo sapiens

. <400> 2

Met Ala Gly Pro Glu Arg Trp Gly Pro Leu Leu Cys Leu Leu Gln 1 5 10 15

Ala Ala Pro Gly Arg Pro Arg Leu Ala Pro Pro Gln Asn Val Thr Leu 20 25 30

Leu Ser Gln Asn Phe Ser Val Tyr Leu Thr Trp Leu Pro Gly Leu Gly 35 40 45

Asn Pro Gln Asp Val Thr Tyr Phe Val Ala Tyr Gln Ser Ser Pro Thr 50 55 60

Arg Arg Arg Trp Arg Glu Val Glu Glu Cys Ala Gly Thr Lys Glu Leu 65 70 75 80

Leu Cys Ser Met Met Cys Leu Lys Lys Gln Asp Leu Tyr Asn Lys Phe 85 90 95

Lys Gly Arg Val Arg Thr Val Ser Pro Ser Ser Lys Ser Pro Trp Val

Glu Ser Glu Tyr Leu Asp Tyr Leu Phe Glu Val Glu Pro Ala Pro Pro 115 120 125

Val Leu Val Leu Thr Gln Thr Glu Glu Ile Leu Ser Ala Asn Ala Thr 130 135 140

Tyr Gln Leu Pro Pro Cys Met Pro Pro Leu Asp Leu Lys Tyr Glu Val 145 150 155 160

Ala Phe Trp Lys Glu Gly Ala Gly Asn Lys Val Gly Ser Ser Phe Pro 165 170 175

3

Ala Pro Arg Leu Gly Pro Leu Leu His Pro Phe Leu Leu Arg Phe Phe 180 185 190

Ser Pro Ser Gln Pro Ala Pro Ala Pro Leu Gln Glu Val Phe Pro 195 200 205

Val His Ser 210

<210> 3

<211> 295

<212> PRT

<213> Homo sapiens

<400> 3

Met Glu Thr Pro Ala Trp Pro Arg Val Pro Arg Pro Glu Thr Ala Val 1 5 10 15

Ala Arg Thr Leu Leu Gly Trp Val Phe Ala Gln Val Ala Gly Ala 20 25 30

Ser Gly Thr Thr Asn Thr Val Ala Ala Tyr Asn Leu Thr Trp Lys Ser 35 40 45

Thr Asn Phe Lys Thr Ile Leu Glu Trp Glu Pro Lys Pro Val Asn Gln 50 55 60

Val Tyr Thr Val Gln Ile Ser Thr Lys Ser Gly Asp Trp Lys Ser Lys 65 70 75 80

Cys Phe Tyr Thr Thr Asp Thr Glu Cys Asp Leu Thr Asp Glu Ile Val 85 90 95

Lys Asp Val Lys Gln Thr Tyr Leu Ala Arg Val Phe Ser Tyr Pro Ala 100 105 110

Gly Asn Val Glu Ser Thr Gly Ser Ala Gly Glu Pro Leu Tyr Glu Asn 115 120 125

Ser Pro Glu Phe Thr Pro Tyr Leu Glu Thr Asn Leu Gly Gln Pro Thr

4

130 135 140

Ile Gln Ser Phe Glu Gln Val Gly Thr Lys Val Asn Val Thr Val Glu
145 150 155 160

Asp Glu Arg Thr Leu Val Arg Arg Asn Asn Thr Phe Leu Ser Leu Arg 165 170 175

Asp Val Phe Gly Lys Asp Leu Ile Tyr Thr Leu Tyr Trp Lys Ser 180 185 190

Ser Ser Ser Gly Lys Lys Thr Ala Lys Thr Asn Thr Asn Glu Phe Leu 195 200 205

Ile Asp Val Asp Lys Gly Glu Asn Tyr Cys Phe Ser Val Gln Ala Val 210 215 220

Ile Pro Ser Arg Thr Val Asn Arg Lys Ser Thr Asp Ser Pro Val Glu 225 230 235 240

Cys Met Gly Gln Glu Lys Gly Glu Phe Arg Glu Ile Phe Tyr Ile Ile 245 250 255

Gly Ala Val Ala Phe Val Val Ile Ile Leu Val Ile Ile Leu Ala Ile 260 265 270

Ser Leu His Lys Cys Arg Lys Ala Gly Val Gly Gln Ser Trp Lys Glu 275 280 285

Asn Ser Pro Leu Asn Val Ser 290 295

<210> 4

<211> 515

<212> PRT

<213> Homo sapiens

<400> 4

Met Leu Leu Ser Gln Asn Ala Phe Ile Phe Arg Ser Leu Asn Leu Val 1 5 10 15

5

Leu Met Val Tyr Ile Ser Leu Val Phe Gly Ile Ser Tyr Asp Ser Pro 20 25 30

Asp Tyr Thr Asp Glu Ser Cys Thr Phe Lys Ile Ser Leu Arg Asn Phe 35 40 45

Arg Ser Ile Leu Ser Trp Glu Leu Lys Asn His Ser Ile Val Pro Thr 50 55 60

His Tyr Thr Leu Leu Tyr Thr Ile Met Ser Lys Pro Glu Asp Leu Lys 65 70 75 80

Val Val Lys Asn Cys Ala Asn Thr Thr Arg Ser Phe Cys Asp Leu Thr 85 90 95

Asp Glu Trp Arg Ser Thr His Glu Ala Tyr Val Thr Val Leu Glu Gly
100 105 110

Phe Ser Gly Asn Thr Thr Leu Phe Ser Cys Ser His Asn Phe Trp Leu 115 120 125

Ala Ile Asp Met Ser Phe Glu Pro Pro Glu Phe Glu Ile Val Gly Phe 130 135 140

Thr Asn His Ile Asn Val Val Val Lys Phe Pro Ser Ile Val Glu 145 150 155 160

Glu Leu Gln Phe Asp Leu Ser Leu Val Ile Glu Glu Gln Ser Glu Gly 165 170 175

Ile Val Lys Lys His Lys Pro Glu Ile Lys Gly Asn Met Ser Gly Asn 180 185 190

Phe Thr Tyr Ile Ile Asp Lys Leu Ile Pro Asn Thr Asn Tyr Cys Val 195 200 205

Ser Val Tyr Leu Glu His Ser Asp Glu Gln Ala Val Ile Lys Ser Pro 210 215 220

Leu Lys Cys Thr Leu Leu Pro Pro Gly Gln Glu Ser Glu Ser Ala Glu 225 230 235 240

Ser Ala Lys Ile Gly Gly Ile Ile Thr Val Phe Leu Ile Ala Leu Val 245 250 255 Leu Thr Ser Thr Ile Val Thr Leu Lys Trp Ile Gly Tyr Ile Cys Leu 260 265 270

Arg Asn Ser Leu Pro Lys Val Leu Asn Phe His Asn Phe Leu Ala Trp 275 280 285

Pro Phe Pro Asn Leu Pro Pro Leu Glu Ala Met Asp Met Val Glu Val 290 295 300

Ile Tyr Ile Asn Arg Lys Lys Lys Val Trp Asp Tyr Asn Tyr Asp Asp 305 310 315

Glu Ser Asp Ser Asp Thr Glu Ala Ala Pro Arg Thr Ser Gly Gly Gly 325 330 335

Tyr Thr Met His Gly Leu Thr Val Arg Pro Leu Gly Gln Ala Ser Ala 340 345 350

Thr Ser Thr Glu Ser Gln Leu Ile Asp Pro Glu Ser Glu Glu Glu Pro 355 360 365

Asp Leu Pro Glu Val Asp Val Glu Leu Pro Thr Met Pro Lys Asp Ser 370 380

Pro Gln Gln Leu Glu Leu Leu Ser Gly Pro Cys Glu Arg Arg Lys Ser 385 390 395

Pro Leu Gln Asp Pro Phe Pro Glu Glu Asp Tyr Ser Ser Thr Glu Gly
405 410 415

Ser Gly Gly Arg Ile Thr Phe Asn Val Asp Leu Asn Ser Val Phe Leu 420 425 430

Arg Val Leu Asp Asp Glu Asp Ser Asp Asp Leu Glu Ala Pro Leu Met 435 440 445

Leu Ser Ser His Leu Glu Glu Met Val Asp Pro Glu Asp Pro Asp Asn 450 . 455 . 460

Val Gln Ser Asn His Leu Leu Ala Ser Gly Glu Gly Thr Gln Pro Thr 465 470 475 480

Phe Pro Ser Pro Ser Ser Glu Gly Leu Trp Ser Glu Asp Ala Pro Ser 485 490 495

WO 02/20569

7

Asp Gln Ser Asp Thr Ser Glu Ser Asp Val Asp Leu Gly Asp Gly Tyr 500 505 510

Ile Met Arg . 515

<210> 5

<211> 325

<212> PRT

<213> Homo sapiens

<400> 5

Met Ala Trp Ser Leu Gly Ser Trp Leu Gly Gly Cys Leu Leu Val Ser 1 5 10 15

Ala Leu Gly Met Val Pro Pro Pro Glu Asn Val Arg Met Asn Ser Val 20 25 30

Asn Phe Lys Asn Ile Leu Gln Trp Glu Ser Pro Ala Phe Ala Lys Gly 35 40 45

Asn Leu Thr Phe Thr Ala Gln Tyr Leu Ser Tyr Arg Ile Phe Gln Asp 50 55 60

Lys Cys Met Asn Thr Thr Leu Thr Glu Cys Asp Phe Ser Ser Leu Ser 65 70 75 80

Lys Tyr Gly Asp His Thr Leu Arg Val Arg Ala Glu Phe Ala Asp Glu 85 90 95

His Ser Asp Trp Val Asn Ile Thr Phe Cys Pro Val Asp Asp Thr Ile 100 105 110

Ile Gly Pro Pro Gly Met Gln Val Glu Val Leu Ala Asp Ser Leu His 115 120 125

Met Arg Phe Leu Ala Pro Lys Ile Glu Asn Glu Tyr Glu Thr Trp Thr 130 135 140

Met Lys Asn Val Tyr Asn Ser Trp Thr Tyr Asn Val Gln Tyr Trp Lys

8

160 155 150 145 Asn Gly Thr Asp Glu Lys Phe Gln Ile Thr Pro Gln Tyr Asp Phe Glu 170 175 165 Val Leu Arg Asn Leu Glu Pro Trp Thr Thr Tyr Cys Val Gln Val Arg 180 185 Gly Phe Leu Pro Asp Arg Asn Lys Ala Gly Glu Trp Ser Glu Pro Val 195 200 Cys Glu Gln Thr Thr His Asp Glu Thr Val Pro Ser Trp Met Val Ala Val Ile Leu Met Ala Ser Val Phe Met Val Cys Leu Ala Leu Leu Gly 230 235 240 Cys Phe Ser Leu Leu Trp Cys Val Tyr Lys Lys Thr Lys Tyr Ala Phe 255 245 250 Ser Pro Arg Asn Ser Leu Pro Gln His Leu Lys Glu Phe Leu Gly His 265 260 Pro His His Asn Thr Leu Leu Phe Phe Ser Phe Pro Leu Ser Asp Glu 275 280

Asn Asp Val Phe Asp Lys Leu Ser Val Ile Ala Glu Asp Ser Glu Ser 290 295 300

Gly Lys Gln Asn Pro Gly Asp Ser Cys Ser Leu Gly Thr Pro Pro Gly 305 310 315 320

Gln Gly Pro Gln Ser 325

<210> 6 .

<211> 231.

<212> PRT

<213> Homo sapiens

<400> 6

9

Met Met Pro Lys His Cys Phe Leu Gly Phe Leu Ile Ser Phe Phe Leu 1 5 10 15

Thr Gly Val Ala Gly Thr Gln Ser Thr His Glu Ser Leu Lys Pro Gln 20 25 30

Arg Val Gln Phe Gln Ser Arg Asn Phe His Asn Ile Leu Gln Trp Gln 35 40 45

Pro Gly Arg Ala Leu Thr Gly Asn Ser Ser Val Tyr Phe Val Gln Tyr 50 55 60

Lys Ile Tyr Gly Gln Arg Gln Trp Lys Asn Lys Glu Asp Cys Trp Gly 65 70 75 80

Thr Gln Glu Leu Ser Cys Asp Leu Thr Ser Glu Thr Ser Asp Ile Gln 85 90 95

Glu Pro Tyr Tyr Gly Arg Val Arg Ala Ala Ser Ala Gly Ser Tyr Ser 100 105 110

Glu Trp Ser Met Thr Pro Arg Phe Thr Pro Trp Trp Glu Thr Lys Ile 115 120 125

Asp Pro Pro Val Met Asn Ile Thr Gln Val Asn Gly Ser Leu Leu Val 130 135 140

Ile Leu His Ala Pro Asn Leu Pro Tyr Arg Tyr Gln Lys Glu Lys Asn 145 150 155 160

Val Ser Ile Glu Asp Tyr Tyr Glu Leu Leu Tyr Arg Val Phe Ile Ile 165 170 175

Asn Asn Ser Leu Glu Lys Glu Gln Lys Val Tyr Glu Gly Ala His Arg 180 185 190

Ala Val Glu Ile Glu Ala Leu Thr Pro His Ser Ser Tyr Cys Val Val 195 200 205

Ala Glu Ile Tyr Gln Pro Met Leu Asp Arg Arg Ser Gln Arg Ser Glu 210 215 220

Glu Arg Cys Val Glu Ile Pro

<210> 7

<211> 553

<212> PRT

<213> Homo sapiens

<220>

<221> MISC_FEATURE

<222> (522)..(522)

<223> unknown amino

<400> 7

Met Arg Ala Pro Gly Arg Pro Ala Leu Arg Pro Leu Pro Leu Pro Pro 1 5 10 15

Leu Leu Leu Leu Leu Ala Ala Pro Trp Gly Arg Ala Val Pro Cys 20 25 30

Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile 35 40 45

Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly . 50 55 60

Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys 65 70 75 80

Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp 85 90 95

Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val 100 105 110

Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg 115 120 125

Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu Val Ala Leu 130 . 135 140

11

Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala Pro Glu Lys 145 150 155 160

Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln Gln Ile Tyr 165 170 175

Ser Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys Ser Asn Arg 180 185 190

Thr Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu Thr Trp Leu 195 200 205

Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe Val Pro Gly 210 215 220

Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala Arg Thr Leu 225 230 235 240

Lys Asp Gln Ser Ser Glu Phe Lys Ala Lys Ile Ile Phe Trp Tyr Val 245 250 255

Leu Pro Ile Ser Ile Thr Val Phe Leu Phe Ser Val Met Gly Tyr Ser 260 265 270

Ile Tyr Arg Tyr Ile His Val Gly Lys Glu Lys His Pro Ala Asn Leu 275 280 285

Ile Leu Ile Tyr Gly Asn Glu Phe Asp Lys Arg Phe Phe Val Pro Ala 290 295 300

Glu Lys Ile Val Ile Asn Phe Ile Thr Leu Asn Ile Ser Asp Asp Ser 305 310 315 320

Lys Ile Ser His Gln Asp Met Ser Leu Leu Gly Lys Ser Ser Asp Val 325 330 335

Ser Ser Leu Asn Asp Pro Gln Pro Ser Gly Asn Leu Arg Pro Pro Gln 340 345 350

Glu Glu Glu Val Lys His Leu Gly Tyr Ala Ser His Leu Met Glu 355 360 365

Ile Phe Cys Asp Ser Glu Glu Asn Thr Glu Gly Thr Ser Leu Thr Gln 370 380

12

Gln Glu Ser Leu Ser Arg Thr Ile Pro Pro Asp Lys Thr Val Ile Glu 385 390 395 400

Tyr Glu Tyr Asp Val Arg Thr Thr Asp Ile Cys Ala Gly Pro Glu Glu
405 410 415

Gln Glu Leu Ser Leu Gln Glu Glu Val Ser Thr Gln Gly Thr Leu Leu 420 425 430

Glu Ser Gln Ala Ala Leu Ala Val Leu Gly Pro Gln Thr Leu Gln Tyr 435 440 445

Ser Tyr Thr Pro Gln Leu Gln Asp Leu Asp Pro Leu Ala Gln Glu His
450 455 460

Thr Asp Ser Glu Glu Gly Pro Glu Glu Glu Pro Ser Thr Thr Leu Val 465 470 475 480

Asp Trp Asp Pro Gln Thr Gly Arg Leu Cys Ile Pro Ser Leu Ser Ser 485 490 495

Phe Asp Gln Asp Ser Glu Gly Cys Glu Pro Ser Glu Gly Asp Gly Leu 500 505 510

Gly Glu Glu Gly Leu Leu Ser Arg Leu Xaa Glu Glu Pro Ala Pro Asp 515 520 525

Arg Pro Pro Gly Glu Asn Glu Thr Tyr Leu Met Gln Phe Met Glu Glu 530 535 540

Trp Gly Leu Tyr Val Gln Met Glu Asn 545 550

<210> 8

<211> 687

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

13

<222> (1)..(684)

<223>

<400 atg	gct	gaa	ctt	tgt	ccg	gcg	gcc	gga	cga	cgg	cgc	ctt	aag	gaa	gcg	48
Met 1	Ala	Glu	Leu	Cys 5	Pro	Ala	Ala	GŢĀ	Arg 10	Arg	Arg	Leu	ГÀЗ	GIU 15	Ala	
gtg Val	cgg Arg	aag Lys	cag Gln 20	gga Gly	caa Gln	gaa Glu	gcc Ala	gcg Ala 25	gga Gly	tct Ser	ctt Leu	cgg Arg	tcc Ser 30	ccc Pro	agg Arg	96
acc Thr	tcc Ser	agg Arg 35	tgc Cys	aga Arg	agt Ser	gac Asp	cgc Arg 40	gga Gly	gac Asp	tct Ser	gct Ala	tca Ser 45	cga Arg	gtt Val	tca Ser	144
gga Gly	gct Ala 50	gct Ala	gaa Glu	aga Arg	ggc Gly	cac His 55	gga Gly	gcg Ala	ccg Pro	gtt Val	ctc Leu 60	agg Arg \	gct Ala	tct Ser	gga Gly	192
ccc Pro 65	gct Ala	gct Ala	gcc Ala	cca Pro	999 Gly 70	gcg Ala	ggc Gly	ctg Leu	cgg Arg	ctg Leu 75	gtg Val	ggc Gly	gag Glu	gcc Ala	ttt Phe 80	240
cac His	tgc Cys	cgg Arg	ctg Leu	cag Gln 85	ggt Gly	ccc Pro	cgc Arg	cgg Arg	gtg Val 90	gac Asp	aag Lys	cgg Arg	acg Thr	ctg Leu 95	gtg Val	288
gag Glu	ctg Leu	cat His	ggt Gly 100	ttc Phe	cag Gln	gct Ala	cct Pro	gct Ala 105	gcc Ala	caa Gln	ggt Gly	gcc Ala	ttc Phe 110	ctg Leu	cga Arg	336
ggc Gly	tcc Ser	ggt Gly 115	ctg Leu	agc Ser	ctg Leu	gcc Ala	tcg Ser 120	ggt Gly	cgg Arg	ttc Phe	acg Thr	gcc Ala 125	ccc Pro	gtg Val	tcc Ser	384
ggc Gly	atc Ile 130	ttc Phe	cag Gln	ttc Phe	tct Ser	gcc Ala 135	agt Ser	ctg Leu	cac His	gtg Val	gac Asp 140	cac His	agt Ser	gag Glu	ctg Leu	432
cag Gln 145	ggc Gly	aag Lys	gcc Ala	cgg Arg	ctg Leu 150	cgg Arg	gcc Ala	cgg Arg	gac Asp	gtg Val 155	gtg Val	tgt Cys	gtt Val	ctc Leu	atc Ile 160	480
tgt Cys	att Ile	gag Glu	.tcc Ser	ctg Leu 165	Cys	Gln	Arg	cac His	Thr	Cys	Leu	Glu	gcc Ala	gtc Val 175	Ser	528
ggc Gly	ctg Leu	gag Glu	agc Ser 180	Asn	agc Ser	agg Arg	gtc Val	ttc Phe 185	Thr	cta Leu	cag Gln	gtg Val	cag Gln 190	Gly ggg	ctg Leu	576
ctg Leu	cag Gln	ctg Leu 195	Gln	gct Ala	gga Gly	cag Gln	tac Tyr 200	Ala	tct Ser	gtg Val	ttt Phe	gtg Val 205	Asp	aat Asn	ggc Gly	624

14

tcc ggg gcc gtc ctc acc atc cag gcg ggc tcc agc ttc tcc ggg ctg

Ser Gly Ala Val Leu Thr Ile Gln Ala Gly Ser Ser Phe Ser Gly Leu

210

215

220

ctc ctg ggc acg tga Leu Leu Gly Thr 225 687

<210> 9

<211> 228

<212> PRT

<213> Homo sapiens

<400> 9

Met Ala Glu Leu Cys Pro Ala Ala Gly Arg Arg Arg Leu Lys Glu Ala 1 5 10 15

Val Arg Lys Gln Gly Gln Glu Ala Ala Gly Ser Leu Arg Ser Pro Arg 20 25 30

Thr Ser Arg Cys Arg Ser Asp Arg Gly Asp Ser Ala Ser Arg Val Ser 35 40 45

Gly Ala Ala Glu Arg Gly His Gly Ala Pro Val Leu Arg Ala Ser Gly 50 55 60

Pro Ala Ala Ala Pro Gly Ala Gly Leu Arg Leu Val Gly Glu Ala Phe 65 70 75 80

His Cys Arg Leu Gln Gly Pro Arg Arg Val Asp Lys Arg Thr Leu Val 85 90 95

Glu Leu His Gly Phe Gln Ala Pro Ala Ala Gln Gly Ala Phe Leu Arg 100 105 110

Gly Ser Gly Leu Ser Leu Ala Ser Gly Arg Phe Thr Ala Pro Val Ser 115 120 125

Gly Ile Phe Gln Phe Ser Ala Ser Leu His Val Asp His Ser Glu Leu 130 135 140

Gln Gly Lys Ala Arg Leu Arg Ala Arg Asp Val Val Cys Val Leu Ile 145 150 155 160

15

Cys Ile Glu Ser Leu Cys Gln Arg His Thr Cys Leu Glu Ala Val Ser 170 Gly Leu Glu Ser Asn Ser Arg Val Phe Thr Leu Gln Val Gln Gly Leu 185 Leu Gln Leu Gln Ala Gly Gln Tyr Ala Ser Val Phe Val Asp Asn Gly 200

Ser Gly Ala Val Leu Thr Ile Gln Ala Gly Ser Ser Phe Ser Gly Leu

Leu Leu Gly Thr 225

195

<210> 10

<211> 1232

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (241)..(1104)

<223>

<400> 10 gggaggccta gggagaaagt agttctcttt cggtggcagg gttgctgtcg agggcaccga gcaggagata ggtcgacaga gacgaggagt tctggctcct cctgcagaca tgcaccagcg getgetggge tegteeetgg geetegeece egegegggg etetgaatge etgeegeege 180 ccccatgaga gcaccggcct gggctcccgc ccctaagcct ctgctcgcgg agactgagcc 240 atg tgg gcc tgg ggc tgg gcc gct gca gcg ctc ctc tgg cta cag act 288 Met Trp Ala Trp Gly Trp Ala Ala Ala Leu Leu Trp Leu Gln Thr gca gga gcc ggg gcc cgg cag gag ctc aag aag tct cgg cag ctg ttt 336 Ala Gly Ala Gly Ala Arg Gln Glu Leu Lys Lys Ser Arg Gln Leu Phe 20

					ccc Pro											3	384
cca Pro	ggc Gly 50	tcc Ser	gtc Val	aag Lys	ccc Pro	ccg Pro 55	gaa Glu	gcc Ala	tct Ser	gga Gly	cct Pro 60	gag Glu	ctc Leu	tca Ser	gat Asp	4	132
gcc Ala 65	cac His	atg Met	acg Thr	tgg Trp	ttg Leu 70	aac Asn	ttt Phe	gtc Val	cga Arg	cgg Arg 75	cca Pro	gat Asp	gat Asp	gly aaa	tcc Ser 80	4	480
ccc Pro	cca Pro	gga Gly	cct Pro	cct Pro 85	ggc Gly	cct Pro	cct Pro	ggt Gly	ccc Pro 90	cct Pro	ggc	tcc Ser	cct Pro	ggt Gly 95	gtg Val		528
ggc Gly	gtt Val	acc Thr	cca Pro 100	gag Glu	gcc Ala	tta Leu	ctg Leu	cag Gln 105	gaa Glu	ttt Phe	cag Gln	gag Glu	ata Ile 110	ctg Leu	aaa Lys	ţ	576
gag Glu	gcc Ala	aca Thr 115	gaa Glu	ctt Leu	cga Arg	ttc Phe	tca Ser 120	Gly 999	cta Leu	cca Pro	gac Asp	aca Thr 125	ttg Leu	tta Leu	ccc Pro	•	624
cag Gln	gaa Glu 130	ccc Pro	agc Ser	caa Gln	cgg Arg	ctg Leu 135	gtg Val	gtt Val	gag Glu	gcc Ala	ttc Phe 140	tac Tyr	tgc Cys	cgt Arg	ttg Leu	(672
aaa Lys 145	ggc Gly	cct Pro	gtg Val	ctg Leu	gtg Val 150	gac Asp	aag Lys	aag Lys	act Thr	ctg Leu 155	gtg Val	gaa Glu	ctg Leu	caa Gln	gga Gly 160	•	720
ttc Phe	caa Gln	gct Ala	cct Pro	act Thr 165	act Thr	cag Gln	ggc	gcc Ala	ttc Phe 170	ctg Leu	cgg Arg	gga Gly	tct Ser	ggc Gly 175	ctg Leu	•	768
agc Ser	ctg Leu	tcc Ser	ttg Leu 180	ggc Gly	cga Arg	ttc Phe	aca Thr	gcc Ala 185	cca Pro	gt <i>c</i> Val	tct Ser	gcc Ala	atc Ile 190	ttc Phe	cag Gln	;	816
ttt Phe	tct Ser	gcc Ala 195	agc Ser	ctg Leu	cac His	gtg Val	gac Asp 200	cac His	agt Ser	gaa Glu	ctg Leu	cag Gln 205	ggc	aga Arg	ggc Gly	;	864
cgg Arg	ttg Leu 210	cgt Arg	acc Thr	cgg Arg	gat Asp	atg Met 215	gtc Val	cgt Arg	gtt Val	ctc Leu	atc Ile 220	Cys	att Ile	gag Glu	tcc Ser		912
ttg Leu 225	Cys	cat His	cgt Arg	cat His	acg Thr 230	tcc Ser	ctg Leu	gag Glu	gct Ala	gta Val 235	tca Ser	ggt Gly	ctg Leu	gag Glu	agc Ser 240		960
aac Asn	agc Ser	agg Arg	gtc Val	ttc Phe 245	aca Thr	gtg Val	cag Gln	gtt Val	cag Gln 250	Gly	ctg Leu	ctg Leu	cat His	cta Leu 255	Gln	1	800
tct Ser	gga Gly	cag Gln	tat Tyr 260	gtc Val	tct Ser	gtg Val	ttc Phe	gtg Val 265	Asp	aac Asn	agt Ser	tct Ser	ggg Gly 270	gca Ala	gtc Val	1	.056

ctc Leu	acc Thr	atc Ile 275	cag Gln	aac Asn	act Thr	tcc Ser	agc Ser 280	ttc Phe	tcg Ser	gga Gly	atg Met	ctt Leu 285	ttg Leu	ggt Gly	acc Thr	1104
tago	ggag	jct g	aaga	aacg	a tt	gtgg	attg	g agg	aacc	aac	acct	tgct	tc t	taga	ggagc	1164
tgaa	aagg	jac t	acto	acto	c cc	tttt	aata	gtt	ttca	tag	caat	aaag	aa c	tcca	aactt	1224
ctto	atct	:														1232
<210)> 1	11														
<211	.> 2	288														
<212	!> I	PRT														
<213	l <	lus n	uscu	lus												
<400)> :	11														
Met 1	Trp	Ala	Trp	Gly 5	Trp	Ala	Ala	Ala	Ala 10	Leu	Leu	Trp	Leu	Gln 15	Thr	
Ala	Gly	Ala	Gly 20	Ala	Arg	Gln	Glu	Leu 25	Lys	Lys	Ser	Arg	Gln 30	Leu	Phe	
Ala	Arg	Val 35	Asp	Ser	Pro	Asn	Ile 40	Thr	Thr	Ser	Asn	Arg 45	Glu	Gly •	Phe	
Pro	Gly 50	Ser	Val	Lys	Pro	Pro 55	Glu	Ala	Ser	Glý	Pro 60	Glu	Leu	Ser	Asp	
Ala 65	His	Met	Thr	Trp	Leu 70	Asn	Phe	Val	Arg	Arg 75	Pro	Asp	Asp	Gly	Ser 80	
Pro	Pro	Gly	Pro	Pro 85	Gly	Pro	Pro	Gly	Pro 90	Pro	Gly	Ser	Pro	Gly 95	Val	
Gly	Val	Thr	Pro 100	Glu	Ala	Leu	Leu	Gln 105	Glu	Phe	Gln	Glu	Ile 110	Leu	Lys	
Glu	Ala	Thr 115	Glu	Leu	Arg	Phe	Ser 120		Leu	Pro	Asp	Thr 125	Leu	Leu	Pro	
Gln	Glu 130		Ser	Gln	Arg	Leu 135		Val	Glu	Ala	Phe 140	Tyr	Cys	Arg	Leu	

18

Lys Gly Pro Val Leu Val Asp Lys Lys Thr Leu Val Glu Leu Gln Gly 145 150

Phe Gln Ala Pro Thr Thr Gln Gly Ala Phe Leu Arg Gly Ser Gly Leu

Ser Leu Ser Leu Gly Arg Phe Thr Ala Pro Val Ser Ala Ile Phe Gln 180 185

Phe Ser Ala Ser Leu His Val Asp His Ser Glu Leu Gln Gly Arg Gly 200

Arg Leu Arg Thr Arg Asp Met Val Arg Val Leu Ile Cys Ile Glu Ser

Leu Cys His Arg His Thr Ser Leu Glu Ala Val Ser Gly Leu Glu Ser 230

Asn Ser Arg Val Phe Thr Val Gln Val Gln Gly Leu Leu His Leu Gln 250 245

Ser Gly Gln Tyr Val Ser Val Phe Val Asp Asn Ser Ser Gly Ala Val 265 260

Leu Thr Ile Gln Asn Thr Ser Ser Phe Ser Gly Met Leu Leu Gly Thr 275 280

<210> 12

<211> 477

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(474)

<223>

gcg ccg cgc gtg gag gcc gct ttc ctc tgc cgc ctg cgc cgg gac gcg 48

Ala 1	Pro	Arg	Val	Glu 5	Ala	Ala	Phe	Leu	Cys 10	Arg	Leu	Arg	Arg	Asp 15	Ala		
					gcg Ala											96	
					ttc Phe											144	
ggc Gly	cag Gln 50	tac Tyr	agg Arg	gcg Ala	ccc Pro	gtg Val 55	gct Ala	ggc Gly	ttc Phe	tac Tyr	gct Ala 60	ctc Leu	gcc Ala	gcc Ala	acg Thr	192	
ctg Leu 65	cac His	gtg Val	gcg Ala	ctc Leu	999 Gly 70	gag Glu	ccg Pro	ccg Pro	agg Arg	agg Arg 75	ggg Gly	ccg Pro	ccg Pro	cgc Arg	ccc Pro 80	240	
					ctg Leu											288	
					gcc Ala											336	
					aat Asn											384	
acc Thr	tcc Ser 130	tgg Trp	gcg Ala	tgt Cys	gag Glu	cgg Arg 135	cca Pro	cca Pro	cag Gln	gcc Ala	ctt Leu 140	cct Pro	ctc Leu	agg Arg	ggc Gly	432	
aaa Lys 145	tgg Trp	agc Ser	aca Thr	gat Asp	cta Leu 150	gac Asp	aat Asn	gtg Val	tgg Trp	aca Thr 155	gtg Val	tca Ser	gag Glu	tag		477	
<210)> :	13															
<21	l> :	158															
<212	2> :	PRT															
<21	3> 1	Homo	sap	iens													
	•				•		•						-	•			
<400)>	13															
Ala	Pro	Arg	Val	Glu 5	Ala	Ala	Phe	Leu	Cys	Arg	Leu	Arg	Arg	Asp 15	Ala		

Leu Val Glu Arg Arg Ala Leu His Glu Leu Gly Val Tyr Tyr Leu Pro 20 25 30

Asp Ala Glu Gly Ala Phe Arg Arg Gly Pro Gly Leu Asn Leu Thr Ser 35 40 45

Gly Gln Tyr Arg Ala Pro Val Ala Gly Phe Tyr Ala Leu Ala Ala Thr 50 55 60

Leu His Val Ala Leu Gly Glu Pro Pro Arg Arg Gly Pro Pro Arg Pro 65 70 75 80

Arg Asp His Leu Arg Leu Leu Ile Cys Ile Gln Ser Arg Cys Gln Arg 85 90 95

Asn Thr Ser Leu Glu Ala Ile Met Gly Leu Glu Ser Ser Glu Leu 100 105 110

Phe Thr Ile Ser Val Asn Gly Val Leu Tyr Leu Gln Met Gly Gln Trp 115 120 125

Thr Ser Trp Ala Cys Glu Arg Pro Pro Gln Ala Leu Pro Leu Arg Gly
130 135 140

Lys Trp Ser Thr Asp Leu Asp Asn Val Trp Thr Val Ser Glu 145 150 155

<210> 14

<211> 3180

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (143)..(2677)

<223>

<400> 14
gctggaagca gcgtcttatt ttaccttgtt ctcccacttc ctgaagatgc taaactcctg 60
gtggactgca gaggagaggg attcagtctt ctcctgatgt gtttgcctgt aggtacctga 120
gttgacaccg aagctcctaa ag atg ctg age ggc gtt tgg ttc ctc agt gtg 172

						Met 1	. Leu	ı Ser	: Gly	Val	Trp	Phe	. Leu	Ser	Val	
tta Leu	acc Thr	gtg Val	gcc Ala	999 Gly 15	atc Ile	tta Leu	cag Gln	aca Thr	gag Glu 20	agt Ser	cgc Arg	aaa Lys	act Thr	gcc Ala 25	aaa Lys	220
gac Asp	att Ile	tgc Cys	aag Lys 30	atc Ile	cgc Arg	tgt Cys	ctg Leu	tgc Cys 35	gaa Glu	gaa Glu	aag Lys	gaa Glu	aac Asn 40	gta Val	ctg Leu	268
aat Asn	atc Ile	aac Asn 45	tgt Cys	gag Glu	aac Asn	aaa Lys	gga Gly 50	ttt Phe	aca Thr	aca Thr	gtt Val	agc Ser 55	ctg Leu	ctc Leu	cag Gln	316
ccc Pro	ccc Pro 60	cag Gln	tat Tyr	cga Arg	atc Ile	tat Tyr 65	cag Gln	ctt Leu	ttt Phe	ctc Leu	aat Asn 70	gga Gly	aac Asn	ctc Leu	ttg Leu	364
aca Thr 75	aga Arg	ctg Leu	tat Tyr	cca Pro	aac Asn 80	gaa Glu	ttt Phe	gtc Val	aat Asn	tac Tyr 85	tcc Ser	aac Asn	gcg Ala	gtg Val	act Thr 90	412
ctt Leu	cac His	cta Leu	ggt Gly	aac Asn 95	aac Asn	G1A 888	tta Leu	cag Gln	gag Glu 100	atc Ile	cga Arg	acg Thr	61 ^y 333	gca Ala 105	ttc Phe	460
agt Ser	ggc Gly	ctg Leu	aaa Lys 110	act Thr	ctc Leu	aaa Lys	aga Arg	ctg Leu 115	cat His	ctc Leu	aac Asn	aac Asn	aac Asn 120	aag Lys	ctt Leu	508
gag Glu	ata Ile	ttg Leu 125	agg Arg	gag Glu	gac Asp	acc Thr	ttc Phe 130	cta Leu	ggc Gly	ctg Leu	gag Glu	agc Ser 135	ctg Leu	gag Glu	tat Tyr	556
ctc Leu	cag Gln 140	gcc Ala	gac Asp	tac Tyr	aat Asn	tac Tyr 145	atc Ile	agt Ser	gcc Ala	atc Ile	gag Glu 150	gct Ala	Gly 999	gca Ala	ttc Phe	604
agc Ser 155	aaa Lys	ctt Leu	aac Asn	aag Lys	ctc Leu 160	aaa Lys	gtg Val	ctc Leu	atc Ile	ctg Leu 165	aat Asn	gac Asp	aac Asn	ctt Leu	ctg Leu 170	652
					aat Asn											700
gac Asp	ctc Leu	agg Arg	999 Gly 190	aat Asn	agg Arg	cta Leu	aaa Lys	gta Val 195	atg Met	cct Pro	ttt Phe	gct Ala	ggc Gly 200	gtc Val	ctt Leu	748
gaa Glu	cat His	att Ile 205	gga Gly	gjy aaa	atc Ile	atg Met	gag Glu 210	Ile	cag Gln	ctg Leu	gag Glu	gaa Glu 215	Asn	cca Pro	tgg Trp	796
aat Asn	tgc Cys 220	act Thr	tgt Cys	gac Asp	tta Leu	ctt Leu 225	cct Pro	ctc Leu	aag Lys	gcc Ala	tgg Trp 230	cta Leu	gac Asp	acc Thr	ata Ile	844

. 22

			gtg Val													892
gly ggg	aaa Lys	gac Asp	gtg Val	acc Thr 255	cag Gln	ctg Leu	acc Thr	agg Arg	caa Gln 260	gac Asp	ctc Leu	tgt Cys	ccc Pro	aga Arg 265	aaa Lys	940
			gat Asp 270													988
caa Gln	agg Arg	ctg Leu 285	tca Ser	cct Pro	aca Thr	atg Met	aat Asn 290	cct Pro	gct Ala	ctc Leu	aac Asn	cca Pro 295	acc Thr	agg Arg	gct Ala	1036
ccg Pro	aaa Lys 300	gcc Ala	agc Ser	cgg Arg	ccg Pro	ccc Pro 305	aaa Lys	atg Met	aga Arg	a'at Asn	cgt Arg 310	cca Pro	act Thr	ccc Pro	cga Arg	1084
gtg Val 315	act Thr	gtg Val	tca Ser	aag Lys	gac Asp 320	agg Arg	caa Gln	agt Ser	ttt Phe	gga Gly 325	ccc Pro	atc Ile	atg Met	gtg Val	tac Tyr 330	1132
cag Gln	acc Thr	aag Lys	tct Ser	cct Pro 335	gtg Val	cct Pro	ctc Leu	acc Thr	tgt Cys 340	ccc Pro	agc Ser	agc Ser	tgt Cys	gtc Val 345	tgc Cys	1180
acc Thr	tct Ser	cag Gln	agc Ser 350	tca Ser	gac Asp	aat Asn	ggt Gly	ctg Leu 355	aat Asn	gta Val	aac Asn	tgc Cys	caa Gln 360	gaa Glu	agg Arg	1228
aag Lys	ttc Phe	act Thr 365	aat Asn	atc Ile	tct Ser	gac Asp	ctg Leu 370	cag Gln	ccc Pro	aaa Lys	ccg Pro	acc Thr 375	agt Ser	cca Pro	aag Lys	1276
aaa Lys	ctc Leu 380	tac Tyr	cta Leu	aca Thr	Gly aaa	aac Asn 385	tat Tyr	ctt Leu	caa Gln	act Thr	gtc Val 390	tat Tyr	aag Lys	aat Asn	gac Asp	1324
ctc Leu 395	Leu	gaa Glu	tac Tyr	agt Ser	tct Ser 400	Leu	gac Asp	tta Leu	ctg Leu	cac His 405	Leu	gga Gly	aac Asn	aac Asn	agg Arg 410	1372
att Ile	gca Ala	gtc Val	att Ile	cag Gln 415	gaa Glu	ggt Gly	gcc Ala	ttt Phe	aca Thr 420	aac Asn	ctg Leu	acc Thr	agt Ser	tta Leu 425	cgc Arg	1420
aga Arg	ctt Leu	tat Tyr	ctg Leu 430	aat Asn	ggc	aat Asn	tac Tyr	ctt Leu 435	gaa Glu	gtg Val	ctg Leu	tac Tyr	cct Pro 440	tct Ser	atg Met	1468
ttt Phe	gat Asp	gga Gly 445	ctg Leu	cag Gln	agc Ser	ttg Leu	caa Gln 450	Tyr	ctc Leu	tat Tyr	tta Leu	gag Glu 455	Tyr	aat Asn	gtc Val	1516
att Ile	aag Lys 460	Glu	att Ile	aag Lys	cct Pro	ctg Leu 465	Thr	ttt Phe	gat Asp	gct Ala	ttg Leu 470	Ile	aac Asn	cta Leu	cag Gln	1564

														•			
cta Leu 475	ctg Leu	ttt Phe	ctg Leu	aac Asn	aac Asn 480	aac Asn	ctt Leu	ctt Leu	cgg Arg	tcc Ser 485	tta Leu	cct Pro	gat Asp	aat Asn	ata Ile 490	1	.612
ttt Phe	Gly 393	gly ggg	acg Thr	gcc Ala 495	cta Leu	acc Thr	agg Arg	ctg Leu	aat Asn 500	ctg Leu	aga Arg _.	aac Asn	aac Asn	cat His 505	ttt Phe	1	.660
tċt Ser	cac His	ctg Leu	ccc Pro 510	gtg Val	aaa Lys	gly ggg	gtt Val	ctg Leu 515	gat Asp	cag Gln	ctc Leu	ccg Pro	gct Ala 520	ttc Phe	atc Ile	1	.708
cag Gln	ata Ile	gat Asp 525	ctg Leu	cag Gln	gag Glu	aac Asn	ccc Pro 530	tgg Trp	gac Asp	tgt Cys	acc Thr	tgt Cys 535	gac Asp	atc Ile	atg Met	1	1756
gly ggg	ctg Leu 540	aaa Lys	gac [°] Asp	tgg Trp	aca Thr	gaa Glu 545	cat His	gcc Ala	aat Asn	tcc Ser	cct Pro 550	gtc Val	atc Ile	att Ile	aat Asn	1	L804
gag Glu 555	gtg Val	act Thr	tgc Cys	gaa Glu	tct Ser 560	cct Pro	gct Ala	aag Lys	cat His	gca Ala 565	Gly 999	gag Glu	ata Ile	cta Leu	aaa Lys 570	1	L852
ttt Phe	ctg Leu	gly aaa	agg Arg	gag Glu 575	gct Ala	atc Ile	tgt Cys	cca Pro	gac Asp 580	agc Ser	cca Pro	aac Asn	ttg Leu	tca Ser 585	gat Asp	:	1900
gga Gly	acc Thr	gtc Val	ttg Leu 590	tca Ser	atg Met	aat Asn	cac His	aat Asn 595	aca Thr	gac Asp	aca Thr	cct Pro	cgg Arg 600	tcg Ser	ctt Leu	:	1948
agt Ser	gtg Val	tct Ser 605	cct Pro	agt Ser	tcc Ser	tat Tyr	cct Pro 610	gaa Glu	cta Leu	cac His	act Thr	gaa Glu 615	gtt Val	cca Pro	ctg Leu	:	1996
tct Ser	gtc Val 620	tta Leu	att Ile	ctg Leu	gga Gly	ttg Leu 625	ctt Leu	gtt Val	gtt Val	ttc Phe	atc Ile 630	tta Leu	tct Ser	gtc Val	tgt Cys	;	2044
ttt Phe 635	gly ggg	gct Ala	ggt Gly	tta Leu	ttc Phe 640	gtc Val	ttt Phe	gtc Val	ttg Leu	aaa Lys 645	cgc Arg	cga Arg	aag Lys	gga Gly	gtg Val 650	:	2092
ccg Pro	agc Ser	gtt Val	ccc Pro	agg Arg 655	aat Asn	acc Thr	aac Asn	aac Asn	tta Leu 660	Asp	gta Val	agc Ser	tcc Ser	ttt Phe 665	Gln	;	2140
tta Leu	cag Gln	tat Tyr	999 Gly 670	Ser	tac Tyr	aac Asn	act Thr	gag Glu 675	act Thr	cac His	gat Asp	aaa Lys	aca Thr 680	Asp	ggc Gly		2188
cat His	gtc Val	tac Tyr 685	Asn	tat Tyr	atc Ile	ccc Pro	cca Pro 690	Pro	gtg Val	ggt Gly	cag Gln	atg Met 695	Cys	caa Gln	aac Asn		2236
ccc Pro	atc	tac Tyr	atg Met	cag Gln	aag Lys	gaa	gga Gly	gac Asp	cca Pro	gta Val	gcc Ala	tat Tyr	tac Tyr	cga Arg	aac Asn		2284

24

700	705	710	
ctg caa gag ttc agc tat Leu Gln Glu Phe Ser Tyr 715 720	Ser Asn Leu		
gcc aca cct gct tac aca Ala Thr Pro Ala Tyr Thr 735	: Ile Ser Ala	act gag ctg cta Thr Glu Leu Leu 740	gaa aag cag 2380 Glu Lys Gln 745
gcc aca cca aga gag cct Ala Thr Pro Arg Glu Pro 750	gag ctg ctg Glu Leu Leu 755	tat caa aat att Tyr Gln Asn Ile	gct gag cga 2428 Ala Glu Arg 760
gtc aag gaa ctt ccc ago Val Lys Glu Leu Pro Ser 765	gca ggc cta Ala Gly Leu 770	gtc cac tat aac Val His Tyr Asn 775	ttt tgt acc 2476 Phe Cys Thr
tta cct aaa agg cag tti Leu Pro Lys Arg Gln Pho 780	gcc cct tcc Ala Pro Ser 785	tat gaa tct cga Tyr Glu Ser Arg 790	cgc caa aac 2524 Arg Gln Asn
caa gac aga atc aat aa Gln Asp Arg Ile Asn Ly: 795 800	Thr Val Leu		
ttt gtg ggg cag tca aaa Phe Val Gly Gln Ser Lys 815	a ccc aac cac s Pro Asn His	cct tta ctg caa Pro Leu Leu Gln 820	gct aag ccg 2620 Ala Lys Pro 825
caa tca gaa ccg gac ta Gln Ser Glu Pro Asp Ty: 830	c ctc gaa gtt c Leu Glu Val 835	ctg gaa aaa caa Leu Glu Lys Gln	act gca atc 2668 Thr Ala Ile 840
agt cag ctg tgaagggaaa Ser Gln Leu 845	tcatttacaa co	ctaaggca tcagagg	gatg 2717
ctgctccgaa ctgttggaaa	caaggacatt ago	ttttgtg tttgttt	ttg ttetecettt 2777
cccagtgtta atgggggact	ttgaaaatgt ttg	ggagata ggatgaag	gtc atgattttgc 2837
ttttgcaagt tttcctttaa	attatttctc tct	egetete eteceet	cct tttttttttt 2897
tttttttt tcttttccc	ttctcttctt agg	gaaccatc agtggac	atg aatgtttcta 2957
caatgcattt cttcatagat	tttgtttatg gtt	ttgtttc tttttc	ttc tttgtttttc 3017
agtgtgggag tgggaagagg	agattatagt gad	ctgaagaa agaatag	gca aacttttcaa 3077
atgaaaatgg atatttagtg	tattttgtag aag	gateteca aagatet	ttt gtgactacaa 3137
cttcttttgt aaataatgat	atatggtatt tco	categica git	3180

<210> 15 .

<211> 845

PCT/US01/28013

<212> PRT

WO 02/20569

<213> Homo sapiens

. <400> 15

Met Leu Ser Gly Val Trp Phe Leu Ser Val Leu Thr Val Ala Gly Ile 1 5 10 15

Leu Gln Thr Glu Ser Arg Lys Thr Ala Lys Asp Ile Cys Lys Ile Arg 20 25 30

Cys Leu Cys Glu Glu Lys Glu Asn Val Leu Asn Ile Asn Cys Glu Asn 35 40 45

Lys Gly Phe Thr Thr Val Ser Leu Leu Gln Pro Pro Gln Tyr Arg Ile 50 55 60

Tyr Gln Leu Phe Leu Asn Gly Asn Leu Leu Thr Arg Leu Tyr Pro Asn 65 70 75 80

Glu Phe Val Asn Tyr Ser Asn Ala Val Thr Leu His Leu Gly Asn Asn 90 95

Gly Leu Gln Glu Ile Arg Thr Gly Ala Phe Ser Gly Leu Lys Thr Leu 100 105 110

Lys Arg Leu His Leu Asn Asn Asn Lys Leu Glu Ile Leu Arg Glu Asp 115 120 125

Thr Phe Leu Gly Leu Glu Ser Leu Glu Tyr Leu Gln Ala Asp Tyr Asn 130 135 140

Tyr Ile Ser Ala Ile Glu Ala Gly Ala Phe Ser Lys Leu Asn Lys Leu 145 150 155 160

Lys Val Leu Ile Leu Asn Asp Asn Leu Leu Ser Leu Pro Ser Asn 165 170 175

Val Phe Arg Phe Val Leu Leu Thr His Leu Asp Leu Arg Gly Asn Arg 180 185 190

Leu Lys Val Met Pro Phe Ala Gly Val Leu Glu His Ile Gly Gly Ile 195 200 205 Met Glu Ile Gln Leu Glu Glu Asn Pro Trp Asn Cys Thr Cys Asp Leu 215 Leu Pro Leu Lys Ala Trp Leu Asp Thr Ile Thr Val Phe Val Gly Glu 230 235 Ile Val Cys Glu Thr Pro Phe Arg Leu His Gly Lys Asp Val Thr Gln 250 Leu Thr Arg Gln Asp Leu Cys Pro Arg Lys Ser Ala Ser Asp Ser Ser 265 260 Gln Arg Gly Ser His Ala Asp Thr His Val Gln Arg Leu Ser Pro Thr Met Asn Pro Ala Leu Asn Pro Thr Arg Ala Pro Lys Ala Ser Arg Pro . 295 300 Pro Lys Met Arg Asn Arg Pro Thr Pro Arg Val Thr Val Ser Lys Asp 305 310 315 Arg Gln Ser Phe Gly Pro Ile Met Val Tyr Gln Thr Lys Ser Pro Val 330 335 325 Pro Leu Thr Cys Pro Ser Ser Cys Val Cys Thr Ser Gln Ser Ser Asp 340 345 Asn Gly Leu Asn Val Asn Cys Gln Glu Arg Lys Phe Thr Asn Ile Ser 360 365 355 Asp Leu Gln Pro Lys Pro Thr Ser Pro Lys Lys Leu Tyr Leu Thr Gly 370 375 380 Asn Tyr Leu Gln Thr Val Tyr Lys Asn Asp Leu Leu Glu Tyr Ser Ser 395 400 390 385 Leu Asp Leu Leu His Leu Gly Asn Asn Arg Ile Ala Val Ile Gln Glu 410 405 -Gly Ala Phe Thr Asn Leu Thr Ser Leu Arg Arg Leu Tyr Leu Asn Gly 425 Asn Tyr Leu Glu Val Leu Tyr Pro Ser Met Phe Asp Gly Leu Gln Ser · 445 435 440

WO 02/20569

Leu Gln Tyr Leu Tyr Leu Glu Tyr Asn Val Ile Lys Glu Ile Lys Pro 450 455 460

. Leu Thr Phe Asp Ala Leu Ile Asn Leu Gln Leu Leu Phe Leu Asn Asn 465 470 475 480

Asn Leu Leu Arg Ser Leu Pro Asp Asn Ile Phe Gly Gly Thr Ala Leu 485 490 495

Thr Arg Leu Asn Leu Arg Asn Asn His Phe Ser His Leu Pro Val Lys 500 505 510

Gly Val Leu Asp Gln Leu Pro Ala Phe Ile Gln Ile Asp Leu Gln Glu 515 520 525

Aşn Pro Trp Asp Cys Thr Cys Asp Ile Met Gly Leu Lys Asp Trp Thr 530 540

Glu His Ala Asn Ser Pro Val Ile Ile Asn Glu Val Thr Cys Glu Ser 545 550 555 560

Pro Ala Lys His Ala Gly Glu Ile Leu Lys Phe Leu Gly Arg Glu Ala 565 570 575

Ile Cys Pro Asp Ser Pro Asn Leu Ser Asp Gly Thr Val Leu Ser Met 580 585 590

Asn His Asn Thr Asp Thr Pro Arg Ser Leu Ser Val Ser Pro Ser Ser 595 600 605

Tyr Pro Glu Leu His Thr Glu Val Pro Leu Ser Val Leu Ile Leu Gly 610 620

Leu Leu Val Val Phe Ile Leu Ser Val Cys Phe Gly Ala Gly Leu Phe 625 630 635 640

Val Phe Val Leu Lys Arg Arg Lys Gly Val Pro Ser Val Pro Arg Asn 645 650 655

Thr Asn Asn Leu Asp Val Ser Ser Phe Gln Leu Gln Tyr Gly Ser Tyr 660 665 670

Asn Thr Glu Thr His Asp Lys Thr Asp Gly His Val Tyr Asn Tyr Ile

								_							
		675					680					685			
Pro	Pro 690	Pro	Val	Gly	Gln	Met 695	Cys	Gln	Asn	Pro	Ile 700	Tyr	Met	Gln	Lys
Glu 705	Gly	Asp	Pro	Val	Ala 710	Tyr	Tyr	Arg	Asn	Leu 715	Gln [°]	Glu	Phe	Ser	Tyr 720
Ser	Asn	Leu	Glu	Glu 725	Lys	Lys	Glu	Glu	Pro 730	Ala	Thr	Pro	Ala	Tyr 735	Thr
Ile	Ser	Ala	Thr 740	Glu	Leu	Leu	Glu	Lys 745	Gln	Ala	Thr	Pro	Arg 750	Glu	Pro
Glu	Leu	Leu 755	Tyr	Gln	Asn	Ile	Ala 760	Glu	Arg	Val	Lys	Glu 765	Leu	Pro	Ser
Ala	Gly 770		Val	His	Tyr	Asn 775	Phe	Сув	Thr	Leu	Pro 780	Lys	Arg	Gln	Phe
Ala 785	Pro	Ser	Tyr	Glu	Ser 790	Arg	Arg	Gln	Asn	Gln 795	Asp	Arg	Ile	Asn	Lys 800
Thr	Val	Leu	Tyr	Gly 805	Thr	Pro	Arg	ГÀЗ	Cys 810	Phe	Val	Gly	Gln	Ser 815	Lys
Pro	Asn	His	Pro 820	Leu	Leu	Gln	Ala	Lys 825	Pro	Gln	Ser	Glu	Pro 830	Asp	Tyr
Leu	Glu	Val 835	Leu	Glu	Lys	Gln	Thr 840	Ala	Ile	Ser	Gln	Leu 845			
<21	0>	16													
<21	1>	469													
<21	2>	DNA													
<21	3>	Mus	musc	ulus											
-40	٥.	16													

<400> 16
ctgaaattcc tgggaaggga ggctatttgt ccagaaaatc ctaacctgtc agatgggact 60
attttgtcaa tgaatcacaa cacagacaca cctagatcac ttagtgtgtc tcctagttct 120
taccccgaac tacacactga agttccactc tccgttttaa ttttaggatt gcttgtggtt 180

.

WO 02/20569 PCT/US01/28013

29

tttatcctgt	ctgtctgttt	tggggcgggg	ttgttcgtct	ttgttctgaa	gcgtcgaaag	240
ggagtgccaa	atgttcccag	gaatgccacc	aacttagatg	taagttcctt	ccagttacaa	300
tatgggtctt	acaacaccga	gactaatgat	aaagctgatg	gccacgtcta	taactacatt	360
cctccacctg	tgggtcagat	gtgccaaaac	cccatctaca	tgcagaagga	aggagaccca	420
gtggcctatt	accgaaatct	gcaggacttc	agctatggca	acctggagg		469

<210> 17

<211> 156

<212> PRT

<213> Mus musculus

<400> 17

Leu Lys Phe Leu Gly Arg Glu Ala Ile Cys Pro Glu Asn Pro Asn Leu 1 5 10 15

Ser Asp Gly Thr Ile Leu Ser Met Asn His Asn Thr Asp Thr Pro Arg 20 25 30

Ser Leu Ser Val Ser Pro Ser Ser Tyr Pro Glu Leu His Thr Glu Val 35 40 45

Pro Leu Ser Val Leu Ile Leu Gly Leu Leu Val Val Phe Ile Leu Ser 50 55 60

Val Cys Phe Gly Ala Gly Leu Phe Val Phe Val Leu Lys Arg Arg Lys 65 70 75 80

Gly Val Pro Asn Val Pro Arg Asn Ala Thr Asn Leu Asp Val Ser Ser 85 90 95

Phe Gln Leu Gln Tyr Gly Ser Tyr Asn Thr Glu Thr Asn Asp Lys Ala 100 105 110

Asp Gly His Val Tyr Asn Tyr Ile Pro Pro Pro Val Gly Gln Met Cys 115 120 125

Gln Asn Pro Ile Tyr Met Gln Lys Glu Gly Asp Pro Val Ala Tyr Tyr 130 135 140

145 150 155	
<210> 18	
<211> 3402	
<212> DNA	
<213> Homo sapiens	
<220>	
<221> CDS	
<222> (89)(2899)	
<223>	
<400> 18 tagacgcgga gcccaaggag gtaaaatgca cacttgctgc cccccagtaa ctttggaaca	60
ggaccttcac agaaaaatgc atagctgg atg ctg cag act cta gcg ttt gct Met Leu Gln Thr Leu Ala Phe Ala	112
1 5	
gta aca tct ctc gtc ctt tcg tgt gca gaa acc atc gat tat tac ggg Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly	160
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20	-
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu	160 208
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 30 35 40	
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu	208
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 30 35 40 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 50 55 cct ccc cgt ttc cca atc tac cac ctc ttg ttg tcc gga aac ctt ttg	208
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 30 35 40 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 50 55	208
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 cct ccc cgt ttc cca atc tac cac ctc ttg ttg tcc gga aac ctt ttg Pro Pro Arg Phe Pro Ile Tyr His Leu Leu Leu Ser Gly Asn Leu Leu 60 aac cgt ctc tat ccc aat gag ttt gtc aat tac act ggg gct tca att	208
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 cct ccc cgt ttc cca atc tac cac ctc ttg ttg tcc gga aac ctt ttg Pro Pro Arg Phe Pro Ile Tyr His Leu Leu Leu Ser Gly Asn Leu Leu 60 65 70	208 256 304
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 30 35 40 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 50 55 cct ccc cgt ttc cca atc tac cac ctc ttg ttg tcc gga aac ctt ttg Pro Pro Arg Phe Pro Ile Tyr His Leu Leu Leu Ser Gly Asn Leu Leu 60 65 70 aac cgt ctc tat ccc aat gag ttt gtc aat tac act ggg gct tca att Asn Arg Leu Tyr Pro Asn Glu Phe Val Asn Tyr Thr Gly Ala Ser Ile 75 80 ttg cat cta qgt agc aat gtt atc cag gac att gag acc ggg gct ttc	208 256 304
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 cct ccc cgt ttc cca atc tac cac ctc ttg ttg tcc gga aac ctt ttg Pro Pro Arg Phe Pro Ile Tyr His Leu Leu Leu Ser Gly Asn Leu Leu 60 aac cgt ctc tat ccc aat gag ttt gtc aat tac act ggg gct tca att Asn Arg Leu Tyr Pro Asn Glu Phe Val Asn Tyr Thr Gly Ala Ser Ile 80	208 256 304
Val Thr Ser Leu Val Leu Ser Cys Ala Glu Thr Ile Asp Tyr Tyr Gly 10 15 20 gaa atc tgt gac aat gca tgt cct tgt gag gaa aag gac ggc att tta Glu Ile Cys Asp Asn Ala Cys Pro Cys Glu Glu Lys Asp Gly Ile Leu 25 30 35 40 act gtg agc tgt gaa aac cgg ggg atc atc agt ctc tct gaa att agc Thr Val Ser Cys Glu Asn Arg Gly Ile Ile Ser Leu Ser Glu Ile Ser 45 50 55 cct ccc cgt ttc cca atc tac cac ctc ttg ttg tcc gga aac ctt ttg Pro Pro Arg Phe Pro Ile Tyr His Leu Leu Ser Gly Asn Leu Leu 60 65 70 aac cgt ctc tat ccc aat gag ttt gtc aat tac act ggg gct tca att Asn Arg Leu Tyr Pro Asn Glu Phe Val Asn Tyr Thr Gly Ala Ser Ile 75 80 85 ttg cat cta ggt agc aat gtt atc cag gac att gag acc ggg gct ttc Leu His Leu Gly Ser Asn Val Ile Gln Asp Ile Glu Thr Gly Ala Phe	208 256 304

gaa Glu	ctt Leu	ctg Leu	cga Arg	gat Asp 125	gat Asp	acc Thr	ttc Phe	ctt Leu	ggc Gly 130	ttg Leu	gag Glu	aac Asn	ctg Leu	gag Glu 135	tac Tyr	496
cta Leu	cag Gln	gtc Val	gat Asp 140	tac Tyr	aac Asn	tac Tyr	atc Il <u>e</u>	agc Ser 145	gtc Val	att Ile	gaa Glu	ccc Pro	aat Asn 150	gct Ala	ttt Phe	544
999 Gly	aaa Lys	ctg Leu 155	cat His	ttg Leu	ttg Leu	cag Gln	gtg Val 160	ctt Leu	atc Ile	ctc Leu	aat Asn	gac Asp 165	aat Asn	ctt Leu	ttg Leu	592
tcc Ser	agt Ser 170	tta Leu	ccc Pro	aac Asn	aat Asn	ctt Leu 175	ttc Phe	cgt Arg	ttt Phe	gtg Val	ccc Pro 180	tta Leu	acg Thr	cac His	ttg Leu	640
gac Asp 185	ctc Leu	cgg Arg	Gly 999	aac Asn	cgg Arg 190	ctg Leu	aaa Lys	ctt Leu	ctg Leu	ccc Pro 195	tac Tyr	gtg Val	gly ggg	ctc Leu	ttg Leu 200	688
cag Gln	cac His	atg Met	gat Asp	aaa Lys 205	gtt Val	gtg Val	gag Glu	cta Leu	cag Gln 210	ctg Leu	gag Glu	gaa Glu	aac Asn	cct Pro 215	tgg Trp	736
aat Asn	tgt Cys	tct Ser	tgt Cys 220	gag Glu	ctg Leu	atc Ile	tct Ser	cta Leu 225	aag Lys	gat Asp	tgg Trp	ttg Leu	gac Asp 230	agc Ser	atc Ile	784
tcc Ser	tat Tyr	tca Ser 235	gcc Ala	ctg Leu	gtg Val	ggg Gly	gat Asp 240	gta Val	gtt Val	tgt Cys	gag Glu	acc Thr 245	ccc Pro	ttc Phe	cgc Arg	832
tta Leu	cac His 250	gga Gly	agg Arg	gac Asp	ttg Leu	gac Asp 255	gag Glu	gta Val	tcc Ser	aag Lys	cag Gln 260	gaa Glu	ctt Leu	tgc Cys	cca Pro	880
agg Arg 265	aga Arg	ctt Leu	att Ile	tct Ser	gac Asp 270	tac Tyr	gag Glu	atg Met	agg Arg	ccg Pro 275	cag Gln	acg Thr	cct Pro	ttg Leu	agc Ser 280	928
acc Thr	acg Thr	ggg	Tyr	tta Leu 285	His	acc Thr	acc Thr	ccg Pro	gcg Ala 290	Ser	gtg Val	aat Asn	tct Ser	gtg Val 295	gcc Ala	976
act Thr	tct Ser	tcc Ser	tct Ser 300	gct Ala	gtt Val	tac Tyr	aaa Lys	ccc Pro 305	Pro	ttg Leu	aag Lys	ccc Pro	cct Pro 310	aag Lys	gjå aaa	1024
act Thr	cgc Arg	caa Gln 315	Pro	aac Asn	aag Lys	ccc Pro	agg Arg 320	gtg Val	cgc	ccc Pro	acc Thr	tct Ser 325	Arg	cag Gln	ccc Pro	1072
tct Ser	aag Lys 330	gac Asp	ttg Leu	ggc Gly	tac Tyr	agc Ser 335	Asn	tat Tyr	ggc	ccc	agc Ser 340	Ile	gcc	tat Tyr	cag Gln	1120
acc Thr	aaa Lys	tcc Ser	ccg Pro	gtg Val	cct Pro	ttg Leu	gag Glu	tgt Cys	ccc	acc Thr	gcg	tgo Cys	tct	tgc Cys	aac Asn	1168

345					350					355					360	
ctg Leu	cag Gln	atc Ile	tct Ser	gat Asp 365	ctg Leu	ggc Gly	ctc Leu	aac Asn	gta Val 370	aac Asn	tgc Cys	cag Gln	gag Glu	cga Arg 375	aag Lys	1216
atc Ile	gag Glu	agc Ser	atc Ile 380	gct Ala	gaa Glu	ctg Leu	cag Gln	ccc Pro 385	aag Lys	ccc Pro	tac Tyr	aat Asn	ccc Pro 390	aag Lys	aaa Lys	1264
atg Met	tat Tyr	ctg Leu 395	aca Thr	gag Glu	aac Asn	tac Tyr	atc Ile 400	gct Ala	gtc Val	gtg Val	cgc Arg	agg Arg 405	aca Thr	gac Asp	ttc Phe	1312
ctg Leu	gag Glu 410	gcc Ala	acg Thr	gjå aaa	ctg Leu	gac Asp 415	ctc Leu	ctg Leu	cac His	ctg Leu	ggg Gly 420	aat Asn	aac Asn	cgc Arg	atc Ile	1360
tcg Ser 425	atg Met	atc Ile	cag Gln	gac Asp	cgc Arg 430	gct Ala	ttc Phe	gly ggg	gat Asp	ctc Leu 435	acc Thr	aac Asn	ctg Leu	agg Arg	cgc Arg 440	1408
ctc Leu	tac Tyr	ctg Leu	aat Asn	ggc Gly 445	aac Asn	agg Arg	atc Ile	gag Glu	agg Arg 450	ctg Leu	agc Ser	ccg Pro	gag Glu	tta Leu 455	ttc Phe	1456
tat Tyr	ggc Gly	ctg Leu	cag Gln 460	agc Ser	ctg Leu	cag Gln	tat Tyr	ctc Leu 465	ttc Phe	ctc Leu	cag Gln	tac Tyr	aat Asn 470	ctc Leu	atc Ile	1504
cgc Arg	gag Glu	att Ile 475	cag Gln	tct Ser	gga Gly	act Thr	ttt Phe 480	gac Asp	ccg Pro	gtc Val	cca Pro	aac Asn 485	ctc Leu	cag Gln	ctg Leu	1552
cta Leu	ttc Phe 490	ttg Leu	aat Asn	aac Asn	aac Asn	ctc Leu 495	ctg Leu	cag Gln	gcc Ala	atg Met	ccc Pro 500	tca Ser	ggc Gly	gtc Val	ttc Phe	1600
tct Ser 505	Gly	ttg Leu	acc Thr	ctc Leu	ctc Leu 510	Arg	cta Leu	aac Asn	ctg Leu	agg Arg 515	Ser	aac Asn	cac His	ttc Phe	acc Thr 520	1648
tcc Ser	ttg Leu	cca Pro	gtg Val	agt Ser 525		gtt Val	ttg Leu	gac Asp	cag Gln 530	Leu	aag Lys	tca Ser	ctc Leu	atc Ile 535	GIn	1696
atc Ile	gac Asp	ctg Leu	cat His 540	Asp	aat Asn	cct Pro	tgg Trp	gat Asp 545	Cys	acc Thr	tgt Cys	gac Asp	att Ile 550	Val	ggc	1744
atg Met	aag Lys	ctg Leu 555	Trp	gtg Val	gag Glu	cag Gln	ctc Leu 560	ГЛЗ	gtg Val	ggc	gtc Val	cta Leu 565	Val	gac Asp	gag Glu	1792
gtg Val	atc Ile 570	Cys	aag Lys	gcg Ala	ccc Pro	aaa Lys 575	Lys	ttc Phe	gct Ala	gag Glu	acc Thr 580	Asp	atg Met	cgc Arg	tcc Ser	1840
att	aag	tcg	gag	cte	ctg	tgo	cct	gac	: tat	tca	gat	gta	gta	gtt	tee	1888

Ile 585	Lys	Ser	Glu	Leu	Leu 590	Сув	Pro	Asp	Tyr	Ser 595	Asp	Val	Val	Val	Ser 600	
acg Thr	ccc Pro	aca Thr	ccc Pro	tcc Ser 605	tct Ser	atc Ile	cag Gln	gtc Val	cct Pro 610	gcg Ala	agg Arg	acc Thr	agc Ser	gcc Ala 615	gtg Val	1936
act Thr	cct Pro	gcg Ala	gtc Val 620	cgg Arg	ttg Leu	aat Asn	agc Ser	acc Thr 625	gly ggg	gcc Ala	ccc Pro	gcg Ala	agc Ser 630	ttg Leu	ggc	1984
gca Ala	ggc Gly	gga Gly 635	gly ggg	gcg Ala	tcg Ser	tcg Ser	gtg Val 640	ccc Pro	ttg Leu	tct Ser	gtg Val	tta Leu 645	att Ile	ctc Leu	agc Ser	2032
ctc Leu	ctg Leu 650	ctg Leu	gtt Val	ttc Phe	atc Ile	atg Met 655	tcc Ser	gtc Val	ttc Phe	gtg Val	gcc Ala 660	gcc Ala	Gly 999	ctc Leu	ttc Phe	2080
gtg Val 665	ctg Leu	gtc Val	atg Met	aag Lys	cgc Arg 670	agg Arg	aag Lys	aag Lys	aac Asn	cag Gln 675	agc Ser	gac Asp	cac His	acc Thr	agc Ser 680	2128
acc Thr	aac Asn	aac Asn	tcc Ser	gac Asp 685	gtg Val	agc Ser	tcc Ser	ttt Phe	aac Asn 690	atg Met	cag Gln	tac Tyr	agc Ser	gtg Val 695	tac Tyr	2176
ggc	ggc	ggc Gly	ggc Gly 700	ggc Gly	acg Thr	ggc Gly	ggc Gly	cac His 705	cca Pro	cac His	gcg Ala	cac His	gtg Val 710	cat His	cac His	2224
cgc Arg	Gly 999	ccc Pro 715	gcg Ala	ctg Leu	ccc Pro	aag Lys	gtg Val 720	aag Lys	acg Thr	ccc Pro	gcg Ala	ggc Gly 725	cac His	gtg Val	tat Tyr	2272
gaa Glu	tac Tyr 730	atc Ile	ccc Pro	cac	cca Pro	ctg Leu 735	ggc Gly	cac His	atg Met	tgc Cys	aaa Lys 740	aac Asn	ccc Pro	atc Ile	tac Tyr	2320
cgc Arg 745	tcc Ser	cga Arg	gag Glu	ggc	aac Asn 750	tcc Ser	gta Val	gag Glu	gat Asp	tac Tyr 755	aaa Lys	gac Asp	ctg Leu	cac His	gag Glu 760	2368
ctc Leu	aag Lys	gtc Val	acc Thr	tac Tyr 765	agc Ser	agc Ser	aac Asn	cac His	cac His 770	Leu	cag Gln	cag Gln	cag Gln	cag Gln 775	Gln	2416
ccg Pro	ccg Pro	ccg Pro	cca Pro 780	Pro	cag Gln	cag Gln	cca	cag Gln 785	Gln	cag Gln	ccc	ccg Pro	ecg Pro 790	Gln	ctg Leu	2464
cag Gln	ctg Leu	cag Gln 795	Pro	gly aaa	gag Glu	gag Glu	gag Glu	Arg	cgg Arg	gaa Glu	ago Ser	cac His	His	ttg Lev	cgg	2512
agc Ser	ccc Pro 810	Ala	tac Tyr	ago Ser	gtc Val	ago Ser 815	Thr	ato Ile	gag Glu	ccc Pro	cgg Arg 820	, Glu	gac Asp	cto Lev	ctg Leu	2560

	2608
tcg ccg gtg cag gac gcc gac cgc ttt tac agg ggc att tta gaa cca Ser Pro Val Gln Asp Ala Asp Arg Phe Tyr Arg Gly Ile Leu Glu Pro 825 830 835 840	2608
gac aaa cac tgc tcc acc acc ccc gcc ggc aat agc ctc ccg gaa tat Asp Lys His Cys Ser Thr Thr Pro Ala Gly Asn Ser Leu Pro Glu Tyr 845 850 855	2656
ccc aaa ttc ccg tgc agc ccc gct gct tac act ttc tcc ccc aac tat Pro Lys Phe Pro Cys Ser Pro Ala Ala Tyr Thr Phe Ser Pro Asn Tyr 860 865 870	2704
gac ctg aga cgc ccc cat cag tat ttg cac ccg ggg gca ggg gac agc Asp Leu Arg Arg Pro His Gln Tyr Leu His Pro Gly Ala Gly Asp Ser 875 880 885	2752
agg cta cgg gaa ccg gtg ctc tac agc ccc ccg agt gct gtc ttt gta Arg Leu Arg Glu Pro Val Leu Tyr Ser Pro Pro Ser Ala Val Phe Val 890 895 900	2800
gaa ccc aac cgg aac gaa tat ctg gag tta aaa gca aaa cta aac gtt Glu Pro Asn Arg Asn Glu Tyr Leu Glu Leu Lys Ala Lys Leu Asn Val 905 910 915 920	2848
gag ccg gac tac ctc gaa gtg ctg gaa aaa cag acc acg ttt agc cag Glu Pro Asp Tyr Leu Glu Val Leu Glu Lys Gln Thr Thr Phe Ser Gln 925 930 935	2896
ttc taaaagcaaa gaaactctct tggagctttt gcatttaaaa caaacaagca Phe	2949
agcagacaca cacagtgaac acatttgatt aattgtgttg tttcaacgtt tagggtgaag	3009
tgccttggca cgggatttet cagcttcggt ggaagatacg aaaagggtgt gcaatttcct	3069
ttaaaattta cacgtgggaa acatttgtgt aaactgggca catcactttc tcttcttgcg	3129
tgtggggcag gtgtggagaa gggctttaag gaggccaatt tgctgcgcgg gtgacctgtg	3189
aaaggtcaca gtcatttttg tagtggttgg aagtgctaag aatggtggat gatggcagag	3249
catagattet actetteete ttttgettee tecceeteee eegeceetge eccacetete	3309
tttctcccct tttaagccat gggtgggtct aactggcttt tgtggagaaa ttagcacacc	3369
ccaactttaa taggaaattt gttctctttt tcc	3402

<210> 19

<211> 937

<212> PRT

<213> Homo sapiens

<400> 19

Met Leu Gln Thr Leu Ala Phe Ala Val Thr Ser Leu Val Leu Ser Cys
1 5 10 15

Ala Glu Thr Ile Asp Tyr Tyr Gly Glu Ile Cys Asp Asn Ala Cys Pro 20 25 30

Cys Glu Glu Lys Asp Gly Ile Leu Thr Val Ser Cys Glu Asn Arg Gly 35 40 45

Ile Ile Ser Leu Ser Glu Ile Ser Pro Pro Arg Phe Pro Ile Tyr His 50 55 60

Leu Leu Leu Ser Gly Asn Leu Leu Asn Arg Leu Tyr Pro Asn Glu Phe 65 70 75 80

Val Asn Tyr Thr Gly Ala Ser Ile Leu His Leu Gly Ser Asn Val Ile 85 90 95

Gln Asp Ile Glu Thr Gly Ala Phe His Gly Leu Arg Gly Leu Arg Arg 100 105 110

Leu His Leu Asn Asn Asn Lys Leu Glu Leu Leu Arg Asp Asp Thr Phe 115 120 125

Leu Gly Leu Glu Asn Leu Glu Tyr Leu Gln Val Asp Tyr Asn Tyr Ile 130 135 140

Ser Val Ile Glu Pro Asn Ala Phe Gly Lys Leu His Leu Leu Gln Val 145 150 155 160

Leu Ile Leu Asn Asp Asn Leu Leu Ser Ser Leu Pro Asn Asn Leu Phe 165 170 175

Arg Phe Val Pro Leu Thr His Leu Asp Leu Arg Gly Asn Arg Leu Lys
180
185
190

Leu Leu Pro Tyr Val Gly Leu Leu Gln His Met Asp Lys Val Val Glu 195 200 205

Leu Gln Leu Glu Glu Asn Pro Trp Asn Cys Ser Cys Glu Leu Ile Ser 210 215 220

Leu Lys Asp Trp Leu Asp Ser Ile Ser Tyr Ser Ala Leu Val Gly Asp

													•		
225					230					235					240
Val	Val	Cys	Glu	Thr 245	Pro	Phe	Arg	Leu	His 250	Gly	Arg	Asp	Leu	Asp 255	Glu
Val	Ser	Lys	Gln 260	Glu	Leu	Cys	Pro	Arg 265	Arg	Leu	Ile	Ser	Asp 270	Tyr	Glu
Met	Arg	Pro 275	Gln	Thr	Pro	Leu	Ser 280	Thr	Thr	Gly	Tyr	Leu 285	His	Thr	Thr
Pro	Ala 290	Ser	Val	Asn	Ser	Val 295	Ala	Thr	Ser	Ser	Ser 300	Ala	Val	Tyr	Lys
Pro 305	Pro	Leu	Lys	Pro	Pro 310	Lys	Gly	Thr	Arg	Gln 315	Pro	Asn	Lys	Pro	Arg 320
Val	Arg	Pro	Thr	Ser 325	Arg	Gln	Pro	Ser	Lys 330	Asp	Leu	Gly	Tyr	Ser 335	Asn
Tyr	Gly	Pro	Ser 340	Ile	Ala	Tyr	Gln	Thr 345	Lys	Ser	Pro	Val	Pro 350	Leu	Glu
Cys	Pro	Thr 355	Ala	Cys	Ser	Cys	Asn 360	Leu	Gln	Ile	Ser	Asp 365	Leu	Gly	Leu
Asn	Val 370		Cys	Gln	Glu	Arg 375	Lys	Ile	Glu	Ser	Ile 380	Ala	Glu	Leu	Gln
Pro 385	Lys	Pro	Tyr	Asn	Pro 390	Lys	Lys	Met	Tyr	Leu 395	Thr	Glu	Asn	Tyr	Ile 400
Ala	Val	Val	Arg	Arg 405		Asp	Phe	Leu	Glu 410		Thr	Gly	Leu	Asp 415	Leu
Leu	His	Leu	Gly 420		Asn	Arg	Ile	Ser 425		Ile	Gln	Asp	Arg 430		Phe
Gly	Asp	Leu 435		. Àsn	Leu	Arg	Arg 440		Tyr	· Leu	Asn	Gly 445	Asn	Arg	Ile
Glu	Arg 450		Ser	Pro	Glu	Leu 455		Tyr	Gly	Leu	Gln 460		Leu	Gln	Tyr

37

Leu Phe Leu Gln Tyr Asn Leu Ile Arg Glu Ile Gln Ser Gly Thr Phe 465 470 475 480

Asp Pro Val Pro Asn Leu Gln Leu Leu Phe Leu Asn Asn Asn Leu Leu 495 496

Gln Ala Met Pro Ser Gly Val Phe Ser Gly Leu Thr Leu Leu Arg Leu 500 505 510

Asn Leu Arg Ser Asn His Phe Thr Ser Leu Pro Val Ser Gly Val Leu 515 520 525

Asp Gln Leu Lys Ser Leu Ile Gln Ile Asp Leu His Asp Asn Pro Trp 530 540

Asp Cys Thr Cys Asp Ile Val Gly Met Lys Leu Trp Val Glu Gln Leu 545 550 555 560

Lys Val Gly Val Leu Val Asp Glu Val Ile Cys Lys Ala Pro Lys Lys 565 570 575

Phe Ala Glu Thr Asp Met Arg Ser Ile Lys Ser Glu Leu Leu Cys Pro 580 585 590

Asp Tyr Ser Asp Val Val Val Ser Thr Pro Thr Pro Ser Ser Ile Gln 595 600 605

Val Pro Ala Arg Thr Ser Ala Val Thr Pro Ala Val Arg Leu Asn Ser 610 615 620

Thr Gly Ala Pro Ala Ser Leu Gly Ala Gly Gly Ala Ser Ser Val 625 630 635 640

Pro Leu Ser Val Leu Ile Leu Ser Leu Leu Leu Val Phe Ile Met Ser 645 650 655

Val Phe Val Ala Ala Gly Leu Phe Val Leu Val Met Lys Arg Arg Lys 660 665 670

Lys Asn Gln Ser Asp His Thr Ser Thr Asn Asn Ser Asp Val Ser Ser 675 680 685

Phe Asn Met Gln Tyr Ser Val Tyr Gly Gly Gly Gly Gly Thr Gly Gly 690 695 700

38

His Pro His Ala His Val His His Arg Gly Pro Ala Leu Pro Lys Val 705 710 Lys Thr Pro Ala Gly His Val Tyr Glu Tyr Ile Pro His Pro Leu Gly 730 His Met Cys Lys Asn Pro Ile Tyr Arg Ser Arg Glu Gly Asn Ser Val 740 745 Glu Asp Tyr Lys Asp Leu His Glu Leu Lys Val Thr Tyr Ser Ser Asn His His Leu Gln Gln Gln Gln Pro Pro Pro Pro Pro Gln Gln Pro Gln Gln Gln Pro Pro Pro Gln Leu Gln Leu Gln Pro Gly Glu Glu Glu 795 Arg Arg Glu Ser His His Leu Arg Ser Pro Ala Tyr Ser Val Ser Thr 810 805 Ile Glu Pro Arg Glu Asp Leu Leu Ser Pro Val Gln Asp Ala Asp Arg 820 Phe Tyr Arg Gly Ile Leu Glu Pro Asp Lys His Cys Ser Thr Thr Pro 835 840 Ala Gly Asn Ser Leu Pro Glu Tyr Pro Lys Phe Pro Cys Ser Pro Ala 850 855 860 Ala Tyr Thr Phe Ser Pro Asn Tyr Asp Leu Arg Arg Pro His Gln Tyr 875 880 865 870 Leu His Pro Gly Ala Gly Asp Ser Arg Leu Arg Glu Pro Val Leu Tyr 885 890 Ser Pro Pro Ser Ala Val Phe Val Glu Pro Asn Arg Asn Glu Tyr Leu 900 Glu Leu Lys Ala Lys Leu Asn Val Glu Pro Asp Tyr Leu Glu Val Leu 920 915

Glu Lys Gln Thr Thr Phe Ser Gln Phe

935

<210> 20
<211> 406
<212> DNA
<213> Mus musculus
·
<pre><400> 20 aagaacccca tctaccggtc tcgagaaggc aattccgtgg aggattacaa agacctgcac</pre>
gageteaagg teacttacag cageaaceae cacetgeage ageageegee geegeegeeg
caacageeee ageageagee eeeteegeag atgeagatge ageetgggga ggaggagagg
cgggaaagcc accatttgag gagccccgcc tacagcgtca gcaccatcga gccccgagag
gacctactgt cgccggtgca ggacgctgat cgcttttaca ggggcatttt agagccagac
aaacactgct ccactacccc tgcgggcagc agcctcccag aataccctaa attcccatgc
ageceggetg cttacaettt etececaaae tatgacegtt eggeeg
<210> 21
<211> 135
<212> PRT
<213> Mus musculus
<400> 21
Lys Asn Pro Ile Tyr Arg Ser Arg Glu Gly Asn Ser Val Glu Asp Tyr
1 5 10 15
Lys Asp Leu His Glu Leu Lys Val Thr Tyr Ser Ser Asn His His Leu
20 25 30
Gln Gln Gln Pro Pro Pro Pro Gln Gln Pro Gln Gln Gln Pro Pro 35 40 45
Pro Gln Met Gln Met Gln Pro Gly Glu Glu Glu Arg Arg Glu Ser His 50 55 60
His Leu Arg Ser Pro Ala Tyr Ser Val Ser Thr Ile Glu Pro Arg Glu 65 70 75 80

											•					
Asp 1	Leu	Leu	Ser	Pro 85	Val	Gln	Asp	Ala	Asp 90	Arg	Phe	Tyr	Arg	Gly 95	Ile	
Leu (Glu	Pro	Asp 100	Lys	His	Cys	Ser	Thr 105	Thr	Pro	Ala	Gly	Ser 110	Ser	Leu	
Pro (Glu	Tyr 115	Pro	Lys	Phe	Pro	Cys 120	Ser	Pro	Ala	Ala	Tyr 125	Thr	Phe	Ser	
Pro .	Asn 130	Tyr	Asp	Arg	Ser	Ala 135						i				
<210	> 2	22														
<211	> 3	3545														
<212	> I	ONA														
<213	> 1	Iomo	sapi	iens												
<220	>															
<221	> (CDS														
<222	>	(112))(3	3042)												
<223	>															
<400 ctga		22 att 1	tgcat	ctcag	gg tt	cca	gece	t gc	gttt	ccta	tat	tgac	ccc 1	ttata	acacga	60
cctg	gcg	ete (cagti	tagg	ga gg	gaga	cgtt	g tt	ttgt:	aatc	aac	cacga	aac q		g aaa : Lys	117
cct Pro	tcc Ser	ata Ile 5	gct Ala	gag Glu	atg Met	ctt Leu	cac His 10	aga Arg	gga Gly	agg Arg	atg Met	ttg Leu 15	tgg Trp	ata Ile	att Ile	165
ctt Leu	cta Leu 20	agc Ser	aca Thr	att Ile	gct Ala	cta Leu 25	gga Gly	tgg Trp	act Thr	acc Thr	ccg Pro 30	att Ile	ccc Pro	cta Leu	ata Ile	213
gag Glu 35	gac Asp	tca Ser	gag Glu	gaa Glu	ata Ile 40	gat Asp	gag Glu	ccc Pro	tgt Cys	ttt Phe 45	gat Asp	cca Pro	tgc Cys	tac Tyr	tgt Cys 50	261
gaa Glu	gtt Val	aaa Lys	gaa Glú	agc Ser 55	ctc Leu	ttt Phe	cat His	ata Ile	cat His 60	tgt Cys	gac Asp	agt Ser	aaa Lys	gga Gly 65	ttt Phe	309

aca Thr	aat Asn	att Ile	agt Ser 70	cag Gln	att Ile	acc Thr	gag Glu	ttc Phe 75	tgg Trp	tca Ser	aga Arg	cct Pro	ttt Phe 80	aaa Lys	ctg Leu	357
	ctg Leu															405
cat His	ttg Leu 100	aat Asn	aat Asn	gct Ala	gtg Val	tct Ser 105	att Ile	aat Asn	ctt Leu	ggg Gly	aac Asn 110	aat Asn	gca Ala	ttg Leu	cag Gln	453
	att Ile															501
	cta Leu															549
ggc	ttg Leu	gaa Glu	agt Ser 150	cta Leu	gaa Glu	tat Tyr	ctg Leu	cag Gln 155	gca Ala	gat Asp	tac Tyr	aat Asn	gtc Val 160	att Ile	aaa Lys	597
cgt Arg	att Ile	gag Glu 165	agt Ser	ggg Gly	gca Ala	ttt Phe	cgg Arg 170	aac Asn	cta Leu	agt Ser	aaa Lys	ttg Leu 175	agg Arg	gtt Val	ctg Leu	645
att Ile	tta Leu 180	aat Asn	gat Asp	aat Asn	ctc Leu	atc Ile 185	ccc Pro	atg Met	ctt Leu	cca Pro	acc Thr 190	aat Asn	tta Leu	ttt Phe	aag Lys	693
	gtc Val															741
ctt Leu	ttt Phe	tac Tyr	cga Arg	gga Gly 215	atg Met	cta Leu	gat Asp	cac His	att Ile 220	ggc Gly	aga Arg	agc Ser	ctg Leu	atg Met 225	gag Glu	789
ctc Leu	cag Gln	ctg Leu	gaa Glu 230	gaa Glu	aac Asn	cct Pro	tgg Trp	aac Asn 235	tgt Cys	aca Thr	tgt Cys	gaa Glu	att Ile 240	Val	caa Gln	837
ctg Leu	aag Lys	agt Ser 245	Trp	ctg Leu	gaa Glu	cgc Arg	att Ile 250	Pro	tat Tyr	act Thr	gcc Ala	ctg Leu 255	Val	gga Gly	gac Asp	885
att	acc Thr 260	Cys	gag Glu	acc Thr	cct Pro	ttc Phe 265	His	ttc Phe	cat His	gga Gly	aag Lys 270	Asp	cta Leu	cga Arg	gaa Glu	933
ato Ile 275	agg Arg	aag Lys	aca Thr	gaa Glu	ctc Leu 280	tgt Cys	ccc Pro	ttg Leu	ttg Leu	tct Ser 285	Asp	tct Ser	gag Glu	gta Val	gag Glu 290	981
	agt Ser															1029

				295					300					305		
cca Pro	act Thr	aag Lys	cct Pro 310	tcc Ser	tca Ser	atg Met	cta Leu	tcc Ser 315	tct Ser	gtt Val	cat His	ttt Phe	act Thr 320	gct Ala	tct Ser	1077
tct Ser	gtc Val	gaa Glu 325	tac Tyr	aag Lys	tcc Ser	tca Ser	aat Asn 330	aaa Lys	cag Gln	cct Pro	aag Lys	ccc Pro 335	acc Thr	aaa Lys	cag Gln	1125
cct Pro	cga Arg 340	aca Thr	cca Pro	agg Arg	cca Pro	ccc Pro 345	tcc Ser	acc Thr	tcc Ser	caa Gln	gct Ala 350	tta Leu	tat Tyr	cct Pro	ggt Gly	1173
cca Pro 355	aac Asn	cag Gln	cct Pro	ccc Pro	att Ile 360	gct Ala	cct Pro	tat Tyr	cag Gln	acc Thr 365	aga Arg	cca Pro	cca Pro	atc Ile	ccc Pro 370	1221
att Ile	ata Ile	tgc Cys	ccc Pro	act Thr 375	ggg Gly	tgt Cys	acc Thr	tgt Cys	aat Asn 380	ttg Leu	cac His	atc Ile	aat Asn	gac Asp 385	ctt Leu	1269
ggc	ttg Leu	act Thr	gtc Val 390	aac Asn	tgc Cys	aaa Lys	gag Glu	cga Arg 395	gga Gly	ttt Phe	aat Asn	aac Asn	att Ile 400	tct Ser	gaa Glu	1317
ctt Leu	ctt Leu	cca Pro 405	agg Arg	ccc Pro	ttg Leu	aat Asn	gcc Ala 410	aag Lys	aaa Lys	ctg Leu	tat Tyr	ctg Leu 415	agt Ser	agc Ser	aat Asn	1365
ctg Leu	att Ile 420	cag Gln	aaa Lys	ata Ile	tac Tyr	cgt Arg 425	tct Ser	gat Asp	ttt Phe	tgg Trp	aat Asn 430	ttt Phe	tct Ser	tcc Ser	ttg Leu	1413
gat Asp 435	ctc Leu	ttg Leu	cat His	ctg Leu	999 Gly 440	aac Asn	aat Asn	cgt Arg	att Ile	tcc Ser 445	tat Tyr	gtc Val	caa Gln	gat Asp	999 Gly 450	1461
gcc Ala	ttt Phe	atc Ile	aac Asn	ttg Leu 455	ccc Pro	aac Asn	tta Leu	aag Lys	agc Ser 460	ctc Leu	ttc Phe	ctt Leu	aat Asn	ggc Gly 465	aac Asn	1509
gat Asp	ata Ile	gag Glu	aag Lys 470	ctg Leu	aca Thr	cca Pro	ggc Gly	atg Met 475	ttc Phe	cga Arg	ggc	cta Leu	cag Gln 480	agt Ser	ttg Leu	1557
cac His	tac Tyr	ttg Leu 485	.Tyr	ttt Phe	gag Glu	ttc Phe	aat Asn 490	gtc Val	atc Ile	cgg Arg	gaa Glu	atc Ile 495	Gln	cct Pro	gca Ala	1605
gcc Ala	ttc Phe 500	agc Ser	ctc Leu	atg Met	Pro	aac Asn 505	ttg Leu	aag Lys	ctg Leu	cta Leu	ttc Phe 510	Leu	aat Asn	aat Asn	aac Asn	1653
tta Leu 515	Leu	agg Arg	act Thr	ctg Leu	cca Pro 520	Thr	gac Asp	gcc Ala	ttt Phe	gct Ala 525	Gly	aca Thr	tcc Ser	ctg Leu	gcc Ala 530	1701
cgg	ctc	aac	ctg	agg	aag	aac	tac	ttc	ctc	tat	ctt	ccc	gtg	gct	ggt	1749

Arg	Leu	Asn	Leu	Arg 535	Lys	Asn	Tyr	Phe	Leu 540	Tyr	Leu	Pro	Val	Ala 545	Gly	
gtc Val																1797
cct Pro	tgg Trp	gac Asp 565	tgc Cys	acc Thr	tgt Cys	gac Asp	ctg Leu 570	gtc Val	ccc Pro	ttt Phe	aaa Lys	cag Gln 575	tgg Trp	atc Ile	gaa Glu	1845
											ctt Leu 590					1893
											gag Glu					1941
											gaa Glu					1989
											agt Ser					2037
											tct Ser					2085
agc Ser	ctg Leu 660	ctg Leu	gtt Val	ctg Leu	ttt Phe	ttc Phe 665	tca Ser	gca Ala	gtc Val	ttt Phe	gtt Val 670	gct Ala	gca Ala	ggc Gly	ctc Leu	2133
ttt Phe 675	gcc Ala	tac Tyr	gtg Val	ctc Leu	cga Arg 680	agg Arg	cgt Arg	cga Arg	aag Lys	aag Lys 685	ctg Leu	ccc Pro	ttc Phe	aga Arg	agc Ser 690	2181
aag Lys	cgg Arg	cag Gln	gaa Glu	ggt Gly 695	gtg Val	gac Asp	ctt Leu	act Thr	ggc Gly 700	atc Ile	caa Gln	atg Met	caa Gln	tgc Cys 705	cac His	2229
agg Arg	ctg Leu	ttt Phe	gag Glu 710	gat Asp	ggt Gly	gga Gly	ggt Gly	ggt Gly 715	ggt Gly	ggc Gly	gga Gly	agt Ser	999 Gly 720	ggt Gly	ggt Gly	2277
ggt Gly	cga Arg	cca Pro 725	act Thr	ctt Leu	tcc Ser	tct Ser	cca Pro 730	Glu	aag Lys	gcc Ala	cct Pro	ccc Pro 735	.Val	ggt Gly	cat His	 2325
gtg Val	tat Tyr 740	gag Glu	tac Tyr	atc Ile	ccc Pro	cac His 745	ccg Pro	gtt Val	acc Thr	caa Gln	atg Met 750	tgc Cys	aac Asn	aac Asn	ccc Pro	2373
atc Ile 755	tac Tyr	aag Lys	cct Pro	cgt Arg	gag Glu 760	Glu	gag Glu	gag Glu	gtg Val	gct Ala 765		tca Ser	tca Ser	gcc Ala	caa Gln 770	2421

gaa gca ggg a Glu Ala Gly s						2469
atg ggt gag g Met Gly Glu A	gct ctc cta Ala Leu Leu 790	Gly Ser G	ag cag tti lu Gln Phe 95	gct gag ac Ala Glu Th	r Pro Lys	2517
gag aac cat a Glu Asn His 8 805						2565
gcc cta gca g Ala Leu Ala V 820	gtg tcc agc Val Ser Ser	tcc cag c Ser Gln L 825	ett aac acc eu Asn Thi	e ata gtg ac Tle Val Th 830	g gtg aat r Val Asn	2613
cac cat cac o His His His I 835				/ Val Ser Gl		2661
ggg gga act g Gly Gly Thr (ggg gga gac Gly Gly Asp 855	ttg gca g Leu Ala G	ggg ttc cgg Gly Phe Arg 860	c cac cat ga g His His Gl	g aaa aat u Lys Asn 865	2709
ggt ggg gtg g		Pro Pro G			er Gly Ser	2757
atg cta cta g Met Leu Leu 2 885	gat cga gag Asp Arg Glu	agg cca c Arg Pro G 890	ag eet ge Eln Pro Ala	c ccc tgc ac a Pro Cys Th 895	a gtg gga ar Val Gly	2805
ttt gtg gac t Phe Val Asp (900	tgt ctc tat Cys Leu Tyr	gga aca g Gly Thr V 905	gtg ccc aaa Val Pro Lys	a tta aag ga s Leu Lys Gl 910	a ctg cac u Leu His	2853
gtg cac cct o Val His Pro 1 915				ı Gln Gln As		2901
ctc aaa gaa a Leu Lys Glu '						2949
caa acc caa a Gln Thr Gln	aaa agt gat Lys Ser Asp 950	Tyr Leu G	gag tta ag Glu Leu Ar 955	g gcc aaa cf g Ala Lys Lo 99	eu Gln Thr	2997
aag ccg gat Lys Pro Asp ' 965	tac ctc gaa Tyr Leu Glu	gtc ctg g Val Leu G 970	gag aag ac Glu Lys Th	a aca tac ag r Thr Tyr A 975	gġ ttc cg Phe	3042
taacagagag a	agaaaatat a	ttagtgctt	ttttttt	c aaaagaaaa	g gaaaataaaa	3102
gaaatatatc c	cttgctccc t	ttacacttg	tcccagtaa	c tecatecte	a cgatctttcc	3162
taccctgaac a	aaactaaaa c	cgcatgata	actagagaa	t acagatgta	t gctctcccct	3222
ctcagatgcg a	itttggagga a	gggccatac	tcagatcat	t aatcaatga	a agtgccttcg	3282

45

cagacttttg	ccagcaaatg	ttatcattat	ttttttatac	tgaaacttga	gactttgact	3342
gtgccatgta	taagatatac	tggggatcat	tgtatggatc	ctaattaagt	aaaattcaat	3402
gtgtcttttt	attttcagta	actattttt	ttatagttgt	agttttgatt	taaagggggg	3462
gaaacaagtt	gacatttgtc	atttgtggct	ttctttctta	tcatcatggc	acagattctg	3522
tacatgtatt	aacaatgcag	ttt				3545

<210> 23

<211> 977

<212> PRT

<213> Homo sapiens

<400> 23

Met Lys Pro Ser Ile Ala Glu Met Leu His Arg Gly Arg Met Leu Trp 1 5 10 .15

Ile Ile Leu Leu Ser Thr Ile Ala Leu Gly Trp Thr Thr Pro Ile Pro 20 25 30

Leu Ile Glu Asp Ser Glu Glu Ile Asp Glu Pro Cys Phe Asp Pro Cys 35 40 45

Tyr Cys Glu Val Lys Glu Ser Leu Phe His Ile His Cys Asp Ser Lys 50 55 60

Gly Phe Thr Asn Ile Ser Gln Ile Thr Glu Phe Trp Ser Arg Pro Phe 65 70 75 80

Lys Leu Tyr Leu Gln Arg Asn Ser Met Arg Lys Leu Tyr Thr Asn Ser 85 90 95

Phe Leu His Leu Asn Asn Ala Val Ser Ile Asn Leu Gly Asn Asn Ala 100 105 110

Leu Gln Asp Ile Gln Thr Gly Ala Phe Asn Gly Leu Lys Ile Leu Lys
115 120 125

Arg Leu Tyr Leu His Glu Asn Lys Leu Asp Val Phe Arg Asn Asp Thr 130 135 140

Phe Leu Gly Leu Glu Ser Leu Glu Tyr Leu Gln Ala Asp Tyr Asn Val Ile Lys Arg Ile Glu Ser Gly Ala Phe Arg Asn Leu Ser Lys Leu Arg . 170 Val Leu Ile Leu Asn Asp Asn Leu Ile Pro Met Leu Pro Thr Asn Leu 185 Phe Lys Ala Val Ser Leu Thr His Leu Asp Leu Arg Gly Asn Arg Leu 200 Lys Val Leu Phe Tyr Arg Gly Met Leu Asp His Ile Gly Arg Ser Leu 215 Met Glu Leu Gln Leu Glu Glu Asn Pro Trp Asn Cys Thr Cys Glu Ile 235 230 Val Gln Leu Lys Ser Trp Leu Glu Arg Ile Pro Tyr Thr Ala Leu Val 245 Gly Asp Ile Thr Cys Glu Thr Pro Phe His Phe His Gly Lys Asp Leu 265 270 Arg Glu Ile Arg Lys Thr Glu Leu Cys Pro Leu Leu Ser Asp Ser Glu 275 280 285 Val Glu Ala Ser Leu Gly Ile Pro His Ser Ser Ser Lys Glu Asn 290 295 Ala Trp Pro Thr Lys Pro Ser Ser Met Leu Ser Ser Val His Phe Thr 305 310 315 Ala Ser Ser Val Glu Tyr Lys Ser Ser Asn Lys Gln Pro Lys Pro Thr 325 330 Lys Gln Pro Arg Thr Pro Arg Pro Pro Ser Thr Ser Gln Ala Leu Tyr 340 - 345 Pro Gly Pro Asn Gln Pro Pro Ile Ala Pro Tyr Gln Thr Arg Pro Pro 355 360 Ile Pro Ile Ile Cys Pro Thr Gly Cys Thr Cys Asn Leu His Ile Asn 370 375

Asp Leu Gly Leu Thr Val Asn Cys Lys Glu Arg Gly Phe Asn Asn Ile 385 390 395 400

Ser Glu Leu Leu Pro Arg Pro Leu Asn Ala Lys Lys Leu Tyr Leu Ser 405 410 415

Ser Asn Leu Ile Gln Lys Ile Tyr Arg Ser Asp Phe Trp Asn Phe Ser 420 425 430

Ser Leu Asp Leu Leu His Leu Gly Asn Asn Arg Ile Ser Tyr Val Gln 435 440 445

Asp Gly Ala Phe Ile Asn Leu Pro Asn Leu Lys Ser Leu Phe Leu Asn 450 455 460

Gly Asn Asp Ile Glu Lys Leu Thr Pro Gly Met Phe Arg Gly Leu Gln 465 470 475 480

Ser Leu His Tyr Leu Tyr Phe Glu Phe Asn Val Ile Arg Glu Ile Gln 485 490 495

Pro Ala Ala Phe Ser Leu Met Pro Asn Leu Lys Leu Leu Phe Leu Asn 500 505 510

Asn Asn Leu Leu Arg Thr Leu Pro Thr Asp Ala Phe Ala Gly Thr Ser 515 520 525

Leu Ala Arg Leu Asn Leu Arg Lys Asn Tyr Phe Leu Tyr Leu Pro Val 530 535 540

Ala Gly Val Leu Glu His Leu Asn Ala Ile Val Gln Ile Asp Leu Asn 545 550 555 560

Glu Asn Pro Trp Asp Cys Thr Cys Asp Leu Val Pro Phe Lys Gln Trp 565 570 575

Ile Glu Thr Ile Ser Ser Val Ser Val Val Gly Asp Val Leu Cys Arg 580 585 590

Ser Pro Glu Asn Leu Thr His Arg Asp Val Arg Thr Ile Glu Leu Glu 595 600 605

Val Leu Cys Pro Glu Met Leu His Val Ala Pro Ala Gly Glu Ser Pro

48

610	615	620

Ala Gln Pro Gly Asp Ser His Leu Ile Gly Ala Pro Thr Ser Ala Ser 625 630 635 640

Pro Tyr Glu Phe Ser Pro Pro Gly Gly Pro Val Pro Leu Ser Val Leu 645 650 655

Ile Leu Ser Leu Leu Val Leu Phe Phe Ser Ala Val Phe Val Ala Ala 660 670

Gly Leu Phe Ala Tyr Val Leu Arg Arg Arg Lys Lys Leu Pro Phe 675 680 685

Arg Ser Lys Arg Gln Glu Gly Val Asp Leu Thr Gly Ile Gln Met Gln 690 695 700

Cys His Arg Leu Phe Glu Asp Gly Gly Gly Gly Gly Gly Gly Ser Gly 705 710 715 720

Gly Gly Arg Pro Thr Leu Ser Ser Pro Glu Lys Ala Pro Pro Val
725 730 735

Gly His Val Tyr Glu Tyr Ile Pro His Pro Val Thr Gln Met Cys Asn 740 745 750

Asn Pro Ile Tyr Lys Pro Arg Glu Glu Glu Glu Val Ala Val Ser Ser

Ala Gln Glu Ala Gly Ser Ala Glu Arg Gly Gly Pro Gly Thr Gln Pro 770 775 780

Pro Gly Met Gly Glu Ala Leu Leu Gly Ser Glu Gln Phe Ala Glu Thr 785 790 795 800

Pro Lys Glu Asn His Ser Asn Tyr Arg Thr Leu Leu Glu Lys Glu Lys 805 810 815

Glu Trp Ala Leu Ala Val Ser Ser Ser Gln Leu Asn Thr Ile Val Thr 820 825 830

Val Asn His His Pro His His Pro Ala Val Gly Gly Val Ser Gly 835 840 845

49

Val Val Gly Gly Thr Gly Gly Asp Leu Ala Gly Phe Arg His His Glu

Lys Asn Gly Gly Val Val Leu Phe Pro Pro Gly Gly Gly Cys Gly Ser 875

Gly Ser Met Leu Leu Asp Arg Glu Arg Pro Gln Pro Ala Pro Cys Thr 890 885

Val Gly Phe Val Asp Cys Leu Tyr Gly Thr Val Pro Lys Leu Lys Glu 905

Leu His Val His Pro Pro Gly Met Gln Tyr Pro Asp Leu Gln Gln Asp 920

Ala Arg Leu Lys Glu Thr Leu Leu Phe Ser Ala Glu Lys Gly Phe Thr

Asp His Gln Thr Gln Lys Ser Asp Tyr Leu Glu Leu Arg Ala Lys Leu 950 955

Gln Thr Lys Pro Asp Tyr Leu Glu Val Leu Glu Lys Thr Thr Tyr Arg 965

Phe

<210> 24

<211> 2631

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (118)..(2628)

<223>

<400> 24 atgatttaca tacaagtaat ttttcaagta atgaccattg aaaaaatgtt ttcttttat 60

ttt	taga	tt a	tttc	tctt	t at	tcag	aago	ata	cagt	tgt	ttgo	tgat	tg c	aaga	.ag	117
atg Met 1	ttt Phe	ctg Leu	tgg Trp	ctg Leu 5	ttt Phe	ctg Leu	att Ile	ttg Leu	tca Ser 10	gcc Ala	ctg Leu	att Ile	tct Ser	tcg Ser 15	aca Thr	165
aat Asn	gca Ala	gat Asp	tct Ser 20	gac Asp	ata Ile	tcg Ser	gtg Val	gaa Glu 25	att Ile	tgc Cys	aat Asn	gtg Val	tgt Cys 30	tcc Ser	tgc Cys	213
gtg Val	tca Ser	gtt Val 35	gag Glu	aat Asn	gtg Val	ctc Leu	tat Tyr 40	gtc Val	aac Asn	tgt Cys	gag Glu	aag Lys 45	gtt Val	tca Ser	gtc Val	261
tac Tyr	aga Arg 50	cca Pro	aat Asn	cag Gln	ctg Leu	aaa Lys 55	cca Pro	cct Pro	tgg Trp	tct Ser	aat Asn 60	ttt Phe	tat Tyr	cac His	ctc Leu	309
aat Asn 65	ttc Phe	caa Gln	aat Asn	aat Asn	ttt Phe 70	tta Leu	aat Asn	att Ile	ctg Leu	tat Tyr 75	cca Pro	aat Asn	aca Thr	ttc Phe	ttg Leu 80	357
aat Asn	ttt Phe	tca Ser	cat His	gca Ala 85	gtc Val	tcc Ser	ctg Leu	cat His	ctg Leu 90	gly ggg	aat Asn	aat Asn	aaa Lys	ctg Leu 95	cag Gln	405
aac Asn	att Ile	gag Glu	gga Gly 100	gga Gly	gcc Ala	ttt Phe	ctt Leu	999 Gly 105	ctc Leu	agt Ser	gca Ala	tta Leu	aag Lys 110	cag Gln	ttg Leu	453
cac His	ttg Leu	aac Asn 115	aac Asn	aat Asn	gaa Glu	tta Leu	aag Lys 120	att Ile	ctc Leu	cga Arg	gct Ala	gac Asp 125	act Thr	ttc Phe	ctt Leu	501
ggc	ata Ile 130	gag Glu	aac Asn	ttg Leu	gag Glu	tat Tyr 135	ctc Leu	cag Gln	gct Ala	gac Asp	tac Tyr 140	aat Asn	tta Leu	atc Ile	aag Lys	549
tat Tyr 145	att Ile	gaa Glu	cga Arg	gga Gly	gcc Ala 150	ttc Phe	aat Asn	aag Lys	ctc Leu	cac His 155	ГЛS	ctg Leu	aaa Lys	gtt Val	ctc Leu 160	597
att Ile	ctt Leu	aat Asn	gac Asp	aat Asn 165	ctg Leu	att Ile	tca Ser	ttc Phe	ctt Leu 170	Pro	gat Asp	aat Asn	att Ile	ttc Phe 175	cga Arg	645
ttc Phe	gca Ala	tct Ser	ttg Leu 180	acc Thr	cat His	ctg Leu	gat Asp	ata Ile 185	Arg	gly ggg	aac Asn	aga Arg	atc Ile 190	Gln	aag Lys	693
ctc Leu	cct Pro	tat Tyr 195	Ile	gly ggg	gtt Val	ctg Leu	gaa Glu 200	His	att	ggc	cgt Arg	gtc Val 205	gtt Val	gaa Glu	ttg Leu	741
caa Gln	ctg Leu 210	Glu	gat Asp	aac Asn	cct Pro	tgg Trp 215	aac Asn	tgt Cys	ago Ser	tgt Cys	gat Asp 220	Leu	ttg Leu	ccc Pro	tta Leu	789
aaa	gct	tgg	ctg	gag	aac	atg	cca	tat	aac	att	: tac	ata	gga	gaa	gct	837

Lys 225	Ala	Trp	Leu	Glu	Asn 230	Met	Pro	Tyr	Asn	Ile 235	Tyr	Ile	Gly	Glu	Ala 240		
atc Ile	tgt Cys	gaa Glu	act Thr	ccc Pro 245	agt Ser	gac Asp	tta Leu	tat Tyr	gga Gly 250	agg Arg	ctt Leu	tta Leu	aaa Lys	gaa Glu 255	acc Thr	885	
aac Asn	aaa Lys	caa Gln	gag Glu 260	cta Leu	tgt Cys	ccc Pro	atg Met	ggc Gly 265	acc Thr	ggc Gly	agt Ser	gat Asp	ttt Phe 270	gac Asp	gtġ Val	933	ı
cgc Arg	atc Ile	ctg Leu 275	cct Pro	cca Pro	tct Ser	cag Gln	ctg Leu 280	gaa Glu	aat Asn	ggc Gly	tac Tyr	acc Thr 285	act Thr	ccc Pro	aat Asn	981	•
ggt Gly	cac His 290	act Thr	acc Thr	caa Gln	aca Thr	tct Ser 295	tta Leu	cac His	aga Arg	tta Leu	gta Val 300	act Thr	aaa Lys	cca Pro	cca Pro	1029	,
aaa Lys 305	aca Thr	aca Thr	aat Asn	cct Pro	tcc Ser 310	aag Lys	atc Ile	tct Ser	gga Gly	atc Ile 315	gtt Val	gca Ala	ggc Gly	aaa Lys	gcc Ala 320	1077	,
ctc Leu	tcc Ser	aac Asn	cgc Arg	aat Asn 325	ctc Leu	agt Ser	cag Gln	att Ile	gtg Val 330	tct Ser	tac Tyr	caa Gln	aca Thr	agg Arg 335	gtg Val	1125	5
cct Pro	cct Pro	cta Leu	aca Thr 340	cct Pro	tgc Cys	ccg Pro	gca Ala	cct Pro 345	tgc Cys	ttc Phe	tgc Cys	aaa Lys	aca Thr 350	cac His	cct Pro	1173	3
tca Ser	gat Asp	ttg Leu 355	gga Gly	cta Leu	agt Ser	gtg Val	aac Asn 360	tgc Cys	caa Gln	gag Glu	aaa Lys	aat Asn 365	ata Ile	cag Gln	tct Ser	122	L
atg Met	tct Ser 370	gaa Glu	ctg Leu	ata Ile	ccg Pro	aaa Lys 375	cct Pro	tta Leu	aat Asn	gcg Ala	aag Lys 380	aag Lys	ctg Leu	cac His	gtc Val	1269	Э
aat Asn 385	ggc	aat Asn	agc Ser	atc Ile	aag Lys 390	Asp	gtg Val	gac Asp	gta Val	tca Ser 395	gac Asp	ttc Phe	act Thr	gac Asp	ttt Phe 400	131	7
gaa Glu	gga Gly	ctg Leu	gat Asp	ttg Leu 405	Leu	cat His	tta Leu	ggc	agc Ser 410	aat Asn	caa Gln	att Ile	aca Thr	gtg Val 415	Ile	136	5
aag Lys	gga Gly	gac Asp	gta Val 420	Phe	cac His	aat Asn	ctc Leu	act Thr 425	Asn	tta Leu	cgc	agg	cta Leu 430	Tyr	ctc Leu	141	3
aat Asn	ggc	aat Asn 435	Gln	att Ile	gag Glu	aga Arg	cto Leu 440	Туг	cct Pro	gaa Glu	ata Ile	ttt Phe 445	Ser	ggt	ctt Leu	146	1
cat His	aac Asn 450	. Leu	cag Gln	tat Tyr	ctg Lev	tat Tyr 455	Leu	gaa Glu	tac Tyr	aat Asn	ttg Lev 460	ı Ile	aag Lys	gaa Glu	atc Ile	150	9

tca Ser 465	gca Ala	ggc Gly	acc Thr	ttt Phe	gac Asp 470	tcc Ser	atg Met	cca Pro	aat Asn	ttg Leu 475	cag Gln	tta Leu	ctg Leu	tac Tyr	tta Leu 480		1557
aac Asn	aat Asn	aat Asn	ctc Leu	cta Leu 485	aag Lys	agc Ser	ctg Leu	cct Pro	gtt Val 490	tac Tyr	atc Ile	ttt Phe	tcc Ser	gga Gly 495	gca Ala		1605
ccc Pro	tta Leu	gct Ala	aga Arg 500	ctg Leu	aac Asn	ctg Leu	agg Arg	aac Asn 505	aac Asn	aaa Lys	ttc Phe	atg Met	tac Tyr 510	ctg Leu	cct Pro		1653
gtc Val	agt Ser	999 Gly 515	gtc Val	ctt Leu	gat Asp	cag Gln	ttg Leu 520	caa Gln	tct Ser	ctt Leu	aca Thr	cag Gln 525	att Ile	gac Asp	ttg Leu		1701
gag Glu	ggc Gly 530	aac Asn	cca Pro	tgg Trp	gac Asp	tgt Cys 535	act Thr	tgt Cys	gac Asp	ttg Leu	gtg Val 540	gca Ala	tta Leu	aag Lys	ctg Leu		1749
tgg Trp 545	gtg Val	gag Glu	aag Lys	ttg Leu	agc Ser 550	gac Asp	Gly 899	att Ile	gtt Val	gtg Val 555	aaa Lys	gaa Glu	ctg Leu	aaa Lys	tgt Cys 560		1797
gag Glu	acg Thr	cct Pro	gtt Val	cag Gln 565	ttt Phe	gcc Ala	aac Asn	att Ile	gaa Glu 570	ctg Leu	aag Lys	tcc Ser	ctc Leu	aaa Lys 575	aat Asn		1845
gaa Glu	atc Ile	tta Leu	tgt Cys 580	ccc Pro	aaa Lys	ctt Leu	tta Leu	aat Asn 585	aag Lys	ccg Pro	tct Ser	gca Ala	cca Pro 590	ttc Phe	aca Thr	•	1893
agc Ser	cct Pro	gca Ala 595	cct Pro	gcc Ala	att Ile	aca Thr	ttc Phe 600	acc Thr	act Thr	cct Pro	ttg Leu	ggt Gly 605	Pro	att Ile	cga Arg		1941
agt Ser	cct Pro 610	Pro	ggt Gly	ggg Gly	cca Pro	gtg Val 615	cct Pro	ctg Leu	tct Ser	att Ile	tta Leu 620	atc Ile	tta Leu	agt Ser	atc Ile		1989
tta Leu 625	Val	gtc Val	ctc Leu	att Ile	tta Leu 630	acg Thr	gtg Val	ttt Phe	gtt Val	gct Ala 635	ttt Phe	tgc Cys	ctt Leu	ctt Leu	gtt Val 640		2037
ttt Phe	gtc Val	ctg Leu	cga Arg	cgc Arg 645	Asn	aag Lys	aaa Lys	ccc Pro	aca Thr 650	Val	aag Lys	cac His	gaa Glu	ggc Gly 655	ctg Leu		2085
gly ggg	aat Asn	cct Pro	gac Asp 660	Cys	ggc	tcc Ser	atg Met	Gln 665	Leu	cag Gln	cta Leu	agg Arg	aag Lys 670	His	gac Asp	•	. 2133
cac His	aaa Lys	acc Thr 675	Asn	aaa Lys	aaa Lys	gat Asp	gga Gly 680	Leu	ago Ser	aca Thr	gaa Glu	gct Ala 685	Phe	att Ile	cca Pro		2181
caa Gln	act Thr 690	Ile	gaa Glu	cag Gln	atg Met	ago Ser 695	. Lys	ago Ser	cac His	act Thr	tgt Cys 700	Gly	ttg Leu	aaa Lys	gag Glu	•	2229

53

tca gaa act ggg ttc atg ttt tca gat cct cca gga cag aaa gtt gtt
Ser Glu Thr Gly Phe Met Phe Ser Asp Pro Pro Gly Gln Lys Val Val
705 710 715 720

atg aga aat gtg gcc gac aag gag aaa gat tta tta cat gta gat acc
Met Arg Asn Val Ala Asp Lys Glu Lys Asp Leu Leu His Val Asp Thr

agg aag aga ctg agc aca att gat gag ctg gat gaa tta ttc cct agc
Arg Lys Arg Leu Ser Thr Ile Asp Glu Leu Asp Glu Leu Phe Pro Ser
740 745 750

730

agg gat tcc aat gtg ttt att cag aat ttt ctt gaa agc aaa aag gag
Arg Asp Ser Asn Val Phe Ile Gln Asn Phe Leu Glu Ser Lys Lys Glu

tat aat agc ata ggt gtc agt ggc ttt gag atc cgc tat cca gaa aaa 2469
Tyr Asn Ser Ile Gly Val Ser Gly Phe Glu Ile Arg Tyr Pro Glu Lys
770 780

Caa cca gac aaa aaa agt aag aag tca ctg ata ggt ggc aac cac agt

Gln Pro Asp Lys Lys Ser Lys Lys Ser Leu Ile Gly Gly Asn His Ser

785 790 795 800

aaa att gtt gtg gaa caa agg aag agt gag tat ttt gaa ctg aag gcg
Lys Ile Val Val Glu Gln Arg Lys Ser Glu Tyr Phe Glu Leu Lys Ala
805
810
815

aaa ctg cag agt tcc cct gac tac cta cag gtc ctt gag gag caa aca
Lys Leu Gln Ser Ser Pro Asp Tyr Leu Gln Val Leu Glu Glu Gln Thr
820 825 830

gct ttg aac aag atc tag 2631 Ala Leu Asn Lys Ile 835

<210> 25

<211> 837

<212> PRT

<213> Homo sapiens

<400> 25

Met Phe Leu Trp Leu Phe Leu Ile Leu Ser Ala Leu Ile Ser Ser Thr 1 5 10 15

Asn Ala Asp Ser Asp Ile Ser Val Glu Ile Cys Asn Val Cys Ser Cys 20 25 30

Val Ser Val Glu Asn Val Leu Tyr Val Asn Cys Glu Lys Val Ser Val

		35					40					45			
Tyr	Arg 50	Pro	Asn	Gln	Leu	Lys 55	Pro	Pro	Trp	Ser	Asn 60	Phe	Tyr	His	Leu
Asn 65	Phe	Gln	Asn	Asn	Phe 70	Leu	Asn	Ile	Leu	Tyr 75	Pro	Asn	Thr	Phe	Leu 80
Asn	Phe	Ser	His	Ala 85	Val	Ser	Leu	His	Leu 90	Gly	Asn	Asn	Lys	Leu 95	Gln
Asn	Ile	Glu	Gly 100	Gly	Ala	Phe	Leu	Gly 105	Leu	Ser	Ala	Leu	Lys 110	Gln	Leu
His	Leu	Asn 115	Asn	Asn	Glu	Leu	Lys 120	Ile	Leu	Arg	Ala	Asp 125	Thr	Phe	Leu
Gly	Ile 130	Glu	Asn	Leu	Glu	Tyr 135	Leu	Gln	Ala	Asp	Tyr 140	Asn	Leu	Ile	Lys
Tyr 145	Ile	Glu	Arg	Gly	Ala 150	Phe	Asn	Lys	Leu	His 155	Lys	Leu	ГÀЗ	Val	Leu 160
Ile	Leu	Asn	Asp	Asn 165	Leu	Ile	Ser	Phe	Leu 170	Pro	Asp	Asn	Ile	Phe 175	Arg
Phe	Ala	Ser	Leu 180	Thr	His	Leu	Asp	Ile 185	Arg	Gly	Asn	Arg	Ile 190	Gln	Lys
Leu	Pro	Tyr 195	Ile	Gly	Val	Leu	Glu 200	His	Ile	Gly	Arg	Val 205	Val	Glu	Leu
Gln	Leu 210	Glu	Asp	Asn	Pro	Trp 215	Asn	Cys	Ser	Cys	Asp 220	Leu	Leu	Pro	Leu
Lys 225		Trp	Leu	Glu	Asn 230	Meț	Pro	Tyr	Asn	Ile 235		Ile	Gly	Glu	Ala 240
Ile	Cys	Glu	Thr	Pro 245	Ser	Asp	Leu	Tyr	Gly 250	Arg	Leu	Leu	Lys	Glu 255	Thr
Asn	Lys	Gln	Glu 260	Leu	Cys	Pro	Met	Gly 265	Thr	Gly	Ser	Asp	Phe 270		Val

55

Arg Ile Leu Pro Pro Ser Gln Leu Glu Asn Gly Tyr Thr Thr Pro Asn 275 280 285

Gly His Thr Thr Gln Thr Ser Leu His Arg Leu Val Thr Lys Pro Pro 290 295 300

Lys Thr Thr Asn Pro Ser Lys Ile Ser Gly Ile Val Ala Gly Lys Ala 305 310 315

Leu Ser Asn Arg Asn Leu Ser Gln Ile Val Ser Tyr Gln Thr Arg Val
325 330 335

Pro Pro Leu Thr Pro Cys Pro Ala Pro Cys Phe Cys Lys Thr His Pro 340 345 350

Ser Asp Leu Gly Leu Ser Val Asn Cys Gln Glu Lys Asn Ile Gln Ser 355 360 365

Met Ser Glu Leu Ile Pro Lys Pro Leu Asn Ala Lys Lys Leu His Val 370 375 380

Asn Gly Asn Ser Ile Lys Asp Val Asp Val Ser Asp Phe Thr Asp Phe 385 390 395 400

Glu Gly Leu Asp Leu Leu His Leu Gly Ser Asn Gln Ile Thr Val Ile 405 410 415

Lys Gly Asp Val Phe His Asn Leu Thr Asn Leu Arg Arg Leu Tyr Leu 420 425 430

Asn Gly Asn Gln Ile Glu Arg Leu Tyr Pro Glu Ile Phe Ser Gly Leu 435 440 445

His Asn Leu Gln Tyr Leu Tyr Leu Glu Tyr Asn Leu Ile Lys Glu Ile 450 455 460

Ser Ala Gly Thr Phe Asp Ser Met Pro Asn Leu Gln Leu Leu Tyr Leu 465 470 475 480

Asn Asn Asn Leu Leu Lys Ser Leu Pro Val Tyr Ile Phe Ser Gly Ala 485 490 495

Pro Leu Ala Arg Leu Asn Leu Arg Asn Asn Lys Phe Met Tyr Leu Pro $50\dot{0}$ 505 510

Val Ser Gly Val Leu Asp Gln Leu Gln Ser Leu Thr Gln Ile Asp Leu 515 520 525

Glu Gly Asn Pro Trp Asp Cys Thr Cys Asp Leu Val Ala Leu Lys Leu 530 535 540

Trp Val Glu Lys Leu Ser Asp Gly Ile Val Val Lys Glu Leu Lys Cys 545 550 555 560

Glu Thr Pro Val Gln Phe Ala Asn Ile Glu Leu Lys Ser Leu Lys Asn 565 570 575

Glu Ile Leu Cys Pro Lys Leu Leu Asn Lys Pro Ser Ala Pro Phe Thr 580 585 590

Ser Pro Ala Pro Ala Ile Thr Phe Thr Thr Pro Leu Gly Pro Ile Arg 595 600 605

Ser Pro Pro Gly Gly Pro Val Pro Leu Ser Ile Leu Ile Leu Ser Ile 610 615 620

Leu Val Val Leu Ile Leu Thr Val Phe Val Ala Phe Cys Leu Leu Val 625 630 635

Phe Val Leu Arg Arg Asn Lys Lys Pro Thr Val Lys His Glu Gly Leu 645 650 655

Gly Asn Pro Asp Cys Gly Ser Met Gln Leu Gln Leu Arg Lys His Asp 660 665 670

His Lys Thr Asn Lys Lys Asp Gly Leu Ser Thr Glu Ala Phe Ile Pro 675 680 685

Gln Thr Ile Glu Gln Met Ser Lys Ser His Thr Cys Gly Leu Lys Glu 690 695 700

Ser Glu Thr Gly Phe Met Phe Ser Asp Pro Pro Gly Gln Lys Val Val 705 710 715 720

Met Arg Asn Val Ala Asp Lys Glu Lys Asp Leu Leu His Val Asp Thr 725 730 735

Arg Lys Arg Leu Ser Thr Ile Asp Glu Leu Asp Glu Leu Phe Pro Ser 740 745 750

PCT/US01/28013

Arg Asp Ser Asn Val Phe Ile Gln Asn Phe Leu Glu Ser Lys Glu 755 760 765

Tyr Asn Ser Ile Gly Val Ser Gly Phe Glu Ile Arg Tyr Pro Glu Lys
770 . 775 780

Gln Pro Asp Lys Lys Ser Lys Lys Ser Leu Ile Gly Gly Asn His Ser 785 790 795 800

Lys Ile Val Val Glu Gln Arg Lys Ser Glu Tyr Phe Glu Leu Lys Ala 805 810 815

Lys Leu Gln Ser Ser Pro Asp Tyr Leu Gln Val Leu Glu Glu Gln Thr 820 825 830

Ala Leu Asn Lys Ile 835

<210> 26

WO 02/20569

<211> 1694

<212> DNA

<213> Homo sapiens

<400> 26 tcactctatg aacagcacat ggtgagcccc atggttcatg tctatagaag tccatccttt 60 120 catctccaaa gaagtctttt ggaacaggaa aatcattcac cactcacagg gtcaaatatg 180 aaatacaaaa ccacgaacca atcaacagaa tttttatcct tccaagatgc cagctcattg 240 tacagaaaca ttttagaaaa agaaagggaa cttcagcaac tgggaatcac agaataccta 300 360 aggaaaaaca ttgctcagct ccagcctgat atggaggcac attatcctgg agcccacgaa 420 gagctgaagt taatggaaac attaatgtac tcacgtccaa ggaaggtatt agtggaacag acaaaaaatg agtattttga acttaaagct aatttacatg ctgaacctga ctatttagaa 480 gtcctggagc agcaaacata gatggagagt ttgagggctt tcgcagaaat gctgtgattc 540 tgttttaagt ccataccttg taaataagtg ccttacgtga gtgtgtcatc aatcagaacc 600 660 taagcacagc agtaaactat ggggaaaaaa aaagaagaag aaaagaaact cagggatcac

tgggagaagc	catggcatta	tcttcaggca	atttagtctg	tcccaaataa	aataaatcct	720
tgcatgtaaa	tcattcaagg	gttatagtaa	tatttcatat	actgaaaagt	gtctcatagg	780
agtcctcttg	cacatctaaa	aaggctgaac	atttaagtat	cccgcaattt	tcttgaattg	840
ctttccctat	agattaaṭta	caattggatt	tcatcattta	aaaaccatac	ttgtatatgt	.900
agttataata	tgtaaggaat	acattgttta	taaccagtat	gtacttcaaa	aatgtgtatt	960
gtcaaacata	cctaactttc	ttgcaataaa	tgcaaaagaa	actggaactt	gacaattata	1020
aatagtaata	gtgaagaaaa	aatagaaagg	ttgcaattat	ataggccatg	ggtggctcaa	1080
aactttgaac	atttgagctt	aaacaaatgc	cactctcatg	cattctaaat	taaaaagtta	1140
aaatgattaa	tagttcaggt	ggaagaaata	agcatacttt	ttgggttttc	tacacatttt	1200
gtgtagacaa	ttttaatgtc	agtgctgctg	tgaactaaag	tatgtcattt	atgctcaaag	1260
tttaattctt	cttcttggga	tattttaaaa	atgctactga	gattctgctg	taaatatgac	1320
tagagaatat	attgggtttg	ctttatttca	taggcttaat	tctttgtaaa	tctgaatgac	1380
cataatagaa	atacatttct	tgtggcaagt	aattcacagt	tgtaaagtaa	ataggaaaaa	1440
ttattttatt	tttattgatg	tacattgata	gatgccataa	atcagtagca	aaaggcactt	1500
ctaaaggtaa	gtggtttaag	ttgcctcaag	agagggacaa	tgtagcttta	ttttacaaga	1560
aggcatagtt	"agatttctat	gaaatattta	ttctgtacag	ttttatatag	ttttggttca	1620
caaaagtaat	tattcttggg	tgcctttcaa	gaaaattaaa	aatactactc	actacaataa	1680
aactaaaatq	222					1694

<210> 27

<211> 841

<212> PRT

<213> Homo sapiens

<400> 27

Met Lys Leu Trp Ile His Leu Phe Tyr Ser Ser Leu Leu Ala Cys Ile 1 10 15

Ser Leu His Ser Gln Thr Pro Val Leu Ser Ser Arg Gly Ser Cys Asp 20 25 30

Ser Leu Cys Asn Cys Glu Glu Lys Asp Gly Thr Met Leu Ile Asn Cys 35 40

WO 02/20569

Glu Ala Lys Gly Ile Lys Met Val Ser Glu Ile Ser Val Pro Pro Ser 50 55 60

Arg Pro Phe Gln Leu Ser Leu Leu Asn Asn Gly Leu Thr Met Leu His 65 70 75 80

Thr Asn Asp Phe Ser Gly Leu Thr Asn Ala Ile Ser Ile His Leu Gly 85 90 95

Phe Asn Asn Ile Ala Asp Ile Glu Ile Gly Ala Phe Asn Gly Leu Gly
100 105 110

Leu Leu Lys Gln Leu His Ile Asn His Asn Ser Leu Glu Ile Leu Lys 115 120 125

Glu Asp Thr Phe His Gly Leu Glu Asn Leu Glu Phe Leu Gln Ala Asp 130 135 140

Asn Asn Phe Ile Thr Val Ile Glu Pro Ser Ala Phe Ser Lys Leu Asn 145 150 155 160

Arg Leu Lys Val Leu Ile Leu Asn Asp Asn Ala Ile Glu Ser Leu Pro 165 170 175

Pro Asn Ile Phe Arg Phe Val Pro Leu Thr His Leu Asp Leu Arg Gly 180 185 190

Asn Gln Leu Gln Thr Leu Pro Tyr Val Gly Phe Leu Glu His Ile Gly 195 200 205

Arg Ile Leu Asp Leu Gln Leu Glu Asp Asn Lys Trp Ala Cys Asn Cys 210 215 220

Asp Leu Leu Gln Leu Lys Thr Trp Leu Glu Asn Met Pro Pro Gln Ser 225 230 . 235 240

Ile Ile Gly Asp Val Val Cys Asn Ser Pro Pro Phe Phe Lys Gly Ser 245 250 255

Ile Leu Ser Arg Leu Lys Lys Glu Ser Ile Cys Pro Thr Pro Pro Val 260 265 270

Tyr Glu Glu His Glu Asp Pro Ser Gly Ser Leu His Leu Ala Ala Thr

 Ser
 Ser
 Ile
 Asn
 Asp
 Ser
 Arg
 Met
 Ser
 Thr
 Lys
 Thr
 Thr
 Ser
 Ile
 Leu

 Lys
 Leu
 Pro
 Thr
 Lys
 Ala
 Pro
 Gly
 Leu
 Ile
 Pro
 Tyr
 Ile
 Thr
 Lys
 Pro
 Pro
 320

 Ser
 Thr
 Gln
 Leu
 Pro
 Gly
 Pro
 Tyr
 Cys
 Pro
 P

Ile Leu Ala Gly Asn Ile Ile His Ser Leu Met Lys Ser Asp Leu Val 370 375 380

360

Glu Tyr Phe Thr Leu Glu Met Leu His Leu Gly Asn Asn Arg Ile Glu 385 390 395 400

Val Leu Glu Glu Gly Ser Phe Met Asn Leu Thr Arg Leu Gln Lys Leu 405 410 415

Tyr Leu Asn Gly Asn His Leu Thr Lys Leu Ser Lys Gly Met Phe Leu 420 425 430

Gly Leu His Asn Leu Glu Tyr Leu Tyr Leu Glu Tyr Asn Ala Ile Lys 435 440 445

Glu Ile Leu Pro Gly Thr Phe Asn Pro Met Pro Lys Leu Lys Val Leu 450 455 460

Tyr Leu Asn Asn Leu Leu Gln Val Leu Pro Pro His Ile Phe Ser

Gly Val Pro Leu Thr Lys Val Asn Leu Lys Thr Asn Gln Phe Thr His 485 490 495

Leu Pro Val Ser Asn Ile Leu Asp Asp Leu Asp Leu Leu Thr Gln Ile 500 505 510 Asp Leu Glu Asp Asn Pro Trp Asp Cys Ser Cys Asp Leu Val Gly Leu 515 520 525

Gln Gln Trp Ile Gln Lys Leu Ser Lys Asn Thr Val Thr Asp Asp Ile 530 540

Leu Cys Thr Ser Pro Gly His Leu Asp Lys Lys Glu Leu Lys Ala Leu 545 550 555 560

Asn Ser Glu Ile Leu Cys Pro Gly Leu Val Asn Asn Pro Ser Met Pro 565 570 575

Thr Gln Thr Ser Tyr Leu Met Val Thr Thr Pro Ala Thr Thr Asn 580 585 590

Thr Ala Asp Thr Ile Leu Arg Ser Leu Thr Asp Ala Val Pro Leu Ser 595 600 605

Val Leu Ile Leu Gly Leu Leu Ile Met Phe Ile Thr Ile Val Phe Cys 610 615 620

Ala Ala Gly Ile Val Val Leu Val Leu His Arg Arg Arg Tyr Lys 625 630 635

Lys Lys Gln Val Asp Glu Gln Met Arg Asp Asn Ser Pro Val His Leu 645 650 655

Gln Tyr Ser Met Tyr Gly His Lys Thr Thr His His Thr Thr Glu Arg 660 670

Pro Ser Ala Ser Leu Tyr Glu Gln His Met Val Ser Pro Met Val His 675 680 685

Val Tyr Arg Ser Pro Ser Phe Gly Pro Lys His Leu Glu Glu Glu 690 695 700

Glu Arg Asn Glu Lys Glu Gly Ser Asp Ala Lys His Leu Gln Arg Ser 705 710 715 720

Leu Leu Glu Gln Glu Asn His Ser Pro Leu Thr Gly Ser Asn Met Lys
725 730 735

Tyr Lys Thr Thr Asn Gln Ser Thr Glu Phe Leu Ser Phe Gln Asp Ala 740 745 750

Ser Ser Leu Tyr Arg Asn Ile Leu Glu Lys Glu Arg Glu Leu Gln Gln 755 760 765
Leu Gly Ile Thr Glu Tyr Leu Arg Lys Asn Ile Ala Gln Leu Gln Pro 770 780
Asp Met Glu Ala His Tyr Pro Gly Ala His Glu Glu Leu Lys Leu Met 785 790 795 800
Glu Thr Leu Met Tyr Ser Arg Pro Arg Lys Val Leu Val Glu Gln Thr 805 810 815
Lys Asn Glu Tyr Phe Glu Leu Lys Ala Asn Leu His Ala Glu Pro Asp 820 825 830
Tyr Leu Glu Val Leu Glu Gln Gln Thr 835 840
<210> 28
<211> 639
<212> DNA
<213> Homo sapiens
<220>·
<221> CDS
<222> (1)(636)
<223>
<400> 28
atg gtt tta ccc tca tat tca aaa tca gag gga ggg tca tta ttg gat 48 Met Val Leu Pro Ser Tyr Ser Lys Ser Glu Gly Gly Ser Leu Leu Asp 1 5 10 15
atc tac tgt tta ctc acg tat tgg atg gag gtg gtg ccc acc ctc ttg Ile Tyr Cys Leu Leu Thr Tyr Trp Met Glu Val Val Pro Thr Leu Leu 20 25 30
gca gag aca aag att cca gcc act gat gtc gct gat gcc agc ctg aat 144 Ala Glu Thr Lys Ile Pro Ala Thr Asp Val Ala Asp Ala Ser Leu Asn 35 40 45

gaa tgt tcc agt acc gaa agg aaa caa gac gta gtg ttg ctg ttc gtg 192

Glu	Cys 50	Ser	Ser	Thr	Glu	Arg 55	Lys	Gln	Asp	Val	Val 60	Leu	Leu	Phe	Val		
acc Thr 65	ttg Leu	tcc Ser	cac His	aca Thr	cag Gln 70	cca Pro	cct Pro	ctg Leu	ttt Phe	cac His 75	ctg Leu	cct Pro	tat Tyr	gtc Val	cag Gln 80	2	240
aaa Lys	ccc Pro	tta Leu	atc Ile	tct Ser 85	aat Asn	gtg Val	gag Glu	cag Gln	ctg Leu 90	atc Ile	ctg Leu	ggg Gly	atc Ile	ccg Pro 95	Gly	2	288
cag Gln	aat Asn	cgc Arg	cgg Arg 100	gag Glu	ata Ile	ggc Gly	cat His	ggc Gly 105	cag Gln	gat Asp	atc Ile	ttt Phe	cca Pro 110	gca Ala	gag Glu	3	336
aag Lys	ctc Leu	tgc Cys 115	cat His	ctg Leu	cag Gln	gat Asp	cgc Arg 120	aag Lys	gtg Val	aac Asn	ctt Leu	cac His 125	aga Arg	gct Ala	gcc Ala		384
tgg Trp	ggc Gly 130	gag Glu	tgt Cys	att Ile	gtt Val	gca Ala 135	ccc Pro	aag Lys	act Thr	ctc Leu	agc Ser 140	ttc Phe	tct Ser	tac Tyr	tgt Cys	4	432
cag Gln 145	Gly ggg	acc Thr	tgc Cys	ccg Pro	gcc Ala 150	ctc Leu	aac Asn	agt Ser	gag Glu	ctc Leu 155	cgt Arg	cat His	tcc Ser	agc Ser	ttt Phe 160	•	480
gag Glu	tgc Cys	tat Tyr	aag Lys	agg Arg 165	gca Ala	gta Val	cct Pro	acc Thr	tgt Cys 170	ccc Pro	tgg Trp	ctc Leu	ttc Phe	cag Gln 175	acc Thr	!	528
tgc Cys	cgt Arg	ccc Pro	acc Thr 180	atg Met	gtc Val	aga Arg	ctc Leu	ttc Phe 185	tcc Ser	ctg Leu	atg Met	gtc Val	cag Gln 190	gat Asp	gac Asp	!	576
gaa Glu	cac His	aag Lys 195	atg Met	agt Ser	gtg Val	cac His	tat Tyr 200	gtg Val	aac Asn	act Thr	tcc Ser	ttg Leu 205	gtg Val	gag Glu	aag Lys		624
_		tgc Cys		tga													639
<21	0 >	29															
<21	1>	212															
<21	2>	PRT											•				
<21	3>	Homo	sap	iens													

<400> 29

Met Val Leu Pro Ser Tyr Ser Lys Ser Glu Gly Gly Ser Leu Leu Asp 1 5 10 15

64

Ile Tyr Cys Leu Leu Thr Tyr Trp Met Glu Val Val Pro Thr Leu Leu 20 25 30

Ala Glu Thr Lys Ile Pro Ala Thr Asp Val Ala Asp Ala Ser Leu Asn 35 40 45

Glu Cys Ser Ser Thr Glu Arg Lys Gln Asp Val Val Leu Leu Phe Val 50 55 60

Thr Leu Ser His Thr Gln Pro Pro Leu Phe His Leu Pro Tyr Val Gln 65 70 75 80

Lys Pro Leu Ile Ser Asn Val Glu Gln Leu Ile Leu Gly Ile Pro Gly 85 90 95

Gln Asn Arg Arg Glu Ile Gly His Gly Gln Asp Ile Phe Pro Ala Glu 100 105 110

Lys Leu Cys His Leu Gln Asp Arg Lys Val Asn Leu His Arg Ala Ala 115 120 125

Trp Gly Glu Cys Ile Val Ala Pro Lys Thr Leu Ser Phe Ser Tyr Cys 130 135 140

Gln Gly Thr Cys Pro Ala Leu Asn Ser Glu Leu Arg His Ser Ser Phe 145 150 155 160

Glu Cys Tyr Lys Arg Ala Val Pro Thr Cys Pro Trp Leu Phe Gln Thr 165 170 175

Cys Arg Pro Thr Met Val Arg Leu Phe Ser Leu Met Val Gln Asp Asp 180 185 190

Glu His Lys Met Ser Val His Tyr Val Asn Thr Ser Leu Val Glu Lys 195 200 205

Cys Gly Cys Ser 210

<210> 30

<211> 1061

<212> DNA

65

<213> Homo sapiens

<220>
<221> CDS
<222> (204)..(860)

<223>

<400> 30

tggccaggca gaggtctgtg gagtggagag gcgaggcctc acggtggaac tctcagatga cagcatgcag gcaccaagag agtggacgca catacagaag acagccatgc actgagctgg 120 180 ggacatgcaa caataacagg tgagttccaa caaattggtt caaaaagagg ggggataaac acgetggece atgetgggea age atg gea eea eet tee agg eac tgt ett ett 233 Met Ala Pro Pro Ser Arg His Cys Leu Leu ctg atc agc act ctg ggt gtc ttt gca ctt aac tgc ttc acc aaa ggt 281 Leu Ile Ser Thr Leu Gly Val Phe Ala Leu Asn Cys Phe Thr Lys Gly cag aag aac agc acg ctc atc ttc aca agg gaa aac acc att cgg aac 329 Gln Lys Asn Ser Thr Leu Ile Phe Thr Arg Glu Asn Thr Ile Arg Asn 35 377 tgc agc tgt tct gcg gac atc cgg gat tgt gac tac agt ttg gcc aac Cys Ser Cys Ser Ala Asp Ile Arg Asp Cys Asp Tyr Ser Leu Ala Asn 50 425 ctg atg tgc aac tgt aaa acc gtc ctg ccc ctt gca gta gag cga acc Leu Met Cys Asn Cys Lys Thr Val Leu Pro Leu Ala Val Glu Arg Thr 473 age tac aat gge cat etg ace ate tgg tte acg gae aca tet geg etg Ser Tyr Asn Gly His Leu Thr Ile Trp Phe Thr Asp Thr Ser Ala Leu 521 ggc cac ctg ctg aac ttc acg ctg gtc caa gac ctg aag ctt tcc ctg Gly His Leu Leu Asn Phe Thr Leu Val Gln Asp Leu Lys Leu Ser Leu 95 100 tgc agc acc aac act ctc ccc act gaa tac ctg gct att tgt ggt ctg 569 Cys Ser Thr Asn Thr Leu Pro Thr Glu Tyr Leu Ala Ile Cys Gly Leu 115 617 aag agg ctg cgc atc aac atg gag gcc aag cat ccc ttc cca gag cag Lys Arg Leu Arg Ile Asn Met Glu Ala Lys His Pro Phe Pro Glu Gln age tta etc atc cat age ggt ggg gae agt gae tec aga gag aag ecc 665 Ser Leu Leu Ile His Ser Gly Gly Asp Ser Asp Ser Arg Glu Lys Pro

	140					145					150		•			
					ggc Gly 160											713
					aac Asn											761
					att Ile											809
aga Arg	acc Thr	ttc Phe 205	cca Pro	atg Met	cca Pro	agc Ser	aac Asn 210	aaa Lys	agc Ser	tat Tyr	gtt Val	gtc Val 215	aca Thr	ttt Phe	att Ile	857
tac Tyr	tago	cataa	ıta a	ictgt	gtco	ca go	etgco	etgga	a act	ttgg	gcaa	atga	itgaa	ita		910
attt	gcag	gaa g	ggaat	ctg	ga aa	ataaç	ggccg	g tga	agata	aggt	atco	ctac	cc a	caa	etgtgc	970
ctct	ctc	ege a	agget	ccat	t to	gcaad	cacaç	g cca	acaca	atac	caat	aaco	ag o	tct	ctgttc	1030
tgct	ctgt	ige (ccaac	tgcg	ga ga	acad	ettt	g								1061
<210)> 3	31														
<211	L> 2	219														
<212	2> I	PRT														
<213	3> I	Tomo	sapi	iens												
<400)> 3	31														
Met 1	Ala	Pro	Pro	Ser 5	Arg	His	Cys	Leu	Leu 10	Leu	Ile	Ser	Thr	Leu 15	Gly	
Val	Phe	Ala	Leu 20	Asn	Cys	Phe	Thr	L уs 25	Gly	Gln	Lys	Asn	Ser 30	Thr	Leu	
Ile	Phe	Thr 35	Arg	Glu	Asn	Thr	Ile 40	Arg	Asn	Cys	Ser	Cys 45	Ser	Ala	Asp	
Ile	Arg 50	Asp	Cys	Asp	Tyr	Ser 55	Leu	Ala	Asn	Leu	Met 60	Cys	Asn	Cys	Lys	
Thr 65	Val	Leu	Pro	Leu	Ala 70	Val	Glu	Arg	Thr	Ser 75	Tyr	Asn	Gly	His	Leu 80	

67

Thr Ile Trp Phe Thr Asp Thr Ser Ala Leu Gly His Leu Leu Asn Phe
85 90 95

Thr Leu Val Gln Asp Leu Lys Leu Ser Leu Cys Ser Thr Asn Thr Leu 100 · 105 110

Pro Thr Glu Tyr Leu Ala Ile Cys Gly Leu Lys Arg Leu Arg Ile Asn 115 120 125

Met Glu Ala Lys His Pro Phe Pro Glu Gln Ser Leu Leu Ile His Ser 130 135 140

Gly Gly Asp Ser Asp Ser Arg Glu Lys Pro Met Trp Leu His Lys Gly 145 150 155

Trp Gln Pro Cys Met Tyr Ile Ser Phe Leu Asp Met Ala Leu Phe Asn 165 170 175

Arg Asp Ser Ala Leu Lys Ser Tyr Ser Ile Glu Asn Val Thr Ser Ile 180 185 190

Ala Asn Asn Phe Pro Asp Phe Ser Tyr Phe Arg Thr Phe Pro Met Pro 195 200 205

Ser Asn Lys Ser Tyr Val Val Thr Phe Ile Tyr 210 215

<210> 32

<211> 921

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (255)..(890)

<223>

<400> 32

acca	gtgg	tg a	cctc	atga	t ct	cctc	gtca	gtt	ctgc	ctg	tgaa	gggt	.cc c	acca	tctct	60
aaca	tcac	ca c	actg	gagc	c to	agct	tctg	aga	cagg	aac	tctt	acag	at c	gagcc	acaga	120
ctag	agca	.cg t	ttat	gcgc	a cc	acgg	gago	aca	tgct	atc	agtg	ctgg	cg c	gagag	tttgg	180
gggt	aagg	ag g	tgac	ctac	a at	ggac	tggc	tca	tgag	gga	gaaa	cagg	aa o	cacac	cagtc	240
catg	ctgg	jac a	aga	atg Met 1	aca Thr	tca Ser	cct Pro	tcc Ser 5	agc Ser	ttc Phe	tgc Cys	ctc Leu	ctt Leu 10	ctg Leu	ctc Leu	290
caa Gln	gcg Ala	cta Leu 15	ggc Gly	atc Ile	gtt Val	gcc Ala	ctt Leu 20	ggc	cac His	ttc Phe	aca Thr	aaa Lys 25	gct Ala	cag Gln	aac Asn	338
aac Asn	aca Thr 30	ctg Leu	att Ile	ttc Phe	aca Thr	aaa Lys 35	gga Gly	aat Asn	acc Thr	att Ile	cgc Arg 40	aac Asn	tgc Cys	agc Ser	tgc Cys	386
cca Pro 45	gta Val	gac Asp	atc Ile	agg Arg	gac Asp 50	tgt Cys	gac Asp	tac Tyr	agt Ser	ttg Leu 55	gct Ala	aac Asn	ttg Leu	ata Ile	tgc Cys 60	434
agc Ser	tgt Cys	aag Lys	tct Ser	atc Ile 65	ctg Leu	cct Pro	tct Ser	gcc Ala	atg Met 70	gag Glu	caa Gln	acc Thr	agc Ser	tat Tyr 75	cat His	482
ggc Gly	cat His	ctg Leu	acc Thr 80	atc Ile	tgg Trp	ttc Phe	aca Thr	gat Asp 85	ata Ile	tcc Ser	aca Thr	ttg Leu	ggc Gly 90	cac His	gtg Val	530
ctg Leu	aag Lys	ttc Phe 95	act Thr	ctg Leu	gtc Val	caa Gln	gac Asp 100	ttg Leu	aag Lys	ctt Leu	tcc Ser	cta Leu 105	tgt Cys	ggt Gly	tcc Ser	578
agc Ser	acc Thr 110	ttc Phe	ccc Pro	acc Thr	aag Lys	tac Tyr 115	ctg Leu	gct Ala	atc Ile	tgt Cys	999 Gly 120	ctg Leu	cag Gln	agg Arg	ctt Leu	626
cgc Arg 125	atc Ile	cat His	act Thr	aag Lys	gcc Ala 130	agg Arg	cat His	ccc Pro	tcc Ser	cgg Arg 135	gjå aaa	cag Gln	agt Ser	ttg Leu	ctc Leu 140	674
atc Ile	cac His	agc Ser	aga Arg	agg Arg 145	gaa Glu	ggc	agt Ser	tcc Ser	ttg Leu 150	tac Tyr	aaa Lys	ggc Gly	tgg Trp	caa Gln 155	aca Thr	722
tgt Cys	atg Met	ttc Phe	atc Ile 160	tca Ser	ttc Phe	tta Leu	gat Asp	gtg Val 165	gct Ala	ctt Leu	ttc Phe	aac Asn	999 Gly 170	gac Asp	tca Ser	770
tct Ser	tta Leu	aag Lys 175	tca Ser	tac Tyr	agt Ser	att Ile	gac Asp 180	Asn	att Ile	tct Ser	agc Ser	ctc Leu 185	Ala	agt Ser	gac Asp	818
ttt Phe	cct Pro 190	Asp	ttt Phe	tct Ser	tac Tyr	ttt Phe 195	Lys	acg Thr	tcc Ser	cca Pro	atg Met 200	Pro	ago Ser	aac Asn	aga Arg	866

920

921

age tat gtt gtc aca gtt att tac tagcatectg tgtccctcca ccaggaactc Ser Tyr Val Val Thr Val Ile Tyr 210 t <210> 33 <211> 212 <212> PRT <213> Mus musculus <400> 33 Met Thr Ser Pro Ser Ser Phe Cys Leu Leu Leu Gln Ala Leu Gly Ile Val Ala Leu Gly His Phe Thr Lys Ala Gln Asn Asn Thr Leu Ile 25 Phe Thr Lys Gly Asn Thr Ile Arg Asn Cys Ser Cys Pro Val Asp Ile 35 40 Arg Asp Cys Asp Tyr Ser Leu Ala Asn Leu Ile Cys Ser Cys Lys Ser Ile Leu Pro Ser Ala Met Glu Gln Thr Ser Tyr His Gly His Leu Thr Ile Trp Phe Thr Asp Ile Ser Thr Leu Gly His Val Leu Lys Phe Thr Leu Val Gln Asp Leu Lys Leu Ser Leu Cys Gly Ser Ser Thr Phe Pro 105 Thr Lys Tyr Leu Ala Ile Cys Gly Leu Gln Arg Leu Arg Ile His Thr Lys Ala Arg His Pro Ser Arg Gly Gln Ser Leu Leu Ile His Ser Arg Arg Glu Gly Ser Ser Leu Tyr Lys Gly Trp Gln Thr Cys Met Phe Ile 155

70

Ser Phe Leu Asp Val Ala Leu Phe Asn Gly Asp Ser Ser Leu Lys Ser 165 170 175

Tyr Ser Ile Asp Asn Ile Ser Ser Leu Ala Ser Asp Phe Pro Asp Phe
180 185 190

Ser Tyr Phe Lys Thr Ser Pro Met Pro Ser Asn Arg Ser Tyr Val Val 195 200 205

Thr Val Ile Tyr 210

<210> 34

<211> 693

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(690)

<223>

		caa ctt gtg Gln Leu Val			
		gtt gcc atg Val Ala Met 25	_		
	r Val Gly Ala	agc att gtg Ser Ile Val			
		gcc aca cac Ala Thr His 55			
gac atc ta Asp Ile Ty 65	t agc acc ctt r Ser Thr Leu 70	ctg ggc ctg Leu Gly Leu	ccc gct gac Pro Ala Asp 75	atc cag ggt Ile Gln Gly	gcc 240 Ala 80

cag gcc atg atg gtg aca tcc agt gca atc tcc tcc ctg gcc tgc att 288

Gln	Ala	Met	Met	Val 85	Thr	Ser	Ser	Ala	Ile 90	Ser	Ser	Leu	Ala	Cys 95	Ile	
												cag Gln				336
gcc Ala	aaa Lys	gac Asp 115	aga Arg	gtg Val	gcg Ala	gta Val	gca Ala 120	ggt Gly	gga Gly	gtc Val	ttt Phe	ttc Phe 125	atc Ile	ctt Leu	gga Gly	384
												cat His				432
cgg Arg 145	gac Asp	ttc Phe	tac Tyr	tca Ser	cca Pro 150	ctg Leu	gtg Val	cct Pro	gac Asp	agc Ser 155	atg Met	aaa Lys	ttt Phe	gag Glu	att Ile 160	480
gga Gly	gag Glu	gct Ala	ctt Leu	tac Tyr 165	ttg Leu	ggc	att Ile	att Ile	tct Ser 170	tcc Ser	ctg Leu	ttc Phe	tcc Ser	ctg Leu 175	ata Ile	528
gct Ala	gga Gly	atc Ile	atc Ile 180	ctc Leu	tgc Cys	ttt Phe	tcc Ser	tgc Cys 185	tca Ser	tcc Ser	cag Gln	aga Arg	aat Asn 190	cgc Arg	tcc Ser	576
aac Asn	tac Tyr	tac Tyr 195	gat Asp	gcc Ala	tac Tyr	caa Gln	gcc Ala 200	caa Gln	cct Pro	ctt Leu	gcc Ala	aca Thr 205	agg Arg	agc Ser	tct Ser	624
cca Pro	agg Arg 210	gct Ala	ggt Gly	caa Gln	cct Pro	ccc Pro 215	aaa Lys	gtc Val	aag Lys	agt Ser	gag Glu 220	ttc Phe	aat Asn	tcc Ser	tac Tyr	672
			Gly 999			tga										693
<21	0> :	35														
<21	1> :	230														
<21	2> :	PRT														
<21	3> 1	Homo	sap	iens				_								
<40	0 >	35														
Met 1	Ala	Ser	Leu	Gly 5	Leu	Gln	Leu	Val	Gly 10	Tyr	Ile	Leu	Gly	Leu 15	Leu	
Gly	Leu	Leu	Gly 20	Thr	Leu	Val	Ala	Met 25	Leu	Leu	Pro	Ser	Trp	Lys	Thr	

WO 02/20569

72

Ser Ser Tyr Val Gly Ala Ser Ile Val Thr Ala Val Gly Phe Ser Lys 35 40 45

Gly Leu Trp Met Glu Cys Ala Thr His Ser Thr Gly Ile Thr Gln Cys
50 . 55 60

Asp Ile Tyr Ser Thr Leu Leu Gly Leu Pro Ala Asp Ile Gln Gly Ala 65 70 75 80

Gln Ala Met Met Val Thr Ser Ser Ala Ile Ser Ser Leu Ala Cys Ile 85 90 95

Ile Ser Val Val Gly Met Arg Cys Thr Val Phe Cys Gln Glu Ser Arg 100 105 110

Ala Lys Asp Arg Val Ala Val Ala Gly Gly Val Phe Phe Ile Leu Gly
115 120 125

Gly Leu Leu Gly Phe Ile Pro Val Ala Trp Asn Leu His Gly Ile Leu 130 135 140

Arg Asp Phe Tyr Ser Pro Leu Val Pro Asp Ser Met Lys Phe Glu Ile 145 150 155 160

Gly Glu Ala Leu Tyr Leu Gly Ile Ile Ser Ser Leu Phe Ser Leu Ile 165 170 175

Ala Gly Ile Ile Leu Cys Phe Ser Cys Ser Ser Gln Arg Asn Arg Ser 180 185 190

Asn Tyr Tyr Asp Ala Tyr Gln Ala Gln Pro Leu Ala Thr Arg Ser Ser 195 200 205

Pro Arg Ala Gly Gln Pro Pro Lys Val Lys Ser Glu Phe Asn Ser Tyr 210 215 220

Ser Leu Thr Gly Tyr Val 225 230

<210> 36

<211> 1002

<212> DNA

73

<213> Homo sapiens

<220>

<221> misc_feature.

<222> (998)..(998)

<223> unknown amino

<400> 36 tgggttccga gttcattact acaggaaaaa ctgttctctt ctgtggcaca gagaaccctg 60 cttcaaagca gaagtagcag ttccggagtc cagctggcta aaactcatcc cagaggataa 120 180 tggcaaccca tgccttagaa atcgctgggc tgtttcttgg tggtgttgga atggtgggca cagtggctgt cactgtcatg cctcagtgga gagtgtcggc cttcattgaa aacaacatcg 240 tggtttttga aaacttctgg gaaggactgt ggatgaattg cgtgaggcag gctaacatca 300 360 ggatgcagtg caaaatctat gattccctgc tggctctttc tccggaccta caggcagcca gaggactgat gtgtgctgct teegtgatgt cettettgge tttcatgatg gecateettg 420 gcatgaaatg caccaggtgc acgggggaca atgagaaggt gaaagctcac attctgctga 480 540 cggctggaat caatctcatc atcacgggca tggtgggggc caaccctgtg aacctggttt 600 ccaatgccat catcagagat ttttttaccc caatagtgaa tgttgcccaa aaacgtgagc ttggagaagc tctctactta ggatggacca cggcactggt gctsattgtt ggaggagctc 660 tgttctgctg cgttttttgy tgcaacgaaa agagcagtag ctacagatac tcgatacctt 720 cccatcgcac aacccaaaaa agttatcaca ccggaaagaa gtcaccgagc gtctactcca 780 gaagtcagta tgtgtagttg tgtatgtttt tttaacttta ctataaagcc atgcaaatga 840 caaaaatcta tattactttc tcaaaatgga ccccaaagaa actttgattt actgttctta 900 960 actgcctaat cttaattaca ggaactgtgc atcagctatt tatgattcta taagctattt 1002 cagcagaatg agatattaaa tccaatgctt tgattgtnct ag

<210> 37

<211> 225

<212> PRT

<213> Homo sapiens

<400> 37

Met Ala Thr His Ala Leu Glu Ile Ala Gly Leu Phe Leu Gly Gly Val 1 5 10 15

Gly Met Val Gly Thr Val Ala Val Thr Val Met Pro Gln Trp Arg Val 20 25 30

Ser Ala Phe Ile Glu Asn Asn Ile Val Val Phe Glu Asn Phe Trp Glu 35 40 45

Gly Leu Trp Met Asn Cys Val Arg Gln Ala Asn Ile Arg Met Gln Cys 50 55 60

Lys Ile Tyr Asp Ser Leu Leu Ala Leu Ser Pro Asp Leu Gln Ala Ala 65 70 75 80

Arg Gly Leu Met Cys Ala Ala Ser Val Met Ser Phe Leu Ala Phe Met 85 90 95

Met Ala Ile Leu Gly Met Lys Cys Thr Arg Cys Thr Gly Asp Asn Glu 100 105 110

Lys Val Lys Ala His Ile Leu Leu Thr Ala Gly Ile Asn Leu Ile Ile 115 120 125

Thr Gly Met Val Gly Ala Asn Pro Val Asn Leu Val Ser Asn Ala Ile 130 135 140

Ile Arg Asp Phe Phe Thr Pro Ile Val Asn Val Ala Gln Lys Arg Glu 145 150 155 160

Leu Gly Glu Ala Leu Tyr Leu Gly Trp Thr Thr Ala Leu Val Leu Ile 165 170 175

Val Gly Gly Ala Leu Phe Cys Cys Val Phe Cys Cys Asn Glu Lys Ser 180 185 190

Ser Ser Tyr Arg Tyr Ser Ile Pro Ser His Arg Thr Thr Gln Lys Ser 195 200 205

Tyr His Thr Gly Lys Lys Ser Pro Ser Val Tyr Ser Arg Ser Gln Tyr 210 215 220

WO 02/20569

75

Val 225 <210> 38 <211> 833 <212> DNA <213> Homo sapiens <220> CDS <221> <222> (159)..(830) <223> <400> 38 60 ccaagttcag tcacagctac tgatttggac taaaacgtta tgggcagcag ccaaggagaa catcatcaaa gactteteta gacteaaaag getteeaegt tetacatett gageatette 120 taccactccg aattgaacca gtcttcaaag taaaggca atg gca ttt tat ccc ttg 176 Met Ala Phe Tyr Pro Leu 5 224 caa att gct ggg ctg gtt ctt ggg ttc ctt ggc atg gtg ggg act ctt Gln Ile Ala Gly Leu Val Leu Gly Phe Leu Gly Met Val Gly Thr Leu 10 gec aca acc ctt ctg cct cag tgg aga gta tca gct ttt gtt ggc agc 272 Ala Thr Thr Leu Leu Pro Gln Trp Arg Val Ser Ala Phe Val Gly Ser 25 aac att att gtc ttt gag agg ctc tgg gaa ggg ctc tgg atg aat tgc 320 Asn Ile Ile Val Phe Glu Arg Leu Trp Glu Gly Leu Trp Met Asn Cys 368 atc cga caa gcc agg gtc cgg ttg caa tgc aag ttc tat agc tcc ttg Ile Arg Gln Ala Arg Val Arg Leu Gln Cys Lys Phe Tyr Ser Ser Leu 60 55 ttg gct ctc ccg cct gcc ctg gaa aca gcc cgg gcc ctc atg tgt gtg 416 Leu Ala Leu Pro Pro Ala Leu Glu Thr Ala Arg Ala Leu Met Cys Val get gtt gct ctc tcc ttg atc gcc ctg ctt att ggc atc tgt ggc atg 464 Ala Val Ala Leu Ser Leu Ile Ala Leu Leu Ile Gly Ile Cys Gly Met 90

aag cag gtc cag tgc aca ggc tct aac gag agg gcc aaa gca tac ctt

PCT/US01/28013

76

									-							
Lys	Gln	Val 105	Gln	Cys	Thr	Gly	Ser 110	Asn	Glu	Arg	Ala	Lys 115	Ala	Tyr	Leu	
													ttc Phe			560
													ttc Phe			608
cca Pro	gcc Ala	atc Ile	cac His	ata Ile 155	ggt Gly	cag Gln	aaa Lys	cga Arg	gag Glu 160	ctg Leu	gga Gly	gca Ala	gca Ala	ctt Leu 165	ttc Phe	656
													ggt Gly 180			704
tgt Cys	gga Gly	ttt Phe 185	tgc Cys	tgc Cys	tgc Cys	aac Asn	aga Arg 190	aag Lys	aag Lys	caa Gln	Gly 999	tac Tyr 195	aga Arg	tat Tyr	cca Pro	752
													aat Asn			800
_		agt Ser	_				_		_	taa						833
<21	0>	39														
<21	1>	224														
<21	2>	PRT														
<21	3>	Homo	sap	iens												
						•										
<40	0>	39														
Met 1	Ala	Phe	Tyr	Pro 5	Leu	Gln	Ile	Ala	Gly 10	Leu	Val	Leu	Gly	Phe 15	Leu	
Gly	Met	Val	Gly 20	Thr	Leu	Ala	Thr	Thr 25	Leu	Leu	Pro	Gln	Trp 30	Arg	Val	
Ser	Ala	Phe 35	Val	Gly	Ser	Asn	Ile	Ile	Val	Phe	Glu	Arg	Leu	Trp	Glu	•

Gly Leu Trp Met Asn Cys Ile Arg Gln Ala Arg Val Arg Leu Gln Cys 50 60

77

Lys Phe Tyr Ser Ser Leu Leu Ala Leu Pro Pro Ala Leu Glu Thr Ala 65 70 75 80

Arg Ala Leu Met Cys Val Ala Val Ala Leu Ser Leu Ile Ala Leu Leu 85 90 95

Ile Gly Ile Cys Gly Met Lys Gln Val Gln Cys Thr Gly Ser Asn Glu 100 $\,$ 105 $\,$ 110 $\,$

Arg Ala Lys Ala Tyr Leu Leu Gly Thr Ser Gly Val Leu Phe Ile Leu 115 120 125

Thr Gly Ile Phe Val Leu Ile Pro Val Ser Trp Thr Ala Asn Ile Ile 130 135 140

Ile Arg Asp Phe Tyr Asn Pro Ala Ile His Ile Gly Gln Lys Arg Glu 145 150 155 160

Leu Gly Ala Ala Leu Phe Leu Gly Trp Ala Ser Ala Ala Val Leu Phe 165 170 175

Ile Gly Gly Leu Leu Cys Gly Phe Cys Cys Cys Asn Arg Lys Lys 180 185

Gln Gly Tyr Arg Tyr Pro Val Pro Gly Tyr Arg Val Pro His Thr Asp 195 200 205

Lys Arg Arg Asn Thr Thr Met Leu Ser Lys Thr Ser Thr Ser Tyr Val 210 215 220

<210> 40

<211> 393

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(390)

<223>

<400	. 4	10														
atg	gcc	gtg Val	act Thr	gcc Ala 5	tgt Cys	cag Gln	ggc Gly	ttg Leu	999 999	ttc Phe	gtg Val	gtt Val	tca Ser	ctg Leu 15	att Ile	48
ejy aaa	att Ile	gcg Ala	ggc Gly 20	atc Ile	att Ile	gct Ala	gcc Ala	acc Thr 25	tgc Cys	atg Met	gcc Ala	cag Gln	tgg Trp 30	agc Ser	acc Thr	96
caa Gln	gac Asp	ttg Leu 35	tac Tyr	aac Asn	aac Asn	ccc Pro	gta Val 40	aca Thr	gct Ala	gtt Val	ttc Phe	aac Asn 45	tac Tyr	cag Gln	gjå aaa	144
ctg Leu	tgg Trp 50	cgc Arg	tcc Ser	tgt Cys	gtc Val	cga Arg 55	gag Glu	agc Ser	tct Ser	ggc Gly	ttc Phe 60	acc Thr	gag Glu	tgc Cys	egg Arg	192
ggc Gly 65	tac Tyr	ttc Phe	acc Thr	ctg Leu	ctg Leu 70	Gly 999	ctg Leu	cca Pro	ggt Gly	aag Lys 75	ggc Gly	cag Gln	gtg Val	tct Ser	ggc Gly 80	240
tgg Trp	ctg Leu	gag Glu	gga Gly	gag Glu 85	att Ile	gga Gly	ggt Gly	gga Gly	gag Glu 90	gaa Glu	act Thr	gca Ala	ggc Gly	tct Ser 95	gtc Val	288
tgg Trp	gca Ala	cca Pro	cga Arg 100	cag Gln	gga Gly	ctg Leu	ctg Leu	999 Gly 105	agg Arg	gag Glu	gaa Glu	ctg Leu	cga Arg 110	ttc Phe	gtg Val	336
ttt Phe	gac Asp	agg Arg 115	ggc Gly	aac Asn	agc Ser	cac His	ctg Leu 120	cac His	cag Gln	ggt Gly	gga Gly	ata Ile 125	gga Gly	gga Gly	cgg Arg	384
_	cct Pro 130	tag														393
<21	0>	41														
<21	1>	130														
<21	2>	PRT														
<21	3 > :	Homo	sap	iens												
<40	0>	41														
Met 1	Ala	Val	Thr	Ala 5	Суз	Gln	Gly	Leu	Gly 10	Phe	Val	Val	Ser	Leu 15	Ile	
Gly	Ile	Ala	Gly 20	Ile	Ile	Ala	Ala	Thr 25	Сув	Met	Ala	Gln	Trp	Ser	Thr	

79

Gln Asp Leu Tyr Asn Asn Pro Val Thr Ala Val Phe Asn Tyr Gln Gly
35 40 45

Leu Trp Arg Ser Cys Val Arg Glu Ser Ser Gly Phe Thr Glu Cys Arg
50 60

Gly Tyr Phe Thr Leu Leu Gly Leu Pro Gly Lys Gly Gln Val Ser Gly 65 70 75 80

Trp Leu Glu Gly Glu Ile Gly Gly Gly Glu Glu Thr Ala Gly Ser Val 85 90 95

Trp Ala Pro Arg Gln Gly Leu Leu Gly Arg Glu Glu Leu Arg Phe Val 100 105 110

Phe Asp Arg Gly Asn Ser His Leu His Gln Gly Gly Ile Gly Gly Arg 115 120 125

Glu Pro 130

<210> 42

<211> 2247

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (742)..(742)

<223> unknown amino

<220>

<221> misc_feature

<222> (747)..(747)

<223> unknown amino

<220>

<221> misc_feature

<222> (793)..(793)

. <223> unknown amino

<220>

<221> misc_feature

<222> (814)..(814)

<223> unknown amino

<220>

<221> misc_feature

<222> (828)..(828)

<223> unknown amino

<220>

<221> misc_feature

<222> (850)..(850)

<223> unknown amino

<220>

<221> misc_feature

<222> (906)..(906)

<223> unknown amino

<220>

<221> CDS

<222> (1)..(2244)

<223>

	_	_													,	
<400 atg Met 1	gag	gca Ala	aat Asn	cag Gln 5	tgc Cys	ccc Pro	ctg Leu	gtt Val	gtg Val 10	gaa Glu	cca Pro	tct Ser	tac Tyr	cca Pro 15	gac Asp	48
ctg Leu	gtc Val	atc Ile	aat Asn 20	gta Val	gga Gly	gaa Glu	gtg Val	act Thr 25	ctt Leu	gga Gly	gaa Glu	gaa Glu	aac Asn 30	aga Arg	aaa Lys	96
aag Lys	ctg Leu	cag Gln 35	aaa Lys	att Ile	cag Gln	aga Arg	gac Asp 40	caa Gln	gag Glu	aag Lys	gag Glu	aga Arg 45	gtt Val	atg Met	cgg Arg	144
gct Ala	gca Ala 50	tgt Cys	gct Ala	tta Leu	tta Leu	aac Asn 55	tca Ser	gga Gly	gga Gly	gga Gly	gtg Val 60	att Ile	cga Arg	atg Met	gcc Ala	192
aag Lys 65	aag Lys	gtt Val	gag Glu	cat His	ccc Pro 70	gtg Val	gag Glu	atg Met	gga Gly	ctg Leu 75	gat Asp	tta Leu	gaa Glu	cag Gln	tct Ser 80	240
ttg Leu	aga Arg	gag Glu	ctt Leu	att Ile 85	cag Gln	tct Ser	tca Ser	gat Asp	ctg Leu 90	cag Gln	gct Ala	ttc Phe	ttt Phe	gag Glu 95	acc Thr	288
aag Lys	caa Gln	caa Gln	gga Gly 100	agg Arg	tgt Cys	ttt Phe	tac Tyr	att Ile 105	ttt Phe	gtt Val	aaa Lys	tct Ser	tgg Trp 110	agc Ser	agt Ser	336
ggc Gly	cct Pro	ttc Phe 115	cct Pro	gaa Glu	gat Asp	cgc Arg	tct Ser 120	gtc Val	aag Lys	ccc Pro	cgc Arg	ctt Leu 125	tgc Cys	agc Ser	ctc Leu	384
agt Ser	tct Ser 130	tca Ser	tta Leu	tac Tyr	cgt Arg	aga Arg 135	tct Ser	gag Glu	acc Thr	tct Ser	gtg Val 140	cgt Arg	tcc Ser	atg Met	gac Asp	432
tca Ser 145	aga Arg	gag Glu	gca Ala	ttc Phe	tgt Cys 150	ttc Phe	ctg Leu	aag Lys	acc Thr	aaa Lys 155	agg Arg	aag Lys	cca Pro	aaa Lys	atc Ile 160	480
ttg Leu	gaa Glu	gaa Glu	gga Gly	cct Pro 165	ttt Phe	cac His	aaa Lys	att Ile	cac His 170	ГЛЯ	ggt Gly	gta Val	tac Tyr	caa Gln 175	gag Glu	528
ctc Leu	cct Pro	aac Asn	tcg Ser 180	Asp	cct Pro	gct Ala	gac Asp	cca Pro 185	Asn	tcg Ser	gat Asp	cct Pro	gct Ala 190	Asp	cta Leu	576
att Ile	ttc Phe	caa Gln 195	Lys	gac Asp	tat Tyr	ctt Leu	gaa Glu 200	Tyr	ggt Gly	gaa Glu	atc	ctg Leu 205	Pro	ttt Phe	cct Pro	624
gag Glu	tct Ser 210	Gln	tta Leu	gta Val	gag Glu	ttt Phe 215	Lys	cag Gln	ttc Phe	tct Ser	aca Thr 220	Lys	cac His	ttc Phe	caa Gln	672

			aaa Lys													720
			ggc Gly													768
			tgt Cys 260													816
aaa Lys	ata Ile	gaa Glu 275	can Thr	gcc Ala	ata Ile	tac Tyr	aaa Lys 280	cta Leu	cct Pro	tgt Cys	ntt Xaa	cat His 285	ttt Phe	tgc Cys	caa Gln	864
ccc Pro	caa Gln 290	cgc Arg	ccg Pro	ata Ile	acc Thr	ttc Phe 295	aca Thr	ctc Leu	aaa Lys	att Ile	gtg Val 300	gat Asp	gtn Val	tta Leu	aaa Lys	912
agg Arg 305	gga Gly	gag Glu	ctc Leu	tat Tyr	ggc Gly 310	tat Tyr	gct Ala	tgc Cys	atg Met	atc Ile 315	aga Arg	gta Val	aat Asn	ccc Pro	ttc Phe 320	960
tgc Cys	tgt Cys	gca Ala	gtg Val	ttc Phe 325	tca Ser	gaa Glu	gct Ala	ccc Pro	aat Asn 330	tca Ser	tgg Trp	ata Ile	gtg Val	gag Glu 335	gac Asp	1008
aag Lys	tac Tyr	gtc Val	tgc Cys 340	agc Ser	ctg Leu	aca Thr	acc Thr	gag Glu 345	aaa Lys	tgg Trp	gta Val	ggc Gly	atg Met 350	atg Met	aca Thr	1056
gac Asp	aca Thr	gat Asp 355	cca Pro	gat Asp	ctt Leu	cta Leu	cag Gln 360	ttg Leu	tct Ser	gaa Glu	gat Asp	ttt Phe 365	gaa Glu	tgt Cys	cag Gln	1104
ctg Leu	agt Ser 370	Leu	tct Ser	agt Ser	gjy aaa	cct Pro 375	ccc Pro	ctt Leu	agc Ser	aga Arg	cca Pro 380	gtg Val	tac Tyr	tcc Ser	aag Lys	1152
aaa Lys 385	Gly	ctg Leu	gaa Glu	cat His	aaa Lys 390	aag Lys	gaa Glu	ctc Leu	cag Gln	caa Gln 395	Leu	tta Leu	ttt Phe	tca Ser	gtc Val 400	1200
cca Pro	cca Pro	gga Gly	tat Tyr	ttg Leu 405	Arg	tat Tyr	act Thr	cca Pro	gag Glu 410	Ser	ctc Leu	tgg Trp	agg Arg	gac Asp 415	Leu	1248
atc Ile	tca Ser	gag Glu	cac His 420	Arg	gga Gly	cta Leu	gag Glu	gag Glu 425	Leu	ata Ile	aat Asn	aag Lys	caa Gln 430	Met	caa Gln	1296
cct Pro	ttc Phe	Phe 435	Arg	gga Gly	att Ile	gtg Val	ato Ile 440	Leu	tct Ser	aga Arg	agc Ser	tgg Trp 445	Ala	gtg Val	gac Asp	1344
ctg Leu	aac Asn 450	Leu	cag Gln	gag Glu	aag Lys	cca Pro 455	Gly	gto Val	ato Ile	tgt Cys	gat Asp 460	Ala	ctg Leu	ctg Leu	ata Ile	1392

_	_		_		ccc Pro 470								-	_	-	:	1440
					tac Tyr												1488
					gjå aaa												1536
					agt Ser												1584
_	_		_	_	gat Asp						_		-				1632
					ctg Leu 550												1680
					agt Ser												1728
					tat Tyr												1776
					cac His												1824
gcc Ala	atg Met 610	aag Lys	atc Ile	atg Met	gag Glu	aag Lys 615	atc Ile	agg Arg	aat Asn	gtg Val	ttt Phe 620	cac His	tgt Cys	gag Glu	gca Ala		1872
					gtt Val 630												1920
agt Ser	gat Asp	aga Arg	aat Asn	atc Ile 645	tgc Cys	cga Arg	gca Ala	gag Glu	acc Thr 650	cgg Arg	aaa Lys	act Thr	ttc Phe	cta Leu 655	aga Arg		1968
					att Ile												2016
ttc Phe	cgt Arg	act Thr 675	gaa Glu	gat Asp	gly ggg	gac Asp	tgg Trp 680	tat Tyr	gly ggg	aag Lys	gca Ala	aaa Lys 685	agc Ser	atc Ile	act Thr		2064
cgg Arg	aga Arg	gca Ala	aag Lys	ggt Gly	ggc Gly	cca Pro	gga Gly	att Ile	ctc Leu	tgg Trp	atc Ile	ttt Phe	ctg Leu	gat Asp	tac Tyr		2112

84

. . .

700 690 695 2160 ttt cag acc agc cac ttg gat tgc agt ggc ctc cct cct ctc tca gac Phe Gln Thr Ser His Leu Asp Cys Ser Gly Leu Pro Pro Leu Ser Asp 710 caa tat cca aga gaa gag ctc acc aga ata gtt cgc aat gca gat cca 2208 Gln Tyr Pro Arg Glu Glu Leu Thr Arg Ile Val Arg Asn Ala Asp Pro 725 2247 ata gcc aag tac tta caa aaa gaa aat gca agt aat tag Ile Ala Lys Tyr Leu Gln Lys Glu Asn Ala Ser Asn <210> 43 <211> 748 <212> PRT <213> Homo sapiens <220> · <221> misc_feature <222> (248)..(248) <223> The 'Xaa' at location 248 stands for Ile, Val, Leu, or Phe. <220> <221> misc_feature <222> (265)..(265) <223> The 'Xaa' at location 265 stands for Ile, Val, Leu, or Phe. <220> <221> misc_feature <222> (272)..(272) <223> The 'Xaa' at location 272 stands for Arg, Gly, or Trp. <220> <221> misc feature <222> (284)..(284) <223> The 'Xaa' at location 284 stands for Ile, Val, Leu, or Phe. <220>

```
<221> misc_feature
  <222> (742)..(742)
  <223> unknown amino
, <220>
  <221> misc_feature
 <222> (747)..(747)
  <223> unknown amino
  <220>
  <221> misc_feature
  <222> (793)..(793)
  <223> unknown amino
  <220>
  <221> misc_feature
  <222> (814)..(814)
  <223> unknown amino
  <220>
  <221> misc_feature
  <222> (828)..(828)
  <223> unknown amino
  <220>
   <221> misc_feature
   <222> (850)..(850)
   <223> unknown amino
  <220>
   <221> misc_feature
   <222> (906)..(906)
   <223> unknown amino
   <400> 43
   Met Glu Ala Asn Gln Cys Pro Leu Val Val Glu Pro Ser Tyr Pro Asp
1 5 10 15
```

- Leu Val Ile Asn Val Gly Glu Val Thr Leu Gly Glu Glu Asn Arg Lys 20 25 30
- Lys Leu Gln Lys Ile Gln Arg Asp Gln Glu Lys Glu Arg Val Met Arg 35 40. 45
- Ala Ala Cys Ala Leu Leu Asn Ser Gly Gly Gly Val Ile Arg Met Ala 50 55 60
- Lys Lys Val Glu His Pro Val Glu Met Gly Leu Asp Leu Glu Gln Ser 65 70 75 80
- Leu Arg Glu Leu Ile Gln Ser Ser Asp Leu Gln Ala Phe Phe Glu Thr 85 90 95
- Lys Gln Gln Gly Arg Cys Phe Tyr Ile Phe Val Lys Ser Trp Ser Ser 100 105 110
- Gly Pro Phe Pro Glu Asp Arg Ser Val Lys Pro Arg Leu Cys Ser Leu 115 120 125
- Ser Ser Ser Leu Tyr Arg Arg Ser Glu Thr Ser Val Arg Ser Met Asp 130 135 140
- Ser Arg Glu Ala Phe Cys Phe Leu Lys Thr Lys Arg Lys Pro Lys Ile 145 150 155 160
- Leu Glu Glu Gly Pro Phe His Lys Ile His Lys Gly Val Tyr Gln Glu 165 170 175
- Leu Pro Asn Ser Asp Pro Ala Asp Pro Asn Ser Asp Pro Ala Asp Leu 180 185 190
- Ile Phe Gln Lys Asp Tyr Leu Glu Tyr Gly Glu Ile Leu Pro Phe Pro 195 · 200 205
- Glu Ser Gln Leu Val Glu Phe Lys Gln Phe Ser Thr Lys His Phe Gln 210 215 220
- Glu Tyr Val Lys Arg Thr Ile Pro Glu Tyr Val Pro Ala Phe Ala Asn 225 230 235 240
- Thr Gly Gly Gly Tyr Leu Phe Xaa Gly Val Asp Asp Lys Ser Arg Glu 245 250 255

WO 02/20569

- Val Leu Gly Cys Ala Lys Glu Asn Xaa Asp Pro Asp Ser Leu Arg Xaa 260 265 270
- . Lys Ile Glu Thr Ala Ile Tyr Lys Leu Pro Cys Xaa His Phe Cys Gln 275 280 285
 - Pro Gln Arg Pro Ile Thr Phe Thr Leu Lys Ile Val Asp Val Leu Lys 290 295 300
 - Arg Gly Glu Leu Tyr Gly Tyr Ala Cys Met Ile Arg Val Asn Pro Phe 305 310 315
 - Cys Cys Ala Val Phe Ser Glu Ala Pro Asn Ser Trp Ile Val Glu Asp 325 330 335
 - Lys Tyr Val Cys Ser Leu Thr Thr Glu Lys Trp Val Gly Met Met Thr 340 345 350
 - Asp Thr Asp Pro Asp Leu Leu Gln Leu Ser Glu Asp Phe Glu Cys Gln 355 360 365
 - Leu Ser Leu Ser Ser Gly Pro Pro Leu Ser Arg Pro Val Tyr Ser Lys 370 375 380
 - Lys Gly Leu Glu His Lys Lys Glu Leu Gln Gln Leu Leu Phe Ser Val 385 390 395 400
 - Pro Pro Gly Tyr Leu Arg Tyr Thr Pro Glu Ser Leu Trp Arg Asp Leu 405 410 415
 - Ile Ser Glu His Arg Gly Leu Glu Glu Leu Ile Asn Lys Gln Met Gln 420 425 430
 - Pro Phe Phe Arg Gly Ile Val Ile Leu Ser Arg Ser Trp Ala Val Asp 435 440 445
 - Leu Asn Leu Gln Glu Lys Pro Gly Val Ile Cys Asp Ala Leu Leu Ile 450 455 460
 - Ala Gln Asn Ser Thr Pro Ile Leu Tyr Thr Ile Leu Arg Glu Gln Asp 465 470 475 480
 - Ala Glu Gly Gln Asp Tyr Cys Thr Arg Thr Ala Phe Thr Leu Lys Gln

88

495 485 490 Lys Leu Val Asn Met Gly Gly Tyr Thr Gly Lys Val Cys Val Arg Ala 505 Lys Val Leu Cys Leu Ser Pro Glu Ser Ser Ala Glu Ala Leu Glu Ala Ala Val Ser Pro Met Asp Tyr Pro Ala Ser Tyr Ser Leu Ala Gly Thr 535 Gln His Met Glu Ala Leu Leu Gln Ser Leu Val Ile Val Leu Leu Gly 555 550 Phe Arg Ser Leu Leu Ser Asp Gln Leu Gly Cys Glu Val Leu Asn Leu Leu Thr Ala Gln Gln Tyr Glu Ile Phe Ser Arg Ser Leu Arg Lys Asn Arg Glu Leu Phe Val His Gly Leu Pro Gly Ser Gly Lys Thr Ile Met 600 605 Ala Met Lys Ile Met Glu Lys Ile Arg Asn Val Phe His Cys Glu Ala 610 615 His Arg Ile Leu Tyr Val Cys Glu Asn Gln Pro Leu Arg Asn Phe Ile 625 630 Ser Asp Arg Asn Ile Cys Arg Ala Glu Thr Arg Lys Thr Phe Leu Arg 650 645 Glu Asn Phe Glu His Ile Gln His Ile Val Ile Asp Glu Ala Gln Asn 660 Phe Arg Thr Glu Asp Gly Asp Trp Tyr Gly Lys Ala Lys Ser Ile Thr 685 · 675 680 Arg Arg Ala Lys Gly Gly Pro Gly Ile Leu Trp Ile Phe Leu Asp Tyr 690 695 Phe Gln Thr Ser His Leu Asp Cys Ser Gly Leu Pro Pro Leu Ser Asp 715 710

Gln Tyr Pro Arg Glu Glu Leu Thr Arg Ile Val Arg Asn Ala Asp Pro
725 730 735

Ile Ala Lys Tyr Leu Gln Lys Glu Asn Ala Ser Asn 740 745

<210> 44

<211> 2676

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(2673)

<223>

<400> 44

atg Met 1	agt Ser	ctt Leu	agg Arg	att Ile 5	gat Asp	gtg Val	gat Asp	aca Thr	aac Asn 10	ttt Phe	cct Pro	gag Glu	tgt Cys	gtt Val 15	gta Val	48
				gtc Val												96
cct Pro	cgc Arg	ctg Leu 35	cgg Arg	gag Glu	aaa Lys	cag Gln	aat Asn 40	gaa Glu	atc Ile	atc Ile	ctg Leu	cga Arg 45	gca Ala	gta Val	tgt Cys	144
				tct Ser												192
				tat Tyr												240
				cat His 85												288
att Ile	ttt Phe	gtg Val	aaa Lys 100	tca Ser	tgg Trp	aac Asn	aca Thr	gag Glu 105	gct Ala	ggt Gly	gtg Val	cca Pro	ctt Leu 110	gct Ala	acc Thr	336
tta	tgc	tcc		ttg											atg	384

Leu Cys Ser Asn Leu Tyr His Arg Glu Arg Thr Ser Thr Asp Val Met

											•					
		115					120				•	125				
										tgc Cys						432
_				_						cca Pro 155		_	_	_		480
										tca Ser						528
										ctc Leu						576
										gac Asp						624
										ttt Phe						672
gga Gly 225	tat Tyr	gta Val	ttt Phe	ttt Phe	ggt Gly 230	gtg Val	cat His	gat Asp	gag Glu	act Thr 235	tgt Cys	caa Gln	gtg Val	att Ile	gga Gly 240	720
										ttg Leu						768
ggc Gly	tgt Cys	att Ile	aag Lys 260	aag Lys	cta Leu	cct Pro	gtc Val	cat His 265	cat His	ttc Phe	tgc Cys	aca Thr	cag Gln 270	agg Arg	cct Pro	816
gag Glu	ata Ile	aaa Lys 275	tat Tyr	gtc Val	ctt Leu	aac Asn	ttc Phe 280	ctt Leu	gaa Glu	gtg Val	cat His	gat Asp 285	aag Lys	gly ggg	gcc Ala	864
ctc Leu	cgt Arg 290	gga Gly	tat Tyr	gtc Val	tgt Cys	gca Ala 295	atc Ile	aag Lys	gtg Val	gag Glu	aaa Lys 300	ttc Phe	tgc Cys	tgt Cys	gcg Ala	912
gtg Val 305	ttt Phe	gcc Ala	aaa Lys	gtg Val	cct Pro 310	agt Ser	tcc Ser	tgg Trp	cag Gln	gtg Val 315	aag Lys	gac Asp	aac Asn	cgt Arg	gtg Val 320	960
aga Arg	caa Gln	ttg Leu	ccc Pro	aca Thr 325	aga Arg	gaa Glu	tgg Trp	act Thr	gct Ala 330	tgg Trp	atg Met	atg Met	gaa Glu	gct Ala 335	gac Asp	1008
cca Pro	gac Asp	ctt Leu	tcc Ser 340	agg Arg	tgt Cys	cct Pro	gag Glu	atg Met 345	gtt Val	ctc Leu	cag Gln	ttg Leu	agt Ser 350	ttg Leu	tca Ser	1056
tct	gcc	acg	ccc	cgc	agc	aag	cct	gtg	tgc	att	cat	aag	aat	tcg	gaa	1104

Ser	Ala	Thr 355	Pro	Arg	Ser	Lys	Pro 360	Val	Cys	Ile	His	Lys 365	Asn	Ser	Glu	
tgt Cys	ctg Leu 370	aaa Lys	gag Glu	cag Gln	cag Gln	aaa Lys 375	cgc Arg	tac Tyr	ttt Phe	cca Pro	gta Val 380	ttt Phe	tca Ser	gac Asp	aga Arg	1152
gtg Val 385	gta Val	tat Tyr	act Thr	cca Pro	gaa Glu 390	agc Ser	ctc Leu	tac Tyr	aag Lys	gaa Glu 395	ctc Leu	ttc Phe	tca Ser	caa Gln	cat His 400	1200
aaa Lys	gga Gly	ctc Leu	aga Arg	gac Asp 405	tta Leu	ata Ile	aat Asn	aca Thr	gaa Glu 410	atg Met	cgc Arg	cct Pro	ttc Phe	tct Ser 415	caa Gln	1248
gga Gly	ata Ile	ttg Leu	att Ile 420	ttt Phe	tct Ser	caa Gln	agc Ser	tgg Trp 425	gct Ala	gtg Val	gat Asp	tta Leu	ggt Gly 430	ctg Leu	caa Gln	1296
gag Glu	aag Lys	cag Gln 435	gga Gly	gtc Val	atc Ile	tgt Cys	gat Asp 440	gct Ala	ctt Leu	cta Leu	att Ile	tcc Ser 445	cag Gln	aac Asn	aac Asn	1344
acc Thr	cct Pro 450	att Ile	ctc Leu	tac Tyr	acc Thr	atc Ile 455	ttc Phe	agc Ser	aag Lys	tgg Trp	gat Asp 460	gcg Ala	Gly aaa	tgc Cys	aag Lys	1392
ggc Gly 465	tat Tyr	tct Ser	atg Met	ata Ile	gtt Val 470	gcc Ala	tat Tyr	tct Ser	ttg Leu	aag Lys 475	cag Gln	aag Lys	ctg Leu	gtg Val	aac Asn 480	1440
aaa Lys	ggc	ggc Gly	tac Tyr	act Thr 485	Gly 999	agg Arg	tta Leu	tgc Cys	atc Ile 490	acc Thr	ccc Pro	ttg Leu	gtc Val	tgt Cys 495	gtg Val	1488
ctg Leu	aat Asn	tct Ser	gat Asp 500	aga Arg	aaa Lys	gca Ala	cag Gln	agc Ser 505	gtt Val	tac Tyr	agt Ser	tcg Ser	tat Tyr 510	tta Leu	caa Gln	1536
att Ile	tac Tyr	cct Pro 515	gaa Glu	tcc Ser	tat Tyr	aac Asn	ttc Phe 520	Met	acc Thr	ccc Pro	cag Gln	cac His 525	atg Met	gaa Glu	gcc Ala	1584
ctg Leu	tta Leu 530	cag Gln	tcc Ser	ctc Leu	gtg Val	ata Ile 535	gtc Val	ttg Leu	ctt Leu	gly aaa	ttc Phe 540	Lys	tcc Ser	ttc Phe	tta Leu	1632
agt Ser 545	gaa Glu	gag Glu	.ctg Leu	ggc Gly	tct Ser 550	Glu	gtt Val	ttg Leu	aac Asn	cta Leu 555	Leu	aca Thr	aat Asn	aaa Lys	cag Gln 560	1680
tat Tyr	gag Glu	ttg Leu	ctt Leu	tca Ser 565	Lys	aac Asn	ctt Leu	cgc Arg	aag Lys 570	Thr	aga Arg	gag Glu	ttg Leu	ttt Phe 575	gtt	1728
cat His	ggc Gly	tta Leu	cct Pro 580	Gly	tca Ser	gly	aag Lys	act Thr 585	Ile	ttg Leu	gct Ala	ctt Leu	agg Arg 590	Ile	atg Met	1776

1	gag Glu	aag Lys	atc Ile 595	agg Arg	aat Asn	gtg Val	ttt Phe	cac His 600	tgt Cys	gaa Glu	ccg Pro	gct Ala	aac Asn 605	att Ile	ctc Leu	tac Tyr	1824
	atc Ile	tgt Cys 610	gaa Glu	aac Asn	cag Gln	ccc Pro	ctg Leu 615	aag Lys	aag Lys	ttg Leu	gtg Val	agt Ser 620	ttc Phe	agc Ser	aag Lys	aaa Lys	1872
	aac Asn 625	atc Ile	tgc Cys	cag Gln	cca Pro	gtg Val 630	acc Thr	cgg Arg	aaa Lys	acc Thr	ttc Phe 635	atg Met	aaa Lys	aac Asn	aac Asn	ttt Phe 640	1920
	gaa Glu	cac His	atc Ile	cag Gln	cac His 645	att Ile	atc Ile	att Ile	gat Asp	gac Asp 650	gct Ala	cag Gln	aat Asn	ttc Phe	cgt Arg 655	act Thr	1968
	gaa Glu	gat Asp	gly ggg	gac Asp 660	tgg Trp	tat Tyr	Gly aaa	aaa Lys	gca Ala 665	aag Lys	ttc Phe	atc Ile	act Thr	cga Arg 670	cag Gln	caa Gln	2016
	agg Arg	gat Asp	ggc Gly 675	cca Pro	gga Gly	gtt Val	ctc Leu	tgg Trp 680	atc Ile	ttt Phe	ctg Leu	gac Asp	tac Tyr 685	ttt Phe	cag Gln	acc Thr	2064
	tat Tyr	cac His 690	ttg Leu	agt Ser	tgc Cys	agt Ser	ggc Gly 695	ctc Leu	ccc Pro	cct Pro	ccc Pro	tca Ser 700	gac Asp	cag Gln	tat Tyr	cca Pro	2112
	aga Arg 705	gaa Glu	gag Glu	atc Ile	aac Asn	aga Arg 710	gtg Val	gtc Val	cgc Arg	aat Asn	gca Ala 715	ggt Gly	cca Pro	ata Ile	gct Ala	aat Asn 720	2160
	tac Tyr	cta Leu	caa Gln	caa Gln	gta Val 725	atg Met	cag Gln	gaa Glu	gcc Ala	cga Arg 730	caa Gln	aat Asn	cct Pro	cca Pro	cct Pro 735	aac Asn	2208
	ctc Leu	ccc Pro	cct Pro	999 Gly 740	tcc Ser	ctg Leu	gtg Val	atg Met	ctc Leu 745	tat Tyr	gaa Glu	cct Pro	aaa Lys	tgg Trp 750	gct Ala	caa Gln	2256
	ggt Gly	gtc Val	cca Pro 755	ggc	aac Asn	tta Leu	gag Glu	att Ile 760	att Ile	gaa Glu	gac Asp	ttg Leu	aac Asn 765	ttg Leu	gag Glu	gag Glu	2304
	ata Ile	ctg Leu 770	Ile	tat Tyr	gta Val	gcg Ala	aat Asn 775	Lys	tgc Cys	cgt Arg	ttt Phe	ctc Leu 780	Leu	cgg Arg	aat Asn	ggt Gly	2352
	tat Tyr 785	tct Ser	ccg Pro	aag Lys	gat Asp	att Ile 790	gct Ala	gtg Val	ctt Leu	ttc Phe	acc Thr 795	Lys	gca Ala	agt Ser	gaa Glu	gtg Val 800	2400
	gaa Glu	aaa Lys	tat Tyr	aaa Lys	gac Asp 805	Arg	ctt Leu	cta Leu	aca Thr	gca Ala 810	Met	agg Arg	aag Lys	aga Arg	aaa Lys 815	ctg Leu	2448
	tct Ser	cag Gln	ctc Leu	cat His 820	Glu	gag Glu	tct Ser	gat Asp	ctg Leu 825	Leu	cta Leu	cag Gln	ato Ile	ggt Gly 830	Asp	gcg Ala	2496

WO 02/20569 PCT/US01/28013

											*					
tcg Ser	gat Asp	gtt Val 835	cta Leu	acc Thr	gat Asp	cac His	att Ile 840	gtg Val	ttg Leu	gac Asp	agt Ser	gtc Val 845	tgt Cys	cga Arg	ttt Phe	2544
												cca Pro				2592
cca Pro 865	ccg Pro	gct Ala	gly ggg	gcc Ala	tac Tyr 870	aat Asn	ctt Leu	ctg Leu	ctc Leu	tgt Cys 875	ttg Leu	gct Ala	tct Ser	agg Arg	gca Ala 880	2640
					att Ile						tga					2676
<21	0> <i>-</i>	45														
<21	1>	891														
<21	2>	PRT														
<21	3> 1	Ното	sap:	iens												
<40	0 >	45														
Met 1		Leu	Arg	Ile 5	Asp	Val	Asp	Thr	Asn 10	Phe	Pro	Glu	Cys	Val 15	Val	
1	Ser			5					10			Glu Gln		15		
1 Asp	Ser Ala	Gly	Lys 20	5 Val	Thr	Leu	Gly	Thr 25	10 Gln	Gln	Arg		Glu 30	15 Met	Asp	
Asp	Ser Ala	Gly Leu 35	Lys 20 Arg	5 Val Glu Ser	Thr	Leu Gln Gly	Gly Asn 40	Thr 25 Glu	Gln Ile	Gln Ile	Arg Leu	Gln Arg 45	Glu 30 Ala	15 Met Val	Asp Cys	
Asp Pro	Ser Ala Arg Leu 50	Gly Leu 35	Lys 20 Arg Asn	Val Glu Ser	Thr Lys	Leu Gln Gly	Gly Asn 40	Thr 25 Glu Ile	Ile	Gln Ile Lys	Arg Leu Ala 60	Gln Arg 45	Glu 30 Ala Ile	Met Val Glu	Asp Cys Asn	
Asp Pro Ala Lys	Ala Arg Leu 50	Gly Leu 35 Leu Tyr	Lys 20 Arg Asn	Val Glu Ser	Thr Lys Gly Glu 70	Gln Gly 55,	Gly Asn 40 Gly	Thr 25 Glu Ile Gly	Gln Ile Ile Val	Gln Ile Lys Gly	Arg Leu Ala 60 Leu	Gln Arg 45 Glu	Glu 30 Ala Ile Val	Met Val Glu Pro	Asp Cys Asn Pro 80	
Asp Pro Ala Lys 65	Ala Arg Leu 50 Gly	Gly Leu 35 Leu Tyr	Lys 20 Arg Asn Asn	Val Glu Ser Tyr His	Thr Lys Gly Glu 70 Leu	Gln Gly 55 Arg Asp	Gly Asn 40 Gly His	Thr 25 Glu Ile Gly Met	Gln Ile Val Gln 90 Ala	Gln Ile Lys Gly 75 Lys	Arg Leu Ala 60 Leu Glu	Gln Arg 45 Glu Asp	Glu 30 Ala Ile Val :	Met Val Glu Pro	Asp Cys Asn Pro 80	

Leu Cys Ser Asn Leu Tyr His Arg Glu Arg Thr Ser Thr Asp Val Met

		115					120					125			
Asp	Ser 130	Gln	Glu	Ala	Leu	Ala 135	Phe	Leu	Lys	Cys	Arg 140	Thr	Gln	Thr	Pro
Thr 145	Asn	Ile	Asn	Val	Ser 150	Asn	Ser	Leu	Gly	Pro 155	Gln	Ala	Ala	Gln	Gly 160
Ser	Val	Gln	Tyr	Glu 165	Gly	Asn	Ile	Asn	Val 170	Ser	Ala	Ala	Ala	Leu 175	Phe
Asp	Arg	ГÀЗ	Arg 180	Leu	Gln	Tyr	Leu	Glu 185	Lys	Leu	Asn	Leu	Pro 190	Glu	Ser
Thr	His	Val 195	Glu	Phe	Val	Met	Phe 200	Ser	Thr	Asp	Val	Ser 205	His	Cys	Val
Lys	Asp 210	Arg	Leu	Pro	rys	Cys 215	Val	Ser	Ala	Phe	Ala 220	Asn	Thr	Glu	Gly
Gly 225	Tyr	Val	Phe	Phe	Gly 230	Val	His	Asp	Glu	Thr 235	Cys	Gln	Val	Ile	Gly 240
Сув	Glu	Lys	Glu	Lys 245	Ile	Asp	Leu	Thr	Ser 250	Leu	Arg	Ala	Ser	Ile 255	Asp
Gly	Суз	Ile	Lys 260	Lys	Leu	Pro	Val	His 265	His	Phe	Cys	Thr	Gln 270	Arg	Pro
Glu	Ile	Lys 275	Tyr	Val	Leu	Asn	Phe 280	Leu	Glu	Val	His	Asp 285	Lys	Gly	Ala
Leu	Arg 290	Gly	Tyr	Val	Cys	Ala 295	Ile	Lys	Val	Glu	Lys 300	Phe	Cys	Cys	Ala
Val 305	Phe	Ala	Lys	Val	Pro 310	Ser	Ser	Trp	Gln	Val 315		Asp	Asn	Arg	Val
Arg	Gln	Leu	Pro	Thr 325	Arg	Glu	Trp	Thr	Ala 330	Trp	Met	Met	Glu	Ala 335	Asp
Pro	Asp	Leu	Ser 340	Arg	Cys	Pro	Glu	Met 345	Val	Leu	Gln	Leu	Ser 350	Leu	Ser

95

Ser Ala Thr Pro Arg Ser Lys Pro Val Cys Ile His Lys Asn Ser Glu 355 360 365

Cys Leu Lys Glu Gln Gln Lys Arg Tyr Phe Pro Val Phe Ser Asp Arg 370 375 380

Val Val Tyr Thr Pro Glu Ser Leu Tyr Lys Glu Leu Phe Ser Gln His 385 390 395 400

Lys Gly Leu Arg Asp Leu Ile Asn Thr Glu Met Arg Pro Phe Ser Gln 405 410 415

Gly Ile Leu Ile Phe Ser Gln Ser Trp Ala Val Asp Leu Gly Leu Gln 420 425 430

Glu Lys Gln Gly Val Ile Cys Asp Ala Leu Leu Ile Ser Gln Asn Asn 435 440 445

Thr Pro Ile Leu Tyr Thr Ile Phe Ser Lys Trp Asp Ala Gly Cys Lys 450 455 460

Gly Tyr Ser Met Ile Val Ala Tyr Ser Leu Lys Gln Lys Leu Val Asn 465 470 475 480

Lys Gly Gly Tyr Thr Gly Arg Leu Cys Ile Thr Pro Leu Val Cys Val
485 490 495

Leu Asn Ser Asp Arg Lys Ala Gln Ser Val Tyr Ser Ser Tyr Leu Gln 500 505 510

Ile Tyr Pro Glu Ser Tyr Asn Phe Met Thr Pro Gln His Met Glu Ala 515 520 525

Leu Leu Gln Ser Leu Val Ile Val Leu Leu Gly Phe Lys Ser Phe Leu 530 535 540

Ser Glu Glu Leu Gly Ser Glu Val Leu Asn Leu Leu Thr Asn Lys Gln 545 550 550 560

Tyr Glu Leu Leu Ser Lys Asn Leu Arg Lys Thr Arg Glu Leu Phe Val 565 570 575

His Gly Leu Pro Gly Ser Gly Lys Thr Ile Leu Ala Leu Arg Ile Met 580 585 590

Glu Lys Ile Arg Asn Val Phe His Cys Glu Pro Ala Asn Ile Leu Tyr 595 600 605

Ile Cys Glu Asn Gln Pro Leu Lys Lys Leu Val Ser Phe Ser Lys Lys 610 615 . 620

Asn Ile Cys Gln Pro Val Thr Arg Lys Thr Phe Met Lys Asn Asn Phe 625 630 635

Glu His Ile Gln His Ile Ile Ile Asp Asp Ala Gln Asn Phe Arg Thr 645 650 655

Glu Asp Gly Asp Trp Tyr Gly Lys Ala Lys Phe Ile Thr Arg Gln Gln 660 665 670

Arg Asp Gly Pro Gly Val Leu Trp Ile Phe Leu Asp Tyr Phe Gln Thr 675 680 685

Tyr His Leu Ser Cys Ser Gly Leu Pro Pro Pro Ser Asp Gln Tyr Pro 690 695 700

Arg Glu Glu Ile Asn Arg Val Val Arg Asn Ala Gly Pro Ile Ala Asn 705 710 715 720

Tyr Leu Gln Gln Val Met Gln Glu Ala Arg Gln Asn Pro Pro Pro Asn 725 730 735

Leu Pro Pro Gly Ser Leu Val Met Leu Tyr Glu Pro Lys Trp Ala Gln 740 745 750

Gly Val Pro Gly Asn Leu Glu Ile Ile Glu Asp Leu Asn Leu Glu Glu 755 760 765

Ile Leu Ile Tyr Val Ala Asn Lys Cys Arg Phe Leu Leu Arg Asn Gly 770 775 780

Tyr Ser Pro Lys Asp Ile Ala Val Leu Phe Thr Lys Ala Ser Glu Val 785 790 795 800

Glu Lys Tyr Lys Asp Arg Leu Leu Thr Ala Met Arg Lys Arg Lys Leu 805 810 815

Ser Gln Leu His Glu Glu Ser Asp Leu Leu Leu Gln Ile Gly Asp Ala 820 825 830

Ser Asp Val Leu Thr Asp His Ile Val Leu Asp Ser Val Cys Arg Phe 835 840 845

Ser Gly Leu Glu Arg Asn Ile Val Phe Gly Ile Asn Pro Gly Val Ala 850 855 860

Pro Pro Ala Gly Ala Tyr Asn Leu Leu Cys Leu Ala Ser Arg Ala 865 870 875 880

Lys Arg His Leu Tyr Ile Leu Lys Ala Ser Val 885 890

<210> 46

<211> 1737

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1734)

<223>

atg aac atc agt gtt gat ttg gaa acg aat tat gcc gag ttg gtt cta Met Asn Ile Ser Val Asp Leu Glu Thr Asn Tyr Ala Glu Leu Val Leu gat gtg gga aga gtc act ctt gga gag aac agt agg aaa aaa atg aag 96 Asp Val Gly Arg Val Thr Leu Gly Glu Asn Ser Arg Lys Lys Met Lys 25 gat tgt aaa ctg aga aaa aag cag aat gaa agg gtc tca cga gct atg 144 Asp Cys Lys Leu Arg Lys Lys Gln Asn Glu Arg Val Ser Arg Ala Met 40 tgt gct ctg ctc aat tct gga ggg gga gtg atc aag gct gaa att gag 192 Cys Ala Leu Leu Asn Ser Gly Gly Gly Val Ile Lys Ala Glu Ile Glu 240 aat gaa gac tat agt tat aca aaa gat gga ata gga cta gat ttg gaa Asn Glu Asp Tyr Ser Tyr Thr Lys Asp Gly Ile Gly Leu Asp Leu Glu

										•						
aat Asn	tct Ser	ttt Phe	agt Ser	aac Asn 85	att Ile	ctg Leu	tta Leu	ttt Phe	gtt Val 90	cct Pro	gag Glu	tac Tyr	tta Leu	gac Asp 95	ttc Phe	288
				aac Asn												336
aac Asn	acc Thr	tct Ser 115	ggt Gly	ctg Leu	cgg Arg	att Ile	acc Thr 120	acc Thr	ttg Leu	agc Ser	tcc Ser	aat Asn 125	ttg Leu	tac Tyr	aaa Lys	384
				tct Ser												432
ttc Phe 145	ctc Leu	aaa Lys	gac Asp	atg Met	aaa Lys 150	aag Lys	act Thr	aga Arg	gly aaa	aga Arg 155	ttg Leu	tat Tyr	tta Leu	aga Arg	cca Pro 160	480
gaa Glu	ttg Leu	ctg Leu	gca Ala	aag Lys 165	agg Arg	ccc Pro	tgt Cys	gtt Val	gat Asp 170	ata Ile	caa Gln	gaa Glu	gaa Glu	aat Asn 175	aac Asn	528
atg Met	aag Lys	gcc Ala	ttg Leu 180	gcc Ala	gly aaa	gtt Val	ttt Phe	ttt Phe 185	gat Asp	aga Arg	aca Thr	gaa Glu	ctt Leu 190	gat Asp	cgg Arg	576
aaa Lys	gaa Glu	aaa Lys 195	ttg Leu	acc Thr	ttt Phe	act Thr	gaa Glu 200	tcc Ser	aca Thr	cat His	gtt Val	gaa Glu 205	att Ile	aaa Lys	aac Asn	624
ttc Phe	tcg Ser 210	aca Thr	gaa Glu	aag Lys	ttg Leu	tta Leu 215	caa Gln	cga Arg	att Ile	aaa Lys	gag Glu 220	att Ile	ctc Leu	cct Pro	caa Gln	672
tat Tyr 225	.gtt Val	tct Ser	gca Ala	ttt Phe	gca Ala 230	aat Asn	act Thr	gat Asp	gga Gly	gga Gly 235	tat Tyr	ttg Leu	ttc Phe	att Ile	ggt Gly 240	720
tta Leu	aat Asn	Glu	Asp	aaa Lys 245	Glu	Ile	Ile	Gly	Phe	Lys	Ala	Glu	atg Met	agt Ser 255	Asp	768
ctc Leu	gat Asp	gac Asp	tta Leu 260	gaa Glu	aga Arg	gaa Glu	atc Ile	gaa Glu 265	aag Lys	tcc Ser	att Ile	agg Arg	aag Lys 270	atg Met	cct Pro	816
gtg Val	cat His	cac His 275	Phe	tgt Cys	atg Met	gag Glu	aag Lys 280	aag Lys	aag Lys	ata Ile	aat Asn	tat Tyr 285	Ser	tgc Cys	aaa Lys	864
ttc Phe	ctt Leu 290	gga Gly	gta Val	tat Tyr	gat Asp	ааа Lys 295	gga Gly	agt Ser	ctt Leu	tgt Cys	gga Gly 300	Tyr	gtc Val	tgt Cys	gca Ala	912
ctc Leu 305	aga Arg	gtg Val	gag Glu	cgc Arg	ttc Phe 310	Cys	tgt Cys	gca Ala	gtg Val	Phe 315	Ala	aaa Lys	gag Glu	cct Pro	gat Asp 320	960

								gtg Val								1008
								gaa Glu 345								1056
_								acg Thr				_				1104
								cag Gln								1152
								gaa Glu								1200
								tta Leu								1248
								tct Ser 425								1296
								ctc Leu								1344
cag Gln	gac Asp 450	agt Ser	cct Pro	cca Pro	gtc Val	cta Leu 455	tac Tyr	acc Thr	ttc Phe	cac His	atg Met 460	gta Val	cag Gln	gat Asp	gag Glu	1392
gag Glu 465	ttt Phe	aaa Lys	ggc Gly	tat Tyr	tct Ser 470	aca Thr	caa Gln	act Thr	gcc Ala	cta Leu 475	acc Thr	tta Leu	aag Lys	cag Gln	aag Lys 480	1440
ctg Leu	gca Ala	aaa Lys	att Ile	ggt Gly 485	Gly	tac Tyr	act Thr	aaa Lys	aaa Lys 490	gtg Val	tgt Cys	gtc Val	atg Met	aca Thr 495	aag Lys	1488
								atg Met 505								1536
agg Arg	tcg Ser	caa Gln 515	gta Val	att Ile	tac Tyr	cct Pro	gaa Glu 520	tcc Ser	tac Tyr	tat Tyr	ttt Phe	aca Thr 525	aga Arg	agg Arg	aaa Lys	1584
								gcc Ala								1632
aga Arg	gac Asp	cag Gln	ttt Phe	tcc Ser	ttt Phe	gca Ala	gaa Glu	aat Asn	cta Leu	tac Tyr	cag Gln	ata Ile	atc Ile	ggt Gly	ata Ile	1680

100

560 555 545 550 gat tgc ttt cag aag aat gat aaa aag atg ttt aaa tct tgt cga agg 1728 Asp Cys Phe Gln Lys Asn Asp Lys Lys Met Phe Lys Ser Cys Arg Arg ctc acc tga 1737 Leu Thr <210> 47 <211> 578 <212> PRT <213> Homo sapiens <400> 47 Met Asn Ile Ser Val Asp Leu Glu Thr Asn Tyr Ala Glu Leu Val Leu 5 Asp Val Gly Arg Val Thr Leu Gly Glu Asn Ser Arg Lys Lys Met Lys 25 20 Asp Cys Lys Leu Arg Lys Lys Gln Asn Glu Arg Val Ser Arg Ala Met 35 Cys Ala Leu Leu Asn Ser Gly Gly Gly Val Ile Lys Ala Glu Ile Glu 50 Asn Glu Asp Tyr Ser Tyr Thr Lys Asp Gly Ile Gly Leu Asp Leu Glu Asn Ser Phe Ser Asn Ile Leu Leu Phe Val Pro Glu Tyr Leu Asp Phe 85 Met Gln Asn Gly Asn Tyr Phe Leu Ile Phe Val Lys Ser Trp Ser Leu 100 Asn Thr Ser Gly Leu Arg Ile Thr Thr Leu Ser Ser Asn Leu Tyr Lys

Arg Asp Ile Thr Ser Ala Lys Val Met Asn Ala Thr Ala Ala Leu Glu

140

115

101

Phe Leu Lys Asp Met Lys Lys Thr Arg Gly Arg Leu Tyr Leu Arg Pro 145 150 155 160

Glu Leu Leu Ala Lys Arg Pro Cys Val Asp Ile Gln Glu Glu Asn Asn 165 170 175

Met Lys Ala Leu Ala Gly Val Phe Phe Asp Arg Thr Glu Leu Asp Arg 180 185 190

Lys Glu Lys Leu Thr Phe Thr Glu Ser Thr His Val Glu Ile Lys Asn 195 200 205

Phe Ser Thr Glu Lys Leu Gln Arg Ile Lys Glu Ile Leu Pro Gln 210 215 220

Tyr Val Ser Ala Phe Ala Asn Thr Asp Gly Gly Tyr Leu Phe Ile Gly 225 230 235 240

Leu Asn Glu Asp Lys Glu Ile Ile Gly Phe Lys Ala Glu Met Ser Asp 245 250 255

Leu Asp Asp Leu Glu Arg Glu Ile Glu Lys Ser Ile Arg Lys Met Pro 260 265 270

Val His His Phe Cys Met Glu Lys Lys Lys Ile Asn Tyr Ser Cys Lys 275 280 285

Phe Leu Gly Val Tyr Asp Lys Gly Ser Leu Cys Gly Tyr Val Cys Ala 290 295 300

Leu Arg Val Glu Arg Phe Cys Cys Ala Val Phe Ala Lys Glu Pro Asp 305 310 315 320

Ser Trp His Val Lys Asp Asn Arg Val Met Gln Leu Thr Arg Lys Glu 325 330 335

Trp Ile Gln Phe Met Val Glu Ala Glu Pro Lys Phe Ser Ser Tyr 340 345 350

Glu Glu Val Ile Ser Gln Ile Asn Thr Ser Leu Pro Ala Pro His Ser 355 360 365

Trp Pro Leu Leu Glu Trp Gln Arg Gln Arg His His Cys Pro Gly Leu 370 375 380

102

Ser Gly Arg Ile Thr Tyr Thr Pro Glu Asn Leu Cys Arg Lys Leu Phe 390 395 Leu Gln His Glu Gly Leu Lys Gln Leu Ile Cys Glu Glu Met Asp Ser 405 410 . 415 Val Arg Lys Gly Ser Leu Ile Phe Ser Arg Ser Trp Ser Val Asp Leu 425 430 420 Gly Leu Gln Glu Asn His Lys Val Leu Cys Asp Ala Leu Leu Ile Ser Gln Asp Ser Pro Pro Val Leu Tyr Thr Phe His Met Val Gln Asp Glu 455 Glu Phe Lys Gly Tyr Ser Thr Gln Thr Ala Leu Thr Leu Lys Gln Lys 465 470 475 Leu Ala Lys Ile Gly Gly Tyr Thr Lys Lys Val Cys Val Met Thr Lys 485 490 Ile Phe Tyr Leu Ser Pro Glu Gly Met Thr Ser Cys Gln Tyr Asp Leu 500 505 510 Arg Ser Gln Val Ile Tyr Pro Glu Ser Tyr Tyr Phe Thr Arg Arg Lys 515 520 Tyr Leu Leu Lys Ala Leu Phe Lys Ala Leu Lys Arg Leu Lys Ser Leu 530 535 540 Arg Asp Gln Phe Ser Phe Ala Glu Asn Leu Tyr Gln Ile Ile Gly Ile 545 550 555 Asp Cys Phe Gln Lys Asn Asp Lys Lys Met Phe Lys Ser Cys Arg Arg 565 570 575

Leu Thr

<210> 48

<211> 2694

<212> DNA

103

<213> Homo sapiens

<220>

<221> CDS .

<222> (1)..(2691)

<223>

<400> 48						
atg gag gca Met Glu Ala 1						48
ctg gtc atc Leu Val Ile						96
aag cta cag Lys Leu Gln 35						144
gcc gcg tgt Ala Ala Cys 50						192
atg gcc aac Met Ala Asn 65	agg gat gag Arg Asp Glu 70	cgt ccc aca Arg Pro Thi	a gag atg Glu Met 75	gga ctg gat Gly Leu Asp	tta gaa Leu Glu 80	240
gaa tcc ttg Glu.Ser Leu	aga aag ctt Arg Lys Leu 85	att cag tat Ile Gln Tyr	cca tat Pro Tyr 90	ttg cag gct Leu Gln Ala	ttc ttt Phe Phe 95	288
gag act aag Glu Thr Lys	caa cac gga Gln His Gly 100	agg tgt ttt Arg Cys Phe 105	Tyr Ile	ttt gtt aaa Phe Val Lys 110	Ser Trp	336
agt ggt gat Ser Gly Asp 115						384
agc ctt agt Ser Leu Ser 130						432
atg aat tca Met Asn Ser 145						480
tcc aaa tat Ser Lys Tyr	aat ctg att Asn Leu Ile 165	aat gaa gg Asn Glu Gly	g tct cca y Ser Pro 170	cct agt aaa Pro Ser Lys	att atg Ile Met 175	528

										•							
aaa Lys	gct Ala	gta Val	tac Tyr 180	cag Gln	aac Asn	ata Ile	tct Ser	gag Glu 185	tca Ser	aat Asn	cct Pro	gca Ala	tat Tyr 190	gaa Glu	gtt Val		576
ttc Phe	caa Gln	act Thr 195	gac Asp	act Thr	att Ile	gaa Glu	tat Tyr 200	ggt Gly	gaa Glu	atc Ile	cta Leu	tct Ser 205	ttt Phe	cct Pro	gag Glu		624
tct Ser	cca Pro 210	tcc Ser	ata Ile	gag Glu	ttt Phe	aaa Lys 215	cag Gln	ttc Phe	tct Ser	aca Thr	aaa Lys 220	cat His	atc Ile	caa Gln	caa Gln		672
tat Tyr 225	gta Val	gaa Glu	aat Asn	ata Ile	att Ile 230	cca Pro	gag Glu	tac Tyr	atc Ile	tct Ser 235	gca Ala	ttt Phe	gca Ala	aac Asn	act Thr 240		720
gag Glu	gga Gly	ggc	tat Tyr	ctt Leu 245	ttt Phe	att Ile	gga Gly	gtg Val	gat Asp 250	gat Asp	aag Lys	agt Ser	agg Arg	aaa Lys 255	gtc Val		768
ctg Leu	gga Gly	tgt Cys	gcc Ala 260	aaa Lys	gaa Glu	cag Gln	gtt Val	gac Asp 265	cct Pro	gac Asp	tct Ser	ttg Leu	aaa Lys 270	aat Asn	gta Val		816
att Ile	gca Ala	aga Arg 275	gca Ala	att Ile	tct Ser	aag Lys	ttg Leu 280	ccc Pro	att Ile	gtt Val	cat His	ttt Phe 285	tgc Cys	tct Ser	tca Ser		864
aaa Lys	cct Pro 290	cgg Arg	gta Val	gag Glu	tac Tyr	agc Ser 295	acc Thr	aaa Lys	atc Ile	gta Val	gaa Glu 300	gtg Val	ttt Phe	tgt Cys	gly ggg		912
aaa Lys 305	gag Glu	ttg Leu	tat Tyr	ggc	tat Tyr 310	ctc Leu	tgt Cys	gtg Val	att Ile	aaa Lys 315	gtg Val	aag Lys	gca Ala	ttc Phe	tgt Cys 320		960
tgt Cys	gtg Val	gtg Val	ttc Phe	tcg Ser 325	gaa Glu	gct Ala	ccc Pro	aag Lys	tca Ser 330	tgg Trp	atg Met	gtg Val	agg Arg	gag Glu 335	aag Lys		1008
tac Tyr	atc Ile	cgc Arg	ccc Pro 340	ttg Leu	aca Thr	act Thr	gag Glu	gaa Glu 345	tgg Trp	gta Val	gag Glu	aaa Lys	atg Met 350	atg Met	gac Asp		1056
gca Ala	gat Asp	cca Pro 355	Glu	ttt Phe	cct Pro	cca Pro	gac Asp 360	ttt Phe	gct Ala	gag Glu	gcc Ala	ttt Phe 365	gag Glu	tct Ser	cag Gln		1104
ttg Leu	agt Ser 370	Leu	tct Ser	gac Asp	agt Ser	cct Pro 375	tca Ser	ctt Leu	tgc Cys	aga Arg	cca Pro 380	gtg Val	tat Tyr	tct Ser	aag Lys	-	1152
aaa Lys 385	Gly	ctg Leu	gaa Glu	cac His	aaa Lys 390	gct Ala	gat Asp	cta Leu	caa Gln	caa Gln 395	His	tta Leu	ttt Phe	cca Pro	gtt Val 400		1200
cca Pro	cca Pro	gga Gly	cat His	ttg Leu 405	gaa Glu	tgt Cys	act Thr	cca Pro	gag Glu 410	Ser	ctc Leu	tgg Trp	aag Lys	gag Glu 415	ctg Leu		1248

tct Ser	tta Leu	cag Gln	cat His 420	gaa Glu	gga Gly	cta Leu	aag Lys	gag Glu 425	tta Leu	ata Ile	cac His	aag Lys	caa Gln 430	atg Met	cga Arg	1296
cct Pro	ttc Phe	tcc Ser 435	cag	gga Gly	att Ile	gtg Val	atc Ile 440	ctc Leu	tct Ser	aga Arg	agc Ser	tgg Trp 445	gct	gtg Val	gac Asp	1344
ctg Leu	aac Asn 450	ttg Leu	cag Gln	gag Glu	aag Lys	cca Pro 455	gga Gly	gtc Val	atc Ile	tgt Cys	gat Asp 460	gct Ala	ctg Leu	ctg Leu	ata Ile	1392
gca Ala 465	cag Gln	aac Asn	agc Ser	acc Thr	ccc Pro 470	att Ile	ctc Leu	tac Tyr	acc Thr	att Ile 475	ctc Leu	agg Arg	gag Glu	cag Gln	gat Asp 480	1440
gca Ala	gag Glu	ggc Gly	cag Gln	gac Asp 485	tac Tyr	tgc Cys	act Thr	cgc Arg	acc Thr 490	gcc Ala	ttt Phe	act Thr	ttg Leu	aag Lys 495	cag Gln	1488
aag Lys	cta Leu	gtg Val	aac Asn 500	atg Met	gly ggg	ggc Gly	tac Tyr	acc Thr 505	gly aaa	aag Lys	gtg Val	tgt Cys	gtc Val 510	agg Arg	gcc Ala	1536
aag Lys	gtc Val	ctc Leu 515	tgc Cys	ctg Leu	agt Ser	cct Pro	gag Glu 520	agc Ser	agc Ser	gca Ala	gag Glu	gcc Ala 525	ttg Leu	gag Glu	gct Ala	1584
gca Ala	gtg Val 530	tct Ser	ccg Pro	atg Met	gat Asp	tac Tyr 535	cct Pro	gcg Ala	tcc Ser	tat Tyr	agc Ser 540	ctt Leu	gca Ala	ggc	acc Thr	1632
cag Gln 545	cac His	atg Met	gaa Glu	gcc Ala	ctg Leu 550	ctg Leu	cag Gln	tcc Ser	ctc Leu	gtg Val 555	att Ile	gtc Val	tta Leu	ctc Leu	ggc Gly 560	1680
ttc Phe	agg Arg	tct Ser	ctc Leu	ttg Leu 565	Ser	gac Asp	cag Gln	ctc Leu	ggc Gly 570	tgt Cys	gag Glu	gtt Val	tta Leu	aat Asn 575	Leu	1728
ctc Leu	aca Thr	gcc Ala	cag Gln 580	cag Gln	tat Tyr	gag Glu	ata Ile	ttc Phe 585	tcc Ser	aga Arg	agc Ser	ctc Leu	cgc Arg 590	Lys	aac Asn	1776
aga Arg	gag Glu	ttg Leu 595	Phe	gtc Val	cac His	ggc	tta Leu 600	cct Pro	ggc	tcạ Ser	gly ggg	aag Lys 605	Thr	atc Ile	atg Met	1824
gcc Ala	atg Met 610	Lys	atc Ile	atg Met	gag Glu	aag Lys 615	atc	agg Arg	aat Asn	gtg Val	ttt Phe 620	His	tgt Cys	gag Glu	gca Ala	1872
cac His 625	Arg	att Ile	ctc Leu	tac Tyr	gtt Val 630	Cys	gaa Glu	aac Asn	cag Gln	cct Pro 635	Leu	agg Arg	aac Asr	ttt Phe	atc : Ile 640	1920
agt Ser	gat Asp	aga Arg	aat Asn	atc Ile	tgc Cys	cga Arg	gca Ala	gag Glu	acc Thr	cgg Arg	gaa Glu	act Thr	tto Phe	cta Lev	aga Arg	1968

				645					650					655			
gaa	aaa	ttt	gaa	cac	att	caa	cac	atc	gtc	att	gac	gaa	gct	cag	aat	2	016
Glu	Lys	Phe	Glu 660	His	Ile	GIn	His	11e 665	Val	TTE	Asp	Glu	670	Gin	Asn		
ttc Phe	cgt Arg	act Thr 675	gaa Glu	gat Asp	gly ggg	gac Asp	tgg Trp 680	tat Tyr	agg Arg	aag Lys	gca Ala	aaa Lys 685	acc Thr	atc Ile	act Thr	2	064
Gln	Arg 690	Glu	Lys	Asp	tgt Cys	Pro 695	Gly	Val	Leu	Trp	Ile 700	Phe	Leu	Asp	Tyr	2	112
					ttg Leu 710											2	160
					gag Glu											2	208
					caa Gln											2	256
cca Pro	att Ile	aat Asn 755	atc Ile	ccc Pro	cat His	glà aaa	tat Tyr 760	ctg Leu	gca Ala	att Ile	ctc Leu	agt Ser 765	gaa Glu	gct Ala	aaa Lys	2	304
					cca Pro											2	352
ttg Leu 785	gag Glu	caa Gln	ata Ile	gtg Val	acc Thr 790	tat Tyr	gtg Val	gca Ala	gac Asp	acc Thr 795	tgc Cys	agg Arg	tgc Cys	ttc Phe	ttt Phe 800	2	400
					cca Pro											2	448
					tat Tyr											2	496
aaa Lys	atg Met	gtg Val 835	gtg Val	cag Gln	ctc Leu	agt Ser	gat Asp 840	gca Ala	tgt Cys	gat Asp	atg Met	ttg Leu 845	Gly	gtg Val	cac His	2	2544
att Ile	gtg Val 850	ttg Leu	gac Asp	agt Ser	gtc Val	cgg Arg 855	cga Arg	ttc Phe	tca Ser	ggc	ctg Leu 860	gaa Glu	agg Arg	agc Ser	ata Ile	2	2592
gtg Val 865	ttt Phe	gly	atc Ile	cat His	cca Pro 870	agg Arg	aca Thr	gct Ala	gac Asp	cca Pro 875	Ala	atc Ile	tta Leu	ecc Pro	aat Asn 880	2	2640
att	ctg	atc	tgt	ctg	gct	tcc	agg	gca	aaa	cag	cac	cta	tat	att	ttt	. 2	2688

107

Ile Leu Ile Cys Leu Ala Ser Arg Ala Lys Gln His Leu Tyr Ile Phe 885 890 895

ctg tga · 2694 Leu

<210> 49

<211> 897

<212> PRT

<213> Homo sapiens

<400> 49

Met Glu Ala Asn His Cys Ser Leu Gly Val Tyr Pro Ser Tyr Pro Asp 1 5 10 15

Leu Val Ile Asp Val Gly Glu Val Thr Leu Gly Glu Glu Asn Arg Lys 20 25 30

Lys Leu Gln Lys Thr Gln Arg Asp Gln Glu Arg Ala Arg Val Ile Arg 35 40 45

Ala Ala Cys Ala Leu Leu Asn Ser Gly Gly Gly Val Ile Gln Met Glu 50 55 60

Met Ala Asn Arg Asp Glu Arg Pro Thr Glu Met Gly Leu Asp Leu Glu 65 70 75 80

Glu Ser Leu Arg Lys Leu Ile Gln Tyr Pro Tyr Leu Gln Ala Phe Phe 85 90 95

Glu Thr Lys Gln His Gly Arg Cys Phe Tyr Ile Phe Val Lys Ser Trp 100 105 110

Ser Gly Asp Pro Phe Leu Lys Asp Gly Ser Phe Asn Ser Arg Ile Cys 115 120 125

Ser Leu Ser Ser Ser Leu Tyr Cys Arg Ser Gly Thr Ser Val Leu His 130 135 140

Met Asn Ser Arg Gln Ala Phe Asp Phe Leu Lys Thr Lys Glu Arg Gln 145 150 155 160

ser	гÀг	ıyr	Asn	165	116	ASI	GIU	GIÀ	170	PIO	Pro	261	пув	175	Mec
Lys	Ala	Val	Tyr 180	Gln	Asn	Ile	Ser	Glu 185	Ser	Asn	Pro	Ala	Tyr 190	Glu	Val
Phe	Gln	Thr 195	Asp	Thr	Ile	Glu	Tyr 200	Gly	Ġlu	Ile	Leu	Ser 205	Phe	Pro	Glu
Ser	Pro 210	Ser	Ile	Glu	Phe	Lys 215	Gln	Phe	Ser	Thr	Lys 220	His	Ile	Gln	Gln
Tyr 225	Val	Glu	Asn	Ile	Ile 230	Pro	Glu	Tyr	Ile	Ser 235	Ala	Phe	Ala	Asn	Thr 240
Glu	Gly	Gly	Tyr	Leu 245	Phe	Ile	Gly	Val	Asp 250	Asp	Lys	Ser	Arg	Lys 255	Val
Leu	Gly	Cys	Ala 260	Lys	Glu	Gln	Val	Asp 265	Pro	Asp	Ser	Leu	Lys 270	Asn	Val
Ile	Ala	Arg 275	Ala	Ile	Ser	Lys	Leu 280	Pro	Ile	Val	His	Phe 285	Cys	Ser	Ser
Lys	Pro 290	Arg	Val	Glu	Tyr	Ser 295	Thr	Lys	Ile	Val	Glu 300	Val	Phe	Сув	Gly
Lys 305	Glu	Leu	Tyr	Gly	Tyr 310	Leu	Cys	Val	Ile	Lys 315	Val	Lys	Ala	Phe	Cys 320
Cys	Val	Val	Phe	Ser 325	Glu	Ala	Pro	Lys	Ser 330	Trp	Met	Val	Arg	Glu 335	Lys
Tyr	Ile	Arg	Pro 340	Leu	Thr	Thr	Glu	Glu 345		Val	Glu	Lys	Met 350	Met	Asp
Ala	Asp	Pro 355	Glu	Phe	Pro	Pro	Asp 360	Phe	Ala	Glu	Ala	Phe 365	Glu	Ser	Gln
Leu	Ser 370	Leu	Ser	Asp	Ser	Pro 375		Leu	Cys	Arg	Pro 380		Tyr	Ser	Lys
Lys 385	Gly	Leu	Glu	His	Lys 390	Ala	Asp	Leu	Gln	Gln 395		Leu	Phe	Pro	Val 400

Pro	Pro	Gly	His	Leu 405	Glu	Cys	Thr	Pro	410	ser	Leu	Trp	гÀг	415	ьeu
Ser	Leu	Gln	His 420	Glu	Gly	Leu	Lys	Glu 425	Leu	Ile	His	Lys	Gln 430	Met	Arg
Pro	Phe	Ser 435	Gln	Gly	Ile	Val	Ile 440	Leu	Ser	Arg	Ser	Trp 445	Ala	Val	Asp
Leu	Asn 450	Leu	Gln	Glu	Lys	Pro 455	Gly	Val	Ile	Сув	Asp 460	Ala	Leu	Leu	Ile
465					470					475				Gln	480
				485					490					Lys 495	
_			500					505					510	Arg	
		515					520					525		Glu	
	530					535					540			Gly	
545					550					555				Leu	560
				565					570					Asn 575	
Leu	Thr	Ala	GIN	GIN	Tyr	GIU	тте	Pne	ser	Arg	ser	ьeu	AT 9	Lys	Hall

Arg Glu Leu Phe Val His Gly Leu Pro Gly Ser Gly Lys Thr Ile Met 595 600 605

Ala Met Lys Ile Met Glu Lys Ile Arg Asn Val Phe His Cys Glu Ala 610 615 620

His Arg Ile Leu Tyr Val Cys Glu Asn Gln Pro Leu Arg Asn Phe Ile

625				·	630					635					640
Ser	Asp	Arg	Asn	Ile 645	Cys	Arg	Ala	Glu	Thr 650	Arg	Glu	Thr	Phe	Leu 655	Arg
Glu	Lys	Phe	Glu 660	His	Ile	Gl'n	His	Ile 665	Val	Ile	Asp	Glu	Ala 670	Gln	Asn
Phe	Arg	Thr 675	Glu	Asp	Gly	Asp	Trp 680	Tyr	Arg	Lys	Ala	Lys 685	Thr	Ile	Thr
Gln	Arg 690	Glu	Lys	Asp	Cys	Pro 695	Gly	Val	Leu	Trp	Ile 700	Phe	Leu	Asp	Tyr
Phe 705	Gln	Thr	Ser	His	Leu 710	Gly	His	Ser	Gly	Leu 715	Pro	Pro	Leu	Ser	Ala 720
Gln	Tyr	Pro	Arg	Glu 725	Glu	Leu	Thr	Arg	Val 730	Val	Arg	Asn	Ala	Asp 735	Glu
Ile	Ala	Glu	Tyr 740	Ile	Gln	Gln	Glu	Met 745	Gln	Leu	Ile	Ile	Glu 750	Asn	Pro
Pro	Ile	Asn 755	Ile	Pro	His	Gly	Tyr 760	Leu	Ala	Ile	Leu	Ser 765	Glu	Ala	Lys
Trp	Val 770	Pro	Gly	Val	Pro	Gly 775	Asn	Thr	Lys	Ile	Ile 780	Lys	Asn	Phe	Thr
Leu 785	Glu	Gln	Ile	Val	Thr 790	Tyr	Val	Ala	Asp	Thr 795	Сув	Arg	Cys	Phe	Phe 800
Glu	Arg	Gly	Tyr	Ser 805	Pro	Lys	Asp	Val	Ala 810		Leu	Val	Ser	Thr 815	Val
Thr	Glu	Val	Glu 820	Gln	Tyr	Gln	Ser	Lys 825	Leu	Leu	Lys	Ala	Met 830	Arg	Lys
Lys	Met	Val 835	Val	Gln	Leu	Ser	Asp 840	Ala	Cys	Asp	Met	Leu 845	Gly	Val	His
Ile	Val 850	Leu	Asp	Ser	Val	Arg 855	Arg	Phe	Ser	Gly	Leu 860	Glu	Arg	Ser	Ile

111

Val Phe Gly Ile His Pro Arg Thr Ala Asp Pro Ala Ile Leu Pro Asn 865 870 875 880

Ile Leu Ile Cys Leu Ala Ser Arg Ala Lys Gln His Leu Tyr Ile Phe 885 890 895

Leu

<210> 50

<211> 1074

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1071)

<223>

<400> 50 atg gag agt Met Glu Ser 1	ctc aag act of Leu Lys Thr	gat act gaa Asp Thr Glu	atg ccg tat Met Pro Tyr 10	cct gag Pro Glu	gta ata Val Ile 15	48
gta gat gtg Val Asp Val	ggc aga gtg Gly Arg Val 20	att ttt gga Ile Phe Gly 25	gaa gaa aac Glu Glu Asn	agg aag Arg Lys 30	aag atg Lys Met	96
acc aac agc Thr Asn Ser 35	tgt ttg aaa Cys Leu Lys	aga tct gag Arg Ser Glu 40	aat tct aga Asn Ser Arg	att atc Ile Ile 45	cgg gct Arg Ala	144
ata tgt gca Ile Cys Ala 50	ctg tta aat Leu Leu Asn	tct gga ggt Ser Gly Gly 55	ggt gtg atc Gly Val Ile 60	aaa gca Lys Ala	gag att Glu Ile	192
gat gat aaa Asp Asp Lys 65	acc tat agt Thr Tyr Ser 70	tac caa tgc Tyr Gln Cys	cat ggg ctg His Gly Leu 75	gga cag Gly Gln	gat ttg Asp Leu 80	240
gaa act tct Glu Thr Ser	ttt caa aag Phe Gln Lys 85	ctc ctt cct Leu Leu Pro	tca ggt tca Ser Gly Ser 90	cag aaa Gln Lys	tac ctt Tyr Leu 95	288
gac tac atg Asp Tyr Met	cag cag ggg . Gln Gln Gly	cac aat ctc His Asn Leu	ctg att ttt Leu Ile Phe	gtg aag Val Lys	tca tgg Ser Trp	_. 336

1	00	105	110	
		-		
	tt ttc agc ctt cc al Phe Ser Leu Pro 12	Leu Arg Ile (
aat ttg tat co Asn Leu Tyr A 130	gg aga gat gtg ac rg Arg Asp Val Th 135	r Ser Ala Ile <i>l</i>	aac ttg agt gct Asn Leu Ser Ala 140	agc 432 Ser
	ag ctt ctc aga ga lu Leu Leu Arg Gl 150			
	tg aag aag ttg ca 'al Lys Lys Leu Hi 165			
Ile Gln Glu G	ag gaa gat atg ag Nu Glu Asp Met Ar 80			
aag gac aaa c Lys Asp Lys L 195	etc atg tat aag ga beu Met Tyr Lys Gl 20	u Lys Leu Asn	ttt act gag tca Phe Thr Glu Ser 205	aca 624 Thr
	tt aaa agg ttc ac The Lys Arg Phe Th 215	r Thr Lys Lys '		
aag gaa atg c Lys Glu Met L 225	etg cct cat tat gt eu Pro His Tyr Va 230	t tct gca ttt g l Ser Ala Phe 2 235	gcc aac act caa Ala Asn Thr Gln	ggg 720 Gly 240
gga tat gtc c Gly Tyr Val L	etc att ggg gtg ga Deu Ile Gly Val As 245	t gat aag agc o p Asp Lys Ser : 250	aaa gaa gtg gtt Lys Glu Val Val 255	gga 768 Gly
Cys Lys Trp G	gaa aaa gtg aat cc Blu Lys Val Asn Pr 160			
	yaa aaa ttg cct ac ilu Lys Leu Pro Th 28	r Phe His Phe		
aag gta aat t Lys Val Asn P 290	tc act aca aaa at Phe Thr Thr Lys Il 295	e Leu Asn Val	tac caa aaa gat Tyr Gln Lys Asp 300	gtc 912 Val
ctg gat ggt t Leu Asp Gly T 305	at gtc tgt gtg at Tyr Val Cys Val Il 310	t caa gtg gag e Gln Val Glu 315	ccc ttc tgt tgc Pro Phe Cys Cys	gtg 960 Val 320
gtg ttt gca g Val Phe Ala G	gag gcc cca gat to Blu Ala Pro Asp Se 325	c tgg atc atg r Trp Ile Met 330	aaa gac aat tct Lys Asp Asn Ser 335	gtc 1008 Val
aca cgg ctg a	aca gct gag cag tg	g gtg gtc atg	atg ctg gat act	cag 1056

113

Thr Arg Leu Thr Ala Glu Gln Trp Val Val Met Met Leu Asp Thr Gln 340 345 . 350

tca ggt aaa ggg aag tga Ser Gly Lys Gly Lys 355 1074

<210> 51

<211> 357

<212> PRT

<213> Homo sapiens

<400> 51

Met Glu Ser Leu Lys Thr Asp Thr Glu Met Pro Tyr Pro Glu Val Ile 1 5 10 15

Val Asp Val Gly Arg Val Ile Phe Gly Glu Glu Asn Arg Lys Lys Met 20 25 30

Thr Asn Ser Cys Leu Lys Arg Ser Glu Asn Ser Arg Ile Ile Arg Ala 35 40 45

Ile Cys Ala Leu Leu Asn Ser Gly Gly Gly Val Ile Lys Ala Glu Ile 50 55 60

Asp.Asp Lys Thr Tyr Ser Tyr Gln Cys His Gly Leu Gly Gln Asp Leu 65 70 75 80

Glu Thr Ser Phe Gln Lys Leu Leu Pro Ser Gly Ser Gln Lys Tyr Leu 85 90 95

Asp Tyr Met Gln Gln Gly His Asn Leu Leu Ile Phe Val Lys Ser Trp 100 105 110

Ser Pro Asp Val Phe Ser Leu Pro Leu Arg Ile Cys Ser Leu Arg Ser 115 120 125

Asn Leu Tyr Arg Arg Asp Val Thr Ser Ala Ile Asn Leu Ser Ala Ser 130 135 140

Ser Ala Leu Glu Leu Leu Arg Glu Lys Gly Phe Arg Ala Gln Arg Gly 145 150 155 160

114

Arg Pro Arg Val Lys Lys Leu His Pro Gln Gln Val Leu Asn Arg Cys 165 170 175

Ile Gln Glu Glu Asp Met Arg Ile Leu Ala Ser Glu Phe Phe Lys
180 185 190

Lys Asp Lys Leu Met Tyr Lys Glu Lys Leu Asn Phe Thr Glu Ser Thr 195 200 205

His Val Glu Phe Lys Arg Phe Thr Thr Lys Lys Val Ile Pro Arg Ile 210 215 220

Lys Glu Met Leu Pro His Tyr Val Ser Ala Phe Ala Asn Thr Gln Gly 225 230 235 240

Gly Tyr Val Leu Ile Gly Val Asp Asp Lys Ser Lys Glu Val Val Gly 245 250 255

Cys Lys Trp Glu Lys Val Asn Pro Asp Leu Leu Lys Lys Glu Ile Glu 260 265 270

Asn Cys Ile Glu Lys Leu Pro Thr Phe His Phe Cys Cys Glu Lys Pro 275 280 285

Lys Val Asn Phe Thr Thr Lys Ile Leu Asn Val Tyr Gln Lys Asp Val 290 295 300

Leu Asp Gly Tyr Val Cys Val Ile Gln Val Glu Pro Phe Cys Cys Val 305 310 315 320

Val Phe Ala Glu Ala Pro Asp Ser Trp Ile Met Lys Asp Asn Ser Val 325 330 335

Thr Arg Leu Thr Ala Glu Gln Trp Val Val Met Met Leu Asp Thr Gln 340 345 350

Ser Gly Lys Gly Lys

<210> 52

<211> 807

<212> DNA

115

<213> Mus musculus

<220>

<221> CDS

<222> (1)..(804)

<223>

<400		52														
					cag Gln											48
agc Ser	aaa Lys	gcc Ala	agg Arg 20	agg Arg	ctg Leu	aag Lys	ctt Leu	ggc Gly 25	ctg Leu	cca Pro	gga Gly	ccc Pro	cca Pro 30	gjå aaa	cca Pro	96
cca Pro	ggt Gly	cct Pro 35	cag Gln	ggc	ccc Pro	cca Pro	ggc Gly 40	ccc Pro	ttt Phe	atc Ile	cca Pro	tct Ser 45	gag Glu	gtt Val	ctg Leu	144
					ctg Leu											192
agc Ser 65	cat His	ctg Leu	gag Glu	cac His	tgc Cys 70	acc Thr	agg Arg	gat Asp	ctc Leu	act Thr 75	aca Thr	cca Pro	gcc Ala	tcg Ser	ggt Gly 80	240
agc Ser	cct Pro	tcc Ser	cgt Arg	gtc Val 85	cca Pro	gcc Ala	gcc Ala	cag Gln	gag Glu 90	ctt Leu	gat Asp	agc Ser	cag Gln	gac Asp 95	cca Pro	. 288
ggg	gca Ala	ttg Leu	tta Leu 100	gct Ala	ctg Leu	ctg Leu	gct Ala	gcg Ala 105	acc Thr	ttg Leu	gcc Ala	cag Gln	ggc Gly 110	ccg Pro	cgg Arg	336
gca Ala	cca Pro	cgt Arg 115	gtg Val	gag Glu	gcc Ala	gca Ala	ttc Phe 120	cac His	tgt Cys	cgc Arg	ttg Leu	cgc Arg 125	cgg Arg	gat Asp	gtg Val	384
cag Gln	gtg Val 130	gat Asp	cgg Arg	cgt Arg	gcg Ala	ttg Leu 135	cac His	gag Glu	ctt Leu	ggg Gly	atc Ile 140	tac Tyr	tac Tyr	ctg Leu	ccc Pro	432
gaa Glu 145	gtt Val	gag Glu	gga Gly	gcc Ala	ttc Phe 150	His	cgg Arg	ggc Gly	cca Pro	ggc Gly 155	ttg Leu	aat Asn	ctg Leu	acc Thr	agc Ser 160	480
ggc	cag Gln	tac Tyr	acc Thr	gca Ala 165	cct Pro	gtg Val	gct Ala	ggc	ttc Phe 170	Tyr	gcg Ala	ctt Leu	gct Ala	gcc Ala 175	act Thr	528

								'	10				•			
		gtg Val														576
cgg Arg	gat Asp	cgt Arg 195	ctg Leu	cgc Arg	ctg Leu	ctg Leu	atc Ile 200	tgc Cys	atc Ile	cag Gln	tct Ser	ctc Leu 205	tgt Cys	cag Gln	cac His	624
aat Asn	gcc Ala 210	tcc Ser	ctg Leu	gag Glu	act Thr	gtg Val 215	atg Met	gly aaa	ctg Leu	gag Glu	aac Asn 220	agc Ser	agc Ser	gag Glu	ctc Leu	672
	Thr	atc Ile														720
act Thr	tct Ser	gtc Val	ttc Phe	ttg Leu 245	gac Asp	aat Asn	gcc Ala	agc Ser	ggc Gly 250	tcc Ser	tcc Ser	ctc Leu	acg Thr	gta Val 255	cgc Arg	768
		tct Ser										tga				807
<21	0>	53														
<21	1>	268														
<21	2>	PRT														
<21	3>	Mus (musci	ulus												
<40	0>	53														
Met 1	Leu	Phe	Val	Lys 5	Gln	Ser	Asp	ГÀЗ	Gly 10	Ile	Asn	Ser	Lys	Arg 15	Arg	
Ser	Lys	Ala	Arg 20	Arg	Leu	Lys	Leu	Gly 25	Leu	Pro	Gly	Pro	Pro 30	Gly	Pro	
Pro	Gly	Pro 35	Gln	Gly	Pro	Pro	Gly 40	Pro	Phe	Ile	Pro	Ser 45	Glu	Val	Leu	
Leu	Lys 50	Glu	Phe	Gln	Leu	Leu 55	Leu	Lys	Gly	Ala	Val 60	Arg	Gln	Arg	Glu	
Ser 65	His	Leu	Glu	His	Cys 70	Thr	Arg	Asp	Leu	Thr 75	Thr	Pro	Ala	Ser	Gly 80	
Ser			_		_		77-	01 =	<i>a</i> 1.,	Ton	n an	Com	G] n	λen	D~o	

Gly Ala Leu Leu Ala Leu Leu Ala Ala Thr Leu Ala Gln Gly Pro Arg 100 105 Ala Pro Arg Val Glu Ala Ala Phe His Cys Arg Leu Arg Arg Asp Val 115 . 120 Gln Val Asp Arg Arg Ala Leu His Glu Leu Gly Ile Tyr Tyr Leu Pro 130 135 Glu Val Glu Gly Ala Phe His Arg Gly Pro Gly Leu Asn Leu Thr Ser 150 Gly Gln Tyr Thr Ala Pro Val Ala Gly Phe Tyr Ala Leu Ala Ala Thr Leu His Val Ala Leu Thr Glu Gln Pro Arg Lys Gly Pro Thr Arg Pro 185 Arg Asp Arg Leu Arg Leu Leu Ile Cys Ile Gln Ser Leu Cys Gln His 200 Asn Ala Ser Leu Glu Thr Val Met Gly Leu Glu Asn Ser Ser Glu Leu 210 . 215 Phe Thr Ile Ser Val Asn Gly Val Leu Tyr Leu Gln Ala Gly His Tyr Thr Ser Val Phe Leu Asp Asn Ala Ser Gly Ser Ser Leu Thr Val Arg 250 245

Ser Gly Ser His Phe Ser Ala Ile Leu Leu Gly Leu

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
☑ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

U OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.